

# Chapter 1

---

# Microorganisms in Fermented Foods and Beverages

---

Jyoti Prakash Tamang, Namrata Thapa,  
Buddhiman Tamang, Arun Rai, and Rajen Chettri

## Contents

1.1	Introduction.....	2
1.1.1	History of Fermented Foods .....	3
1.1.2	History of Alcoholic Drinks .....	4
1.2	Protocol for Studying Fermented Foods .....	5
1.3	Microorganisms .....	6
1.3.1	Isolation by Culture-Dependent and Culture-Independent Methods.....	8
1.3.2	Identification: Phenotypic and Biochemical.....	8
1.3.3	Identification: Genotypic or Molecular.....	9
1.4	Main Types of Microorganisms in Global Food Fermentation.....	10
1.4.1	Bacteria.....	10
1.4.1.1	Lactic Acid Bacteria .....	11
1.4.1.2	Non-Lactic Acid Bacteria.....	11
1.4.2	Yeasts .....	11
1.4.3	Fungi .....	12
1.4.4	Pathogenic Contaminants.....	12
1.4.5	Gut Microflora.....	12
1.5	Types of Fermented Foods.....	13
1.5.1	Fermented Milks .....	13
1.5.2	Fermented Cereal Foods .....	20
1.5.3	Fermented Vegetable Products .....	29
1.5.4	Fermented Legumes.....	36
1.5.5	Fermented Root Crop and Tuber Products .....	48
1.5.6	Fermented Fruit Products .....	51
1.5.7	Fermented Meat Products.....	51

## 2 ■ Health Benefits of Fermented Foods and Beverages

1.5.8	Fermented, Dried and Smoked Fish Products.....	51
1.5.9	Vinegar.....	64
1.5.10	Ethnic Fermented Tea.....	64
1.5.11	Bacterial Cellulose.....	64
1.5.12	Cocoa/Chocolates.....	68
1.5.13	Coffee Cherries.....	68
1.5.14	Fermented Eggs.....	68
1.6	Types of Fermented Beverages.....	68
1.6.1	Amyolytic Mixed Starters.....	68
1.6.2	Alcoholic Beverages and Drinks.....	84
1.6.3	Nondistilled and Unfiltered Alcoholic Beverages Produced by Amyolytic Starters.....	84
1.6.4	Nondistilled and Filtered Alcoholic Beverages Produced by Amyolytic Starters.....	85
1.6.5	Distilled Alcoholic Beverages Produced by Amyolytic Starters.....	85
1.6.6	Alcoholic Beverages Produced by Human Saliva.....	85
1.6.7	Alcoholic Beverages Produced by Mono Fermentation.....	85
1.6.8	Alcoholic Beverage Produced from Honey.....	86
1.6.9	Alcoholic Beverages Produced from Plants.....	86
1.6.10	Alcoholic Beverages Produced by Malting or Germination.....	86
1.6.11	Alcoholic Beverages Produced from Fruits without Distillation.....	86
1.6.12	Distilled Alcoholic Beverages without Amyolytic Starters.....	87
1.6.13	Recommendations.....	87
1.7	Conclusion.....	87
	References.....	87

### 1.1 Introduction

Traditionally boiled rice is the staple diet with ethnic fermented and nonfermented legume products, vegetables, pickles, fish and meat on the side in the Far East, South and North Asia and the Indian subcontinent excluding western and northern India. In the west and north of India, wheat/barley-based breads/loaves is the staple diet together with milk and fermented milk products, meat and fermented meats. This diet is also followed in West Asia, Europe, North America and in Australia and New Zealand. Sorghum/maize porridges are the main diet with ethnic fermented and nonfermented sorghum/maize/millet, cassava, wild legume seeds, meat and milk products in Africa and South America (Tamang 2012a). Fermented foods are popular throughout the world and in some regions make a significant contribution to the diet of millions of individuals. The fermented food products supply protein, minerals, and other nutrients that add variety and nutritional fortification to otherwise starchy, bland diets.

Fermentation was traditionally a process which enabled the preservation of perishable food and has been used for centuries ([www.eolss.net/sample-chapters/c06/e6-34-09-09.pdf](http://www.eolss.net/sample-chapters/c06/e6-34-09-09.pdf); Hansen 2004). The term fermentation comes from the Latin word *fermentum* (to ferment). Throughout the world there are many different types of fermented foods, in which a range of different substrates are metabolized by a variety of microorganisms to yield products with unique and appealing characteristics. Fermented foods and alcoholic beverages are produced from raw materials or substrates of plant or animal origins mostly by natural fermentation or in the case of a few products by

black-slopping or the addition of a traditionally prepared starter culture(s) containing functional microorganisms which modify the substrates biochemically and organoleptically into edible products that are socially acceptable to consumers (Tamang 2010b).

### 1.1.1 History of Fermented Foods

Methods for fermentation of vegetables might have developed in Asia (Pederson 1979), or in the Mediterranean (Hulse 2004), or in Europe (Tamang and Samuel 2010). Methods for pickling vegetables were well established during the Song dynasty in China (960–1279 AD), and have remained more or less the same to the present day (Tamang and Samuel 2010). *Suau cai*, the ethnic fermented mixed vegetable product of China, was one of the main meals for workers during the construction of the Great Wall of China in 300 BC (Pederson 1979). The history of *kimchi*, a fermented vegetable product of Korea, was traced back to 3–4 AD (Chang 1975). The traditional preparation of *kimchi* was mentioned in some Korean historical documents recorded during 1759–1829 AD (Cheigh 2002). *Sauerkraut* or *sauerkohl* meaning sour cabbage in German was documented in the seventeenth century (Pederson and Albury 1969). Olives were preserved or fermented with various methods in Roman times during 50–150 AD (Sealey and Tyers 1989). The oral history of the origin of some common Himalayan fermented vegetables such as *gundruk*, ethnic fermented leafy vegetables and *sinki*, ethnic fermented radish tap roots were documented by Tamang (2010a).

Bread-making is one of the oldest food processing practices in the food history of human beings. There are ample archaeological remains of tools and installations which were used to make bread in ancient Egypt (Samuel 2002), and descriptions of baking in Roman texts and tombs are available (Curtis 2001). About 250 bakeries were reported to have operated in ancient Rome around 100 BC (Pederson 1979). *Dosa*, the ethnic fermented pancake made from rice and black gram in South India was first noted in the Tamil Sangam literature in India about the sixth century AD (Srinivasa 1930). The traditional preparation of *idli*, an ethnic fermented rice-black gram food of India and Sri Lanka, eaten at breakfast, has been described by the poet Chavundaraya of South India in 1025 AD (Iyengar 1950). *Dhokla*, a fermented mixture of wheat and Bengal gram of western India was first mentioned in 1066 AD (Prajapati and Nair 2003). *Jalebi*, the fermented cereal-based pretzel-like product of India and Pakistan has been known since 1450 AD and is probably of Arabic or Persian origin (Gode 1943).

Soybean was probably introduced to India from China through the Himalayas several centuries ago and some believe that soybeans were also brought via Myanmar by traders from Indonesia (Shurtleff and Aoyagi 2010). *Kinema*, a fermented sticky soybean food of India, Nepal and Bhutan might have originated in east Nepal around 600 BC to 100 AD during the Kirat dynasty (Tamang 2010a). The word *kinema* was derived from the word *kinamba* of the Limboo language of the Kirat race, *ki* means fermented and *namba* means flavor (Tamang 2001). A hypothetical triangle “*natto*-triangle” was initially proposed by Nakao (1972) and was based on the distribution of plasmids (Hara et al. 1986, 1995) and the 16s RNA sequencing (Tamang et al. 2002) of the *Bacillus* species from common nonsalted sticky fermented soybean foods of Asia. An imaginary triangle known as “*Kinema-Natto-Thua nao* triangle” (KNT-triangle) was proposed by Tamang (2010a). Within the KNT-triangle in Asia only *Bacillus*-fermented soybean foods are prepared and consumed starting from Japan (*natto*), touching the Korean peninsula (*chungkok-jang*), South China (*douchi*), North Thailand (*thua-nao*), Myanmar (*pepok*), Cambodia and Laos (*sieng*), southern Bhutan, the Darjeeling hills and Sikkim in India, and eastern Nepal (*kinema*), and the North East Indian states of Meghalaya (*tungrymbai*), Manipur (*hawaijar*), Mizoram

(*bekang*), Nagaland (*aakhone*), and Arunachal Pradesh (*perayaan*). The proposed “KNT triangle” does not include nonsticky and nonbacilli fermented soybean products such as *tempeh*, *miso*, *sufu*, *shoyu*, and so on.

Asian fermented soybean foods might have originated from *douchi* or *tau-shi*, during the Han dynasty in southern China around 206 BC (Bo 1984a,b, Zhang and Liu 2000). *Natto*, a fermented sticky soybean, was introduced to Japan from China by Buddhist priests during the Nara period around 710–794 AD (Itoh et al. 1996, Kiuchi 2001). Production of *shoyu* and *miso* in China was recorded around 1000 BC with the transfer of knowledge of production of *shoyu* and *miso* to Japan in around 600 AD (Yokotsuka 1985). *Tempeh*, the mold-fermented soybean food of Indonesia was originally introduced by ethnic Chinese traders in the early seventeenth century and the earliest record of the word *tempeh* appeared in the Serat Centini manuscript around 1815 AD (Astuli 1999).

The oldest sacred books of the Hindus, the *Rig Veda* and the *Upanishads* mentioned the origin of *dahi*, one of the oldest yoghurt-like fermented milk products of India during 6000–4000 BC (Yegna Narayan Aiyar 1953). Some milk products of Sudan and Egypt consumed in modern Africa such as *rob* (made from cow/goat/sheep milk), *gariss* (from camel milk), *biruni* (cow/camel milk) of Sudan, and *mishb* (cow/camel milk) were mentioned by medieval Arab travelers (Odufa 1988, Dirar 1993). The ancient Turkish people in Asia who lived as nomads were the first to make yoghurt, called “yoghurut” (Rasic and Kurmann 1978). The Babylonian records refer to cheese in 2000 BC (Davis 1964). The importance of cheese in the food habits of people in Greece (1500 BC) and Rome (750 BC) has been well documented (Scott 1986).

The Mekong basin of south-west China, Laos and northern and north-west Thailand were the most probable place of origin of fermented fish products in Asia (Ishige 1993, Ruddle 1993). Consumption of sausage by the ancient Babylonians was recorded around 1500 BC (Pederson 1979). The name *salami* is believed to have originated from the city of Salamis located on the east coast of Cyprus, which was destroyed in 449 BC (Lücke 1985).

### 1.1.2 History of Alcoholic Drinks

Fermented beverages appeared in 5000 BC in Babylon, 3150 BC in Ancient Egypt, 2000 BC in Mexico and 1500 BC in Sudan ([www.eols.net/sample-chapters/c06/e6-34-09-09.pdf](http://www.eols.net/sample-chapters/c06/e6-34-09-09.pdf)). The earliest evidence of the grape in Egypt is seeds in jars imported from the Levant, dating to about 3150 BC indicating that wine was possibly produced in Egypt itself by 3000 BC (Murray 2000). Archaeological findings and chemical analyses of residues recovered from the Neolithic (6th millennium BC) Hajji Firuz Tepe, and the Early Bronze Age (4th millennium BC) Godin Tepe, both in western Iran, is commonly reported to represent the earliest evidence of wine making (Renfrew 1999, Wilson 1999).

Early Mesopotamian beer was based on barley malt, that is, sprouted and dried grain (Curtis 2001). Analysis of ancient beer residues using scanning electron microscopy was more focused on archaeological evidence (Samuel 1996, 2000). The whole starch remaining in the unheated malt would have broken down more slowly and the source of fermentation is uncertain (Samuel 2000). Archaeobotanical evidence from northern Germany and Scandinavia shows that hops and sweet gale (*Myrica gale*) became important beer flavorings in early medieval times (ninth to tenth century AD) (Behre 1984).

*Pulque*, one of the ancient alcoholic beverages of South America, which is fermented from *agave* juice and now is the national drink of Mexico was inherited from the Aztecs (Goncalves de Lima 1975). Another ancient alcoholic drink of the Andes Indians living in the lower altitude

regions of South America is *chicha*, prepared from maize through human saliva which serves as the source of amylase for conversion of starch into fermentation sugars (Escobar 1977).

During the *Vedic* period of Indian history (2500–200 BC), based originally around the Indus River system, alcoholic drink *soma rus* was common and was worshipped as the liquid god *Soma* because of its medicinal attributes (Bose 1922, Sarma 1939). Drinking of alcohol in India has been mentioned in the *Ramayana* during 300–75 BC (Prakash 1961). The malting process as well as wine fermentation is rarely used in traditional fermentation processes in Asia, instead, amylolytic mixed starters prepared from the growth of molds and yeasts on raw or cooked cereals are more commonly used (Tamang 2010c). The use of traditionally prepared amylolytic mixed starters, common to the Himalayas and South East Asia, might have its origins during the time of Euhok, the daughter of the legendary king of Woo of China, known as the goddess of rice-wine in Chinese culture in 4000 BC (Lee 1984). The first documentation of *chu*, the mixed amylolytic starter of China (use for production of fermented beverages and alcoholic drinks), was found in the Shu-Ching document written during the Chou dynasty (1121–256 BC) (Haard et al. 1999). The use of *chu* for fermentation of rice-based alcoholic beverages and drinks was recorded in the beginning of the Three Nations' Periods in Korea during first century BC to second century AD (Lee 1995). The word *ragi*, an amylolytic starter of Indonesia, was first noted on an ancient inscription called the Kembang Arum near Yogyakarta in Java around 903 AD (Astuli 1999). *Kodo ko jaanr*, the ethnic fermented finger millet alcoholic beverage of the Himalayas was mentioned in the history of Nepal during the Kirat dynasty in 625 BC to 100 AD (Adhikari and Ghimirey 2000). The Newar community of Nepal used to ferment alcoholic beverages from rice during the Malla dynasty in 880 AD (Khatri 1987). There are brief descriptions of the fermented millet beverages of the Darjeeling hills and Sikkim in India in historical documents (Hooker 1854, Risley 1928, Gorer 1938).

## 1.2 Protocol for Studying Fermented Foods

Protocols for studies of ethnic fermented foods and alcoholic beverages primarily focus on the following parameters (Tamang 2014):

1. Documentation of the indigenous knowledge of the local people of traditional preparation, culinary practices, mode of consumption, ethnical values, therapeutic uses, socio-economy, market survey of marginal producers of fermented foods/alcoholic beverages using standard format; photography.
2. Sensory character(s) of a fermented food or alcoholic beverage by a researcher: taste, texture, aroma/flower, and appearance.
3. Calculation of per capita consumption and annual production of ethnic fermented foods and alcoholic beverages of the particular region/village.
4. Determination of pH, temperature of the product *in situ*.
5. Collection of samples aseptically in pre-sterile containers.
6. Microbiology of fermented foods: determination of microbial loads of microorganisms (bacteria, yeasts, molds) and pathogenic contaminants (colony forming unit per gram or liter of sample), isolation of cultivable microorganisms.
7. Identification of isolates: phenotypic (morphological, physiological, and biochemical tests) and molecular identifications, assigning the proper identification of functional microorganisms following the standard norm of ICBN for microorganisms and well-authenticated taxonomical keys.

8. Isolation of uncultured microorganisms directly from food samples, and identification by culture-independent molecular techniques.
9. Preservation of identified strains of microorganisms in 15% glycerol at 80°C, or lyophilized, and deposited at authorized culture collection centers.
10. Experiments on fermentation dynamics or microbial changes during *in situ* fermentation to understand the role of functional or nonfunctional microorganism(s) using culture-independent techniques during natural fermentation.
11. Determination of the proximate composition and nutritional values of the products.
12. Determination of functional properties: probiotics, antioxidants, antimicrobial activities, tyrosinase inhibition, cell proliferation and MMP-2 inhibition activities, degradation of antinutritive compounds, bioactive compounds, unsaturated fatty acids, including omega 3-fatty acids, isoflavones and saponins, total phenolic compounds, total anthocyanin, saponin, and so on.
13. Studies on food safety: occurrence of pathogenic and spoilage microorganisms, toxin production, shelf-life of the products.
14. Optimization of a traditional process using a starter culture(s), consisting either of a pure culture or of a consortium of identified native microorganism(s) with desirable functional properties.
15. Organoleptic evaluation and consumer preference trials of the product prepared on laboratory scale.

### 1.3 Microorganisms

Microorganisms determine the characteristics of fermented food, for example, acidity, flavor and texture, as well as the health benefits that go beyond simple nutrition (Vogel et al. 2011). Microorganisms may be present as the indigenous microbiota of the food or as a result of the intentional addition of microorganisms as starter cultures in an industrial food fermentation process (Stevens and Nabors 2009). Also, microbial cultures can be used to produce several compounds (enzymes, flavors, fragrances, etc.) either specifically for application as food additives or *in situ* as a part of food fermentation processes (Longo and Sanromán 2006).

With an estimated 5000 varieties of fermented foods and beverages, worldwide, only a small fraction of these artisanal products have been subjected to scientific studies so far (Tamang 2010b). In many of these foods, the biological and microbiological bases of the fermentation processes are poorly understood. What little information is available often deals with the identification and perhaps preliminary characterization of the primary microbiota in the finished product. This in turn will necessitate a more thorough understanding of the microorganisms involved, in terms of the types and their specific activities, so that the fermentation process can be made more reliable and predictable.

With the discovery of microorganisms, it became possible to understand and manage food fermentation. Methods for isolating and purifying microbial cultures became available in the nineteenth century. Sterilization or pasteurization of the raw materials prior to inoculation with well-defined cultures allowed the fermentation processes to be managed with little variation. The use of defined cultures became the industrial standard in breweries by the nineteenth century. During the twentieth century, the wine, dairy, and meat industries also shifted production procedures toward the use of well-characterized and defined starter cultures. In the beginning, starter cultures were isolates from earlier fermentations that were maintained and propagated at the site



of production. The application of microbiology and process technology resulted in large improvements in the quality of the fermented food products. The quality improvements have been so great that today all significant production of fermented food is industrial. The small amount of “home fermentation” conducted in the form of baking, home brewing and private cheese making usually rely on commercially available yeast and bacterial cultures. The maintenance of the microorganisms differs between the different food industries in the sense that some fermentation industries, such as breweries and vinegar producers maintain their own stains and inocula. In the dairy industry, as well as in the meat industry and bakeries, cultures are usually obtained from suppliers dedicated to the production of high-quality food ingredients (Mogensen et al. 2002, Hansen 2004).

Fermentation can basically be performed either by natural or spontaneous fermentation, by back-slopping or by the addition of starter cultures. By spontaneous fermentation, the raw material and its initial treatment will encourage the growth of an indigenous microbiota. In most spontaneous fermentation, a microbial succession takes place: quite often lactic acid bacteria (LAB) will initially dominate followed by various species of yeasts. Molds only grow aerobically, limiting their occurrence in certain types of fermented products. LAB produce lactic acid and other antimicrobial substances that inhibit the growth of harmful bacteria along with reducing the sugar content, thereby prolonging the shelf life of the product. Yeasts mostly produce aroma components and alcohols. When molds are involved in fermentation, they generally contribute by producing both intra- and extracellular proteolytic and lipolytic enzymes that highly influence the flavor and texture of the product (Tamang and Fleet 2009).

In back-slopping, a part of a precious batch of a fermented product is used to inoculate the new batch. This procedure produces a higher initial number of beneficial microorganisms than that found in the raw material and ensures a faster and more reliable fermentation than that which occurs in spontaneous fermentation. Examples of back-slopping are home-made fermentation of milk, vegetables, and cereals (Josephsen and Jespersen 2004).

Novel insights into the metabolism of LAB offer perspectives for the application of a new generation of starter cultures. Functional starter cultures are starters that possess at least one inherent functional property. These can contribute to food safety and/or offer one or more organoleptic, technological, nutritional, or health advantage. The implementation of carefully selected strains as starter cultures or co-cultures in fermentation processes can help to achieve *in situ* expression of the desired property, maintaining a perfectly natural and healthy product. A functional LAB starter culture is able to produce antimicrobial substances, sugar polymers, sweeteners, aromatic compounds, useful enzymes, or nutraceuticals, while the so-called probiotic strains, mainly LAB, exhibit health-promoting properties. Such functionalities also lead to a wider application area and higher flexibility of starter cultures (Leroy and De Vuyst 2004). Nowadays, there are many commercial suppliers of starter cultures worldwide, such as Alce and CSL (Italy), ASCRS (Australia), Chr. Hansen and Danisco (Denmark), CSK and DSM (the Netherlands), Degussa and Gewürzmüller (Germany), Lallemand (Canada), NZDRI (New Zealand), Quest International (UK), Rhodia (France), and so on. All companies can easily be found on the Internet/World wide Web (Hansen 2002, 2004).

Fermented foods are the hubs of consortia of microbiota and mycobiota (functional, nonfunctional, and pathogenic contaminants), which may be present as natural indigenous microbiota in uncooked plant or animal substrates, utensils, containers, earthen pots, or environments (Hesseltine 1983, Tamang 1998), or as a result of the intentional addition of the microorganisms as starter cultures in an industrial food fermentation process (Stevens and Nabors 2009). Functional microorganisms transform the chemical constituents of raw

materials of plant/animal sources during fermentation thereby enhancing the nutritional value of the products, enriching them with improved flavor and texture, prolonging their shelf life, and fortifying them with health-promoting bio-active compounds (Farhad et al. 2010). Species of lactic acid bacteria and *Bacillus*, and amylolytic and alcohol-producing yeasts and filamentous molds are the major microbiota in the fermented foods and alcoholic beverages of Asia, whereas LAB or a combination of bacteria (LAB, non-LAB, micrococci)-yeast mixtures and filamentous molds are more prominent in Africa. Filamentous molds and bacilli are rare in the fermented foods and beverages of Africa, Europe, Australia, and America (Tamang 1998), although fermented legume products, based on *Bacillus* fermentation, are common in West Africa (Oguntoyinbo et al. 2007).

### 1.3.1 Isolation by Culture-Dependent and Culture-Independent Methods

The classical phenotypic identification methods are totally based on culture-dependent techniques for only cultivable microorganisms in culture media, ignoring several unknown uncultivable microorganisms that may have major or minor functional roles in the production of fermented foods. Direct DNA extraction from samples of fermented foods commonly referred to as culture-independent methods are nowadays commonly used in food microbiology to profile both cultivable and uncultivable microbial populations from fermented foods (Cocolin and Ercolini 2008). Culture independent techniques first appeared in food microbiology at the end of the 1990s and since then they have been applied extensively. These methods do not rely on cultivation and target nucleic acids (DNA and RNA) to identify and follow the changes that occur in the main populations present in a specific ecosystem (Cocolin et al. 2013). The most popular culture-independent technique being used in the isolation of microorganisms from fermented foods is a PCR-DGGE analysis to profile bacterial populations (Cocolin et al. 2011, Tamang 2014) and yeast populations in fermented foods (Cocolin et al. 2002, Jianzhong et al. 2009). Culture-dependent and culture-independent methods are contradictory to each other (Alegría et al. 2011), but for microbial taxonomy both methods are equally important and complementary. Both cultivable and uncultivable microorganisms from any fermented food and beverage may be identified using culture-dependent and -independent methods to document a complete profile of native functional and nonfunctional microorganisms, and to study both inter- and intra-species diversity within a particular genus or among genera (Ramos et al. 2010, Yan et al. 2013, Tamang 2014). Greppi et al. (2013a) first reported the combination of both culture-dependent and independent methods to reveal predominant yeast species and biotypes in the traditional fermented maize foods of Benin. The DGGE analysis on the DNA directly extracted from fermented maize products demonstrated the presence of *Dekkera bruxellensis* and *Debaryomyces hansenii*, not detected by the culture-based approach (Greppi et al. 2013b).

### 1.3.2 Identification: Phenotypic and Biochemical

Phenotypic characteristics include colony appearance, cell morphology, Gram staining, growth at different temperatures (8–65°C), pH (3.9–9.6), and salt tolerance (4.0%–18%). Biochemical tests are based on the metabolic activities of microorganisms such as carbon and nitrogen sources, energy sources, sugar fermentation, secondary metabolite formation, and enzyme production (Tamang 2014). A few ready-to-use commercial identification kits are commonly used such as API 50CH (bioMérieux, France) for rapid sugar fermentation tests, and the Biolog Microbial Identification System, for identification of different groups of bacteria and yeasts.



### 1.3.3 Identification: Genotypic or Molecular

Molecular identification is emerging as an accurate and reliable identification tool, and is widely used in identification of both culture-dependent and culture-independent microorganisms from fermented foods. Owing to a variety of tools that provide advanced molecular differentiation of microorganisms, microbial populations can be quantified and new microbial species isolated and identified (Giraffa and Carminati 2008). Some common molecular tools used in identification of microorganisms isolated from fermented foods are species-specific PCR, qPCR, rep-PCR, AFLP, RAPD, DGGE, TGGE, ARDRA, mtDNA-RFLP, mCOLD-PCR, MLSA, and MLST.

Species-specific PCR primers are used to identify a particular species of the genus (Tamang et al. 2005); this technique is widely applied in the identification of LAB isolated from fermented foods (Robert et al. 2009). The application of real-time quantitative PCR (qPCR) with specific primers provides the specific detection and quantification of LAB species in fermented foods (Park et al. 2009). The repetitive extragenic palindromic sequence-based PCR (rep-PCR) technique permits typing at a subspecies level and reveals significant genotypic differences between strains of the same bacterial species from fermented foods (Tamang et al. 2008). Amplified fragment-length polymorphism (AFLP) is a technique based on the selective amplification and separation of genomic restriction fragments, and its applicability is for the identification of various LAB strains in fermented foods (Tanigawa and Watanabe 2011). Random amplification of polymorphic DNA (RAPD) is commonly used for discrimination of LAB strains from fermented foods (Schillinger et al. 2003, Chao et al. 2008). The amplified ribosomal DNA restriction analysis (ARDRA) technique using restriction enzymes is also useful in identification of uncultivable microorganisms from fermented foods (Jeyaram et al. 2010). The mt DNA-RFLP technique showed discriminating power similar to microsatellite typing and interdelta analysis (Schuller et al. 2007) and is considered as a useful genetic marker for *Saccharomyces cerevisiae* in amylolytic starters (Jeyaram et al. 2011). Techniques of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) have been developed to profile microbial communities directly from fermented foods, and are based on sequence-specific distinctions of 16S rDNA or 26S rDNA amplicons produced by PCR (Ongol and Asano 2009, Alegría et al. 2011). A modified CO-amplification at lower denaturation temperature PCR (mCOLD-PCR) method has been developed to detect low-abundant microorganisms using a double-strand RNA probe to inhibit the amplification of the sequence of a major microorganism in wine fermentation (Takahashi et al. 2014). Multilocus sequence analysis (MLSA) using housekeeping genes as molecular markers alternative to the 16S rRNA genes, has been proposed for LAB species identification from fermented foods (Naser et al. 2005, de Bruyne et al. 2007, 2008, 2010). Multilocus sequence typing (MLST) has also been used for discriminating LAB strains from fermented foods (Diancourt et al. 2007, Picozzi et al. 2010, Tanigawa and Watanabe 2011).

Effective tools of next-generation sequencing (NGS) such as phylobiomics, metagenomics, and metatranscriptomics are required for the documentation of cultures in traditional fermented products, for sensory quality and safety improvements, in some cases for starter culture design for commercialization and potentially for supporting sustainable food systems (van Hijum et al. 2013). A proteomics identification method using matrix-assisted laser desorption ionizing-time of flight mass spectrometry (MALDI-TOF MS) is used to identify the species of *Bacillus* in the fermented foods of Africa (Savadogo et al. 2011), *Lactobacillus* strains isolated from fermented foods (Dušková et al. 2012, Sato et al. 2012), sub-speciation of *Lactococcus (Lc.) lactis* (Tanigawa et al. 2010), LAB from traditional fermented vegetables of Vietnam (Nguyen et al. 2013b), probiotics from yoghurt (Angelakis et al. 2011), and *Tetragenococcus halophilus* and *Tetra. muriaticus* from fermented foods

(Kuda et al. 2014). The application of NGS such as metagenomic approaches using massively parallel pyrosequencing of tagged 16S rRNA gene amplicons provide detailed information on microbial communities associated with *kimchi* (Jung et al. 2011, Park et al. 2012), *nukadoko*, a fermented rice bran mash used for pickling vegetables in Japan (Sakamoto et al. 2011), *narezushi*, a fermented salted fish with cooked rice in Japan (Kiyohara et al. 2012), and *ben-saalga*, a traditional gruel of pearl millet in Burkina Faso (Humblot and Guyot 2009). The 16S rRNA gene sequence-based pyrosequencing method enables a detailed, comprehensive and high-throughput analysis of microbial ecology (Sakamoto et al. 2011), and this method has been applied to various traditional fermented foods (Oki et al. 2014). Compared to the pyrosequencing analysis, the DGGE method only revealed some of the major bacterial species such as *Bacillus thermoamylovorans* and *B. licheniformis* in *chungkokjang* (the sticky fermented soybean food of Korea) and could not detect a large number of predominant or diverse rare bacterial species identified in the pyrosequencing analysis. Also the regional differences of the bacterial community were more clearly represented in the pyrosequencing method than in the DGGE analysis (Nam et al. 2011).

Metabolomics, also called metabonomics or metabolic profiling, deals with the simultaneous determination and quantitative analysis of intracellular metabolites or low-molecular-mass molecules and can be used as a tool for the comprehensive understanding of fermented and functional foods with LAB (Mozzi et al. 2013). The application of species-independent functional gene microarray for identification of LAB in fermented foods has been developed (Weckx et al. 2009). Up-to-date analytical methods have also been applied in the identification and discrimination of some microbial species/strains from fermented foods such as length heterogeneity PCR (LH-PCR), high-throughput sequencing (HTS), BOX-PCR (Zhu et al. 2014), and so on.

## 1.4 Main Types of Microorganisms in Global Food Fermentation

Main types (with genera) of microorganisms associated with global fermented foods and beverages are grouped as follows (Bernardeau et al. 2006, Tamang and Fleet 2009, Tamang 2010a,b,c, Bourdichon et al. 2012, Alexandraki et al. 2013):

- Bacteria: *Acetobacter*, *Arthrobacter*, *Bacillus*, *Bifidobacterium*, *Brachybacterium*, *Brevibacterium*, *Carnobacterium*, *Corynebacterium*, *Enterobacter*, *Enterococcus*, *Gluconacetobacter*, *Hafnia*, *Halomonas*, *Klebsiella*, *Kocuria*, *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Macroccoccus*, *Microbacterium*, *Micrococcus*, *Oenococcus*, *Pediococcus*, *Propionibacterium*, *Staphylococcus*, *Streptococcus*, *Streptomyces*, *Tetragenococcus*, *Weisella*, *Zymomonas*.
- Fungi: *Actinomuor*, *Aspergillus*, *Fusarium*, *Lecanicillium*, *Mucor*, *Neurospora*, *Penicillium*, *Rhizopus*, *Scopulariopsis*, *Sperendonema*.
- Yeasts: *Candida*, *Cyberlindnera*, *Cystofilobasidium*, *Debaryomyces*, *Dekkera*, *Hanseniaspora*, *Kazachstania*, *Galactomyces*, *Geotrichum*, *Guehomuces*, *Kluveromyces*, *Lachancea*, *Metschnikowia*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, *Schwanniomyces*, *Starmerella*, *Torulaspora*, *Trigonopsis*, *Wickerhamomyces*, *Yarrowia*, *Zygosaccharomyces*, *Zygorulaspora*.

### 1.4.1 Bacteria

Bacteria are the most dominant microorganisms in both naturally fermented foods or foods fermented by the use of starter cultures. Among the bacteria, lactic acid bacteria are commonly

associated with acidic fermented foods, while non-LAB bacteria such as *Bacillus*, micrococcaceae, *Bifidobacterium*, *Brachybacterium*, *Brevibacterium*, and *Propionibacterium* etc., are also involved in food fermentation, frequently as minor or secondary groups.

#### 1.4.1.1 Lactic Acid Bacteria

Lactic acid bacteria are Gram-positive, catalase-negative bacteria that produce large amounts of lactic acid. The bacterial groups that make up the LAB are among the most familiar to humans, because of their association with the human environment, and with a wide range of naturally fermented dairy products, grain crops, vegetables, and so on. The LAB comprise a large bacterial group consisting of about 380 species in 40 genera of 6 families, belonging phylogenetically to the order *Lactobacillales* within the phylum *Firmicutes* (Stiles and Holzapfel 1997). Common genera of the LAB isolated from various fermented foods of the world are *Alkalibacterium*, *Carnobacterium*, *Enterococcus*, *Lactococcus*, *Lactobacillus*, *Leuconostoc*, *Oenococcus*, *Pediococcus*, *Streptococcus*, *Tetragenococcus*, *Vagococcus*, and *Weissella* (Carr et al. 2002, Salminen et al. 2004, MetaMicrobe.com/ Lactic Acid Bacteria 2013).

#### 1.4.1.2 Non-Lactic Acid Bacteria

*Bacillus* is reported from the alkaline-fermented foods of Asia and Africa (Parkouda et al. 2009). Species of *Bacillus* present in fermented foods mostly soybean-based foods are *B. amyloliquefaciens*, *B. circulans*, *B. coagulans*, *B. firmus*, *B. licheniformis*, *B. megaterium*, *B. pumilus*, *B. subtilis*, *B. subtilis* variety *natto* and *B. thuringiensis* (Kiers et al. 2000, Kubo et al. 2011), while strains of *B. cereus* have been isolated from the fermentation of *Prosopis africana* seeds for the production of *okpehe* in Nigeria (Oguntoyinbo et al. 2007). Some strains of *B. subtilis* produce  $\lambda$ -polyglutamic acid (PGA) which is an amino acid polymer commonly present in Asian fermented soybean foods giving the characteristic sticky texture to the product (Urushibata et al. 2002, Meerak et al. 2007, Nishito et al. 2010).

Species of *Bifidobacterium*, *Brachybacterium*, *Brevibacterium*, and *Propionibacterium* have been isolated from cheese and other fermented milks (Bourdichon et al. 2012). Several species of *Kocuria*, *Micrococcus*, and *Staphylococcus* have been reported from fermented milk products, fermented sausages and meat and fish products (Wu et al. 2000, Martín et al. 2006, Coton et al. 2010). *Enterobacter cloacae*, *Klebsiella pneumoniae*, *K. pneumoniae* subsp. *ozaenae*, *Haloanaerobium*, *Halobacterium*, *Halococcus*, *Propionibacterium*, *Pseudomonas*, and so on, are also present in many fermented foods (Tamang 2010b). Species of *Arthrobacter* and *Hafnia* are involved in meat fermentation (Bourdichon et al. 2012).

### 1.4.2 Yeasts

The role of yeasts in food fermentation is to ferment sugar, produce secondary metabolites, inhibit growth of mycotoxin-producing molds and display several enzymatic activities such as lipolytic, proteolytic, pectinolytic, glycosidasic and urease activities (Aidoo et al. 2006, Romano et al. 2006). Genera of yeasts reported from fermented foods, alcoholic beverages and nonfood mixed amylolytic starters are *Brettanomyces*, *Candida*, *Cryptococcus*, *Debaryomyces*, *Dekkera*, *Galactomyces*, *Geotrichum*, *Hansenula*, *Hanseniaspora*, *Hyphopichia*, *Issatchenkia*, *Kazachstania*, *Kluyveromyces*, *Metschnikowia*, *Pichia*, *Rhodotorula*, *Rhodospiridium*, *Saccharomyces*,

*Saccharomyces*, *Saccharomycopsis*, *Schizosaccharomyces*, *Sporobolomyces*, *Torulaspota*, *Torulopsis*, *Trichosporon*, *Yarrowia*, and *Zygosaccharomyces* (Watanabe et al. 2008, Tamang and Fleet 2009, Kurtzman et al. 2011, Lv et al. 2013).

### 1.4.3 Fungi

The major roles of fungi, mostly filamentous molds, in fermented foods and alcoholic beverages are the production of intra- and extracellular proteolytic and lipolytic enzymes that highly influence the flavor and texture of the product, and also the degradation of antinutritive factors improving bioavailability of minerals (Josephsen and Jespersen 2004, Aidoo and Nout 2010). Species of *Actinomucor*, *Amylomyces*, *Aspergillus*, *Monascus*, *Mucor*, *Neurospora*, *Penicillium*, *Rhizopus*, and *Ustilago* are reported from many fermented foods, Asian nonfood amylolytic starters and alcoholic beverages (Hesseltine 1991, Nout and Aidoo 2002).

### 1.4.4 Pathogenic Contaminants

About 80% of fermented foods are produced by natural fermentation and may contain functional, nonfunctional, and pathogenic microorganisms during initial fermentation. Pathogenic bacteria commonly reported for fermented foods are *Escherichia coli*, *Listeria monocytogenes*, *Yersinia enterocolitica*, *Bacillus cereus*, *Clostridium botulinum*, and so on (Lindqvist and Lindblad 2009, Rossi et al. 2011).

### 1.4.5 Gut Microflora

The human gastrointestinal tract (GIT) houses over  $10^{14}$  microbial cells with over 1000 diverse bacterial types, mostly in the colon (Lepage et al. 2012, Purchiaroni et al. 2013). Colonization of the gut is initiated before birth following ingestion of microbe-containing amniotic fluid by the fetus (Mshvildadze and Neu 2010, Aagaard et al. 2014). The majority of bacteria in the adult gut are nonsporing anaerobes, the most numerically predominant of which include species of *Bacteroides*, *Bifidobacterium*, *Eubacterium*, *Clostridium*, *Lactobacillus*, *Fusobacterium*, and various Gram-positive cocci and bacteria that are present in lower numbers include *Enterococcus* spp., Enterobacteriaceae, methanogens, and dissimilatory sulfate-reducing bacteria (Wallace et al. 2011). Microorganisms colonize different parts of GIT and bacterial population density varies along the GIT (Romano-Keeler et al. 2014). The GIT is one of the most complex ecosystems of microorganisms ranging from bacteria (mostly LAB), archaea (e.g., methanogens), and eukarya (fungi, helminthes, and protozoa) as well as viruses (Mitsuoka 1992, Holzappel et al. 1997, Norman et al. 2014). In healthy adults, 80% of the identified fecal microbiota belong to four dominant phyla: the Gram-negative *Bacteroides* and *Proteobacteria* and the Gram-positive *Actinobacteria* and *Firmicutes* which include at least 17 families, corresponding to no less than 1250 different species of bacteria (Schuijt et al. 2013). The composition and distribution of gut microbiota (Purchiaroni et al. 2013) are as follows: in the stomach (*Lb. reuteri*, *Lb. delbrueckii*, *Lb. gastricus*, *Lb. antri*), in the small intestine (*Lb. reuteri*, *Lb. bulgaricus*, *Lb. acidophilus*, *Enterococcus avium*, *Ent. dispar*, *Ent. durans*, *Ent. faecalis*, *Ent. faecium*, *Ent. flavescens*, *Ent. gallinarum*, *Ent. hirae*, *Ent. mundtii*, *Ent. raffinosus*), and in the large intestine (*Ent. faecalis*, *Bacteroides*, *Bifidobacterium*, *Eubacterium*, *Peptococcus*, *Clostridium*, *Lactobacillus*).

## 1.5 Types of Fermented Foods

The major groups of substrates-based fermented foods are as follows:

1. Fermented milk foods
2. Fermented cereal foods
3. Fermented vegetable foods
4. Fermented soybean and non-soybean foods
5. Fermented meat products
6. Fermented fish products
7. Fermented root/tuber products
8. Fermented beverages and Asian amylolytic starters
9. Miscellaneous fermented products (fermented tea, cocoa, vinegar, *nata*, *pidan*, etc.)

### 1.5.1 Fermented Milks

Fermented milks (Table 1.1) are classified into two major groups based on the presence of dominant microorganisms: (i) lactic fermentations which are dominated by species of LAB, and consist of the thermophilic type (e.g., yogurt, Bulgarian buttermilk), probiotic type (acidophilus milk, yakult, bifidus milk), and the mesophilic type (e.g., natural fermented milk, cultured milk, cultured cream, cultured buttermilk); and (ii) fungal-lactic fermentations where LAB and yeasts species cooperate to generate the final product and consist of alcoholic milks (e.g., *kefir*, *koumiss*, acidophilus-yeast milk), and moldy milks (e.g., *viili*) (Mayo et al. 2010). Starter cultures in milk fermentation are of two types depending on the principal function, primary cultures to participate in the acidification, and secondary cultures for flavor, aroma, and maturing activities (Topisirovic et al. 2006). The main species involved as primary cultures in milk fermentation are *Lactococcus lactis* subsp. *cremoris*, *Lc. lactis* subsp. *lactis*, *Lactobacillus delbrueckii* subsp. *delbrueckii*, *Lb. delbrueckii* subsp. *lactis*, *Lb. helveticus*, *Leuconostoc* spp., and *Streptococcus thermophilus* (Parente and Cogan 2004). Secondary cultures used in cheese making are *Brevibacterium linens*, *Propionibacterium freudenreichii*, *Debaryomyces hansenii*, *Geotrichum candidum*, *Penicillium camemberti*, and *P. roqueforti* for the development of flavor and texture during the ripening of cheese (Coppola et al. 2006, Quigley et al. 2011). Besides primary and secondary cultures, some non-starter lactic acid bacteria (NSLAB) microbiota are usually present in high numbers which include *Enterococcus durans*, *Ent. faecium*, *Lb. casei*, *Lb. plantarum*, *Lb. salivarius*, and *Staphylococcus* spp. (Briggiler-Marcó et al. 2007). Yogurt is a widely consumed highly nutritious fermented milk as a coagulated milk product resulting from the fermentation of milk by *Strep. thermophilus* and *Lb. delbrueckii* subsp. *bulgaricus* (formerly *Lb. bulgaricus*) (Tamime and Robinson 2007). *Lb. acidophilus*, *Lb. casei*, *Lb. rhamnosus*, *Lb. gasseri*, *Lb. johnsonii*, and *Bifidobacterium* spp., are among the most common adjunct cultures in yogurt fermentation (Guarner et al. 2005). Fermented milk products that are manufactured using starter cultures containing yeasts include acidophilus-yeast milk, *kefir*, *koumiss*, and *viili* (de Ramesh et al. 2006). *Lb. acidophilus*, *Lb. amylovorus*, *Lb. crispatus*, *Lb. gallinarum*, *Lb. gasseri*, and *Lb. johnsonii* are reported from acidophilus milk (Berger et al. 2007).

Natural fermented milks are one of the oldest methods of milk fermentation using raw or boiled milk to ferment spontaneously or by using the back-slopping method (Robinson and Tamime 2006). In back-slopping, a part of a precious batch of a fermented product is used to inoculate the new batch

**Table 1.1 Some Common and Uncommon Ethnic Fermented Milk Products of the World**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
Acidophilus milk	Cow milk	Acidic, sour, drink	Species of <i>Lactobacillus</i> , <i>Lactococcus</i>	Russia, East Europe, Greece, Turkey, North America, Scandinavia	Mayo et al. (2010)
<i>Airag</i>	Mare or camel milk	Acidic, sour, mild alcoholic, drink	<i>Lb. helveticus</i> , <i>Lb. kefiranofaciens</i> , <i>Bifidobacterium mongoliense</i> , <i>Kluyveromyces marxianus</i>	Mongolia	Watanabe et al. (2008, 2009b)
<i>Ayib</i>	Goat milk		<i>Canida</i> sp., <i>Saccharomyces</i> sp., <i>Lactobacillus</i> sp., <i>Leuconostoc</i> sp.	East and Central Africa	Odunfa and Oyewole (1997)
<i>Biruni</i>	Cow/camel milk	Acidic, semi-liquid, drink	LAB	Sudan	Jung (2012)
Butter	Animal milk	Soft paste, butter	LAB	Worldwide	Mayo et al. (2010)
<i>Buttermilk</i>		Acid fermented butter milk	<i>Lb. bulgaricus</i>	Bulgaria	Mayo et al. (2010)
<i>Cheese</i>	Animal milk	Soft or hard, solid; side dish, salad	<i>Lc. lactis</i> subsp. <i>cremoris</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lb. delbrueckii</i> subsp. <i>delbrueckii</i> , <i>Lb. delbrueckii</i> subsp. <i>lactis</i> , <i>Lb. helveticus</i> , <i>Lb. casei</i> , <i>Lb. plantarum</i> , <i>Lb. salivarius</i> , <i>Leuconostoc</i> spp., <i>Strep. thermophilus</i> , <i>Ent. durans</i> , <i>Ent. faecium</i> , and <i>Staphylococcus</i> spp., <i>Brevibacterium linens</i> , <i>Propionibacterium freudenreichii</i> , <i>Debaryomyces hansenii</i> , <i>Geotrichum candidum</i> , <i>Penicillium camemberti</i> , <i>P. roqueforti</i> .	Worldwide	Parente and Cogan (2004), Quigley et al. (2011)



<i>Chhu</i>	Yak/cow milk	Cheese like product, curry, soup	<i>Lb. farciminis</i> , <i>Lb. brevis</i> , <i>Lb. alimentarius</i> , <i>Lb. salivarius</i> , <i>Lact. lactis</i> , <i>Saccharomycopsis</i> sp., <i>Candida</i> sp.	India, Nepal, Bhutan, China (Tibet)	Dewan. and Tamang (2006)
<i>Chhurpi (hard)</i>	Yak/cow milk	Chewable milk, masticator	<i>Lb. farciminis</i> , <i>Lb. casei</i> , <i>Lb. biofermentans</i> , <i>W. confusus</i>	India, Nepal, Bhutan, China (Tibet)	Tamang (2010a)
<i>Chhurpi (soft)</i>	Yak/cow milk	Cheese-like product, soup, curry, pickle	<i>Lb. farciminis</i> , <i>Lb. paracasei</i> , <i>Lb. biofermentans</i> , <i>Lb. plantarum</i> , <i>Lb. curvatus</i> , <i>Lb. fermentum</i> , <i>Lb. alimentarius</i> , <i>Lb. kefir</i> , <i>Lb. hilgardii</i> , <i>W. confusus</i> , <i>Ent. faecium</i> , <i>Leuc. mesenteroides</i>	India, Nepal, Bhutan, China (Tibet)	Tamang et al. (2000)
<i>Dahi</i>	Cow/buffalo milk, starter culture	Curd, savory	<i>Lb. bif fermentans</i> , <i>Lb. alimentarius</i> , <i>Lb. paracasei</i> , <i>Lact. lactis</i> , <i>Strep. cremoris</i> , <i>Strep. lactis</i> , <i>Strep. thermophilus</i> , <i>Lb. bulgaricus</i> , <i>Lb. acidophilus</i> , <i>Lb. helveticus</i> , <i>Lb. cremoris</i> , <i>Ped. pentosaceus</i> , <i>P. acidilactici</i> , <i>W. cibara</i> , <i>W. paramesenteroides</i> , <i>Lb. fermentum</i> , <i>Lb. delbrueckii</i> subsp. <i>indicus</i> , <i>Saccharomycopsis</i> sp., <i>Candida</i> sp.	India, Nepal, Sri Lanka, Bangladesh, Pakistan	Harun-ur-Rashid et al. (2007), Patil et al. (2010)
<i>Ergo</i>	Milk	Acid fermented butter milk	<i>Lactobacillus</i> sp., <i>Lactococcus</i> sp.	Ethiopia	Steinkraus (1996)
<i>Filmjök</i>	Cow milk	Less-sour than yoghurt, yoghurt-like	<i>Lc. lactis</i> and <i>Leuc. mesenteroides</i>	Sweden	Kosikowski and Mistry (1997)
<i>Gariss</i>	Camel milk	Acidic, liquid, refreshing beverage	LAB	Sudan	Akabanda et al. (2013)
<i>Gheu/ghee</i>	Cow milk	Soft, oily mass, solid, butter	<i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lc. lactis</i> subsp. <i>cremoris</i>	India, Nepal, Bhutan, Bangladesh, Pakistan	Tamang (2010a)

(Continued)

**Table 1.1 (Continued) Some Common and Uncommon Ethnic Fermented Milk Products of the World**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Kefir</i>	Goat, sheep, cow	Alcoholic fermented milk, effervescent milk	<i>Tor. holmii</i> , <i>Tor. delbruechii</i> , <i>Lb. brevis</i> , <i>Lb. caucasicus</i> , <i>Strep. thermophilus</i> , <i>Lb. bulgaricus</i> , <i>Lb. plantarum</i> , <i>Lb. casei</i> , <i>Lb. brevis</i>	Russia	Bernardeau et al. (2006)
<i>Kesong Puti</i> , <i>Keso</i> , <i>Kesiyo</i>	Carabao's (buffalo) milk or cow carabao's milk, salt, Abomasal extracts coagulant, starter	White cheese, soft cheese	<i>Lb. helveticus</i> , <i>Lact. lactis</i> , <i>Lb. rhamnosus</i> , <i>Leuc. mesenteroides</i> , <i>Lb. acidophilus</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. curvatus</i>	Philippines	Kisworo (2003)
<i>Kishk</i>	Milk, wheat	Fermented milk wheat mix, drink	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. bulgaricus</i> , <i>Lb. casei</i> , <i>Strep. thermophilus</i>	Egypt	Bernardeau et al. (2006)
<i>Kurut</i>	Yak milk	Naturally fermented milk, drink	LAB	China	Sun et al. (2010)
<i>Kushuk</i>	Milk, wheat	Fermented milk wheat mix, drink	<i>Lb. plantarum</i> , <i>Lb. brevis</i>	Iraq	Bernardeau et al. (2006)
<i>Koumiss</i>	Milk	Acid fermented milk, drink	<i>Lb. bulgaricus</i> , <i>Torula</i> sp., <i>Lb. salivarius</i> , <i>Lb. buchneri</i> , <i>Lb. heveticus</i> , <i>Lb. plantarum</i> , <i>Lb. acidophilus</i>	Russia	Hao et al. (2010)
<i>Laban rayeb</i>	Milk	Acid fermented milk, yoghurt-like	<i>Lb. casei</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lact. lactis</i> , <i>Sacch. kefir</i> , <i>Leuconostoc</i> sp.	Egypt	Bernardeau et al. (2006)

<i>Laban zeer</i>	Milk	Acid fermented milk	<i>Lb. casei</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lc. lactis</i> , <i>Lc. lactis</i>	Egypt	Bernardeau et al. (2006)
<i>Lassi</i>	Cow milk	Acidic, buttermilk, refreshing beverage	<i>Lb. Acidophilus</i> , <i>Strep. thermophilus</i>	India, Nepal, Bhutan, Bangladesh, Pakistan	Patidar and Prajapati (1998)
<i>Långfil</i>	Cow milk	Elastic texture, sour, yoghurt-like	LAB	Sweden	Tamime (2005)
<i>Leben/Lben</i>	Cow milk	Sour milk	<i>Candida</i> sp., <i>Saccharomyces</i> sp., <i>Lactobacillus</i> sp., <i>Leuconostoc</i> sp.,	North, East Central Africa	Odufa and Oyewole (1997)
<i>Liban-argeel</i>	Sheep, goat, cow, buffalo milk	Acid fermented milk	LAB	Iraq	Bernardeau et al. (2006)
<i>Maa</i>	Yak milk	Mild-acidic, viscous, butter	LAB, yeasts	China (Tibet), India, Bhutan	Tamang (2010a)
<i>Maziwa lala</i>	Milk	Yoghurt-like	<i>Strep. lactis</i> , <i>Strep. thermophilus</i>	Kenya	Olasupo et al. (2010)
<i>Mohi</i>	Cow milk	Acidic, buttermilk, refreshing beverage	<i>Lb. alimentarius</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lc. lactis</i> subsp. <i>cremoris</i> ; <i>Saccharomycopsis</i> spp. and <i>Candida</i> spp.	Nepal, India, Bhutan	Dewan and Tamang (2007)
<i>Mish</i>	Cow/camel milk	Acidic, semi-liquid, refreshing beverage	LAB	Sudan, Egypt	Bernardeau et al. (2006)
<i>Misti dahi</i> ( <i>mishti doi</i> , <i>lal dahi</i> , <i>payodhi</i> )	Buffalo/cow milk	Mild-acidic, thick-gel, sweetened curd, savory	<i>Strep. salivarius</i> subsp. <i>thermophilus</i> , <i>Lb. acidophilus</i> , <i>Lb. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Sacch. cerevisiae</i> .	India, Bangladesh	Ghosh and Rajorhia (1990), Gupta et al. (2000)

(Continued)

**Table 1.1 (Continued) Some Common and Uncommon Ethnic Fermented Milk Products of the World**

Product	Substrate	Sensory Property and Nature	Microorganisms	Country	References
Nunu	Raw cow milk	Naturally fermented milk	<i>Lb. fermentum</i> , <i>Lb. plantarum</i> , <i>Lb. helveticus</i> , <i>Leuc. mesenteroides</i> , <i>Ent. faecium</i> , <i>Ent. italicus</i> , <i>Weissella confuse</i> ; <i>Candida parapsilosis</i> , <i>C. rugosa</i> , <i>C. tropicalis</i> , <i>Galactomyces geotrichum</i> , <i>Pichia kudriavzevii</i> , <i>Sacch. cerevisiae</i>	Ghana	Akabanda et al. (2013)
Paneer	Buffalo or cow milk	Whey, soft, cheese-like product, fried snacks, curry	LAB	India, Nepal, Pakistan, Bangladesh, Middle East	Tamang (2012b)
Phrung	Yak milk	Mild-acidic, hard-mass like <i>chhurpi</i> , masticator	Unknown	India, China (Tibet)	Tamang (2010a)
Philu	Cow or yak milk, bamboo vessels	Cream like product, curry	<i>Lb. paracasei</i> , <i>Lb. bifementans</i> , <i>Ent. faecium</i>	India, Nepal, Tibet (China)	Dewan and Tamang (2007)
Pheuja or suja	Tea-yak butter, salt	Salty with buttery flavor, liquid, Refreshing tea	Unknown	India, China (Tibet), Bhutan, Nepal	Tamang (2010a)
Rob	Cow, goat, sheep milk	Mild-acidic, savory	LAB	Sudan	Akabanda et al. (2013)
Shrikhand	Cow, buffalo milk	Acidic, concentrated sweetened viscous, savory	<i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lc. lactis</i> subsp. <i>diacetylactis</i> , <i>Lc. lactis</i> subsp. <i>cremoris</i> , <i>Strep. thermophilus</i> , <i>Lb. delbruecki</i> subsp. <i>Bulgaricus</i>	India	Aneja et al. (2002)

<i>Somar</i>	Yak or cow milk	Buttermilk	<i>Lb. paracasei</i> , <i>Lact. Lactis</i>	India, Nepal	Dewan and Tamang. (2007)
<i>Sour milk kerbah</i>	Milk	Acid fermented milk	<i>Lact. lactis</i> , <i>Sacch. kefir</i> , <i>Lb. casei</i> , <i>Lb. brevis</i> , <i>Lb. plantarum</i>	Egypt	Mayo et al. (2010)
<i>Sua chua</i>	Dried skim milk, starter, sugar	Acid fermented milk	<i>Lb. bulgaricus</i> , <i>Strep. thermophilus</i>	Vietnam	Alexandraki et al. (2013)
<i>Shyow</i>	Yak milk	Acidic, thick-gel viscous, curd-like, savory	LAB, yeasts	China (Tibet), Bhutan, India	Tamang (2010a)
<i>Tarag</i>	Cow, yak, goat milk	Acidic, sour, drink	<i>Lb. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>Lb. helveticus</i> , <i>Strep. thermophilus</i> , <i>Sacch. cerevisiae</i> , <i>Issatchenkia orientalis</i> , <i>Kazachstania unispora</i>	Mongolia	Watanabe et al. (2008)
<i>Viili</i>	Cow milk	Thick and sticky, sweet taste, breakfast	<i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lc. lactis</i> subsp. <i>cremoris</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> biovar. <i>Diacetylactis</i> , <i>Leuc. mesenteroides</i> subsp. <i>cremoris</i> , <i>G. candidum</i> , <i>K. marxianus</i> , <i>P. fermentans</i>	Finland	Kahala et al. (2008)
<i>Wara</i>	Milk	Sweet taste, beverage	<i>Lc. lactis</i> , <i>Lactobacillus</i> sp.	West Africa	Olasupo et al. (2010)
Yoghurt	Animal milk	Acidic, thick-gel viscous, curd-like product, savory	<i>Strep. thermophilus</i> , <i>Lb. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>Lb. acidophilus</i> , <i>Lb. casei</i> , <i>Lb. rhamnosus</i> , <i>Lb. gasserii</i> , <i>Lb. johnsonii</i> , <i>Bifidobacterium</i> spp.	Europe, Australia, America	Tamime and Robinson (2007)

(Josephsen and Jespersen 2004). Examples of naturally fermented milks are *dahi*, *lassi*, *misti dahi*, *srikhand*, *chhu*, *chhurpi*, *mohi*, *philu*, *shoyu*, *somar* (cow/buffalo/yak milk) of India, Nepal, Pakistan, Bhutan, and Bangladesh (Harun-ur-Rashid et al. 2007, Sarkar 2008, Patil et al. 2010, Tamang 2010a, Tamang et al. 2012), *kurut* of China (Sun et al. 2010), *aaruuul*, *airag*, *byasulag*, *chi-gee*, *eezgii*, *tarag*, and *khoormog* of Mongolia (Watanabe et al. 2008, Takeda et al. 2011, Oki et al. 2014), *ergo* of Ethiopia, *kad*, *lben*, *laban*, *rayeb*, *zabady*, *zeer* of Morocco and Northern African and Middle East countries, *rob* (from camel milk), *biruni* (cow/camel milk), *mish* (cow/camel milk) of Sudan, *amasi* (*hodzeko*, *mukaka wakakora*) of Zimbabwe, *nunu* (from raw cow milk) of Ghana (Akabanda et al. 2013), *filmjök* and *långfil* of Sweden (Mayo et al. 2010), *koumiss* or *kumis* or *kumys* or *kymys* of the Caucasian area (Wu et al. 2009).

*Lc. lactis* subsp. *cremoris*, and *Lc. lactis* subsp. *lactis* are found among the dominant microbiota along with other mesophilic lactobacilli (*Lb. casei*/*Lb. paracasei*, *Lb. fermentum*, *Lb. helveticus*, *Lb. plantarum*, and/or *Lb. acidophilus*), *Ent. faecium*, and species of *Leuconostoc* and *Pediococcus* in naturally fermented milks (Tamang et al. 2000, Mathara et al. 2004, Patrignani et al. 2006, Dewan and Tamang 2006, 2007, Yu et al. 2011, Akabanda et al. 2013). Yeasts present in naturally fermented milks are *Candida lusitanae*, *C. parapsilosis*, *C. rugosa*, *C. tropicalis*, *Kluyveromyces* (*Kl.*) *marxianus*, *Sacch. cerevisiae*, *Galactomyces geotrichum*, *Pichia kudriavzevii*, and others (Gadaga et al. 2000, Dewan and Tamang 2006, Akabanda et al. 2013). *Koumiss* or *kumis* or *kumys* or *kymys* is a natural fermented dairy product of the Caucasian area. *Lb. casei*, *Lb. coryniformis*, *Lb. curvatus*, *Lb. helveticus*, *Lb. kefiranofaciens*, *Lb. kefir*, *Lb. paracasei*, *Lb. plantarum*, *Lb. fermentum*, and *Leuc. mesenteroides* (Ying et al. 2004, Watanabe et al. 2008, Wu et al. 2009), *Lb. acidophilus*, *Lb. fermentum*, and *Lb. kefiranofaciens* (dominant LAB), *E. faecalis*, *Lc. lactis*, *Lb. buchneri*, *Lb. jensenii*, *Lb. kefir*, *Lb. kitasatonis*, *Lb. paracasei*, *Leuc. mesenteroides*, and *Strep. thermophilus* (Hao et al. 2010), yeasts *Sacch. cerevisiae*, *Issatchenkia orientalis*, *Kazachstania unispora*, *Kl. marxianus*, *Pichia mandshurica* (Watanabe et al. 2008) were isolated from *koumiss*.

### 1.5.2 Fermented Cereal Foods

The well-documented fermented cereal foods of the world (Table 1.2) are sourdough of Europe, America, and Australia (de Vuyst et al. 2009), *selroti* of India and Nepal (Yonzan and Tamang 2009), *idli* of India and Sri Lanka (Sridevi et al. 2010), *dosa* of India and Sri Lanka (Soni et al. 1986), *mawè* and *gowé* of Benin (Vieira-Dalodé et al. 2007), *ben-saalga* of Burkino Faso and Ghana (Humblot and Guyot 2009), *kisra* of Sudan (Hamad et al. 1997), *kenkey* of Ghana (Oguntoyinbo et al. 2011), *togwa* of Tanzania (Mugula et al. 2003), *ting* of Botswana (Sekwati-Monang and Gänzle 2011), *ogi* and *kunu-zaki* of Nigeria (Oguntoyinbo et al. 2011), and *tarhana* of Turkey, Cyprus and Greece (Sengun et al. 2009). Cereal fermentation is characterized by a complex microbial ecosystem, mainly represented by the species of LAB and yeasts (Corsetti and Settanni 2007), whose fermentation confers to the resulting bread its characteristic features such as palatability and high sensory quality (Blandino et al. 2003). The species of *Enterococcus*, *Lactococcus*, *Lactobacillus*, *Leuconostoc*, *Pediococcus*, *Streptococcus*, and *Weissella* are commonly associated with cereal fermentation (Guyot 2010). A native strain of *Sacch. cerevisiae* is the principal yeast of most bread fermentations (Hammes et al. 2005). Other non-*Saccharomyces* yeasts are also significant in many cereal fermentations which include *Candida*, *Debaryomyces*, *Hansenula*, *Pichia*, *Trichosporon*, *Yarrowia* (Foschino et al. 2004, Veinocchi et al. 2006). *Lb. plantarum*, *Lb. panis*, *Lb. sanfranciscensis*, *Lb. pontis*, *Lb. brevis*, *Lb. curvatus*, *Lb. sakei*, *Lb. alimentarius*, *Lb. fructivorans*, *Lb. paralimentarius*, *Lb. pentosus*, *Lb. spicheri*, *Lb. crispatus*, *Lb. delbrueckii*, *Lb. fermentum*, *Lb. reuteri*, *Lb. acidophilus*, *Lc. lactis*, *Leuc. mesenteroides*, *Ped. pentosaceus*, *W. confuse*. The yeasts



**Table 1.2 Some Common and Uncommon Ethnic Fermented Cereal Foods of the World**

<i>Product</i>	<i>Raw Material/ Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Abreh</i>	Sorghum	Solid state and submerged	<i>Lb. plantarum</i>	Sudan	Odufa and Oyewole (1997)
<i>Aliha</i>	Maize, sorghum	Nonalcoholic beverage	LAB	Ghana, Togo, Benin	Odufa and Oyewole (1997)
<i>Ambali</i>	Millet, rice	Acidic, pan cake Shallow-fried, staple	LAB	India	Tamang (2010a)
<i>Ang-kak</i>	Red rice	Colorant	<i>Monascus purpureus</i>	China, Taiwan, Thailand, Phillipines	Steinkraus (1996)
<i>Banku</i>	Maize and cassava	Staple food	<i>Lactobacillus</i> sp., yeasts	Ghana	Campbell-Platt (1987)
<i>Bahtura</i>	Wheat flour	Deep-fried bread	LAB, yeasts	India	Campbell-Platt (1987)
<i>Boza</i>	Cereals	Sour refreshing liquid	<i>Lactobacillus</i> sp., <i>Lactococcus</i> sp., <i>Pediococcus</i> sp., <i>Leuconostoc</i> sp., <i>Sacch. cerevisiae</i>	Bulgaria, Balkan	Blandino et al. (2003)
<i>Burukutu</i>	Sorghum and cassava	Creamy, liquid, drink	<i>Sacch. cerevisiae</i> , <i>Sacch. chavelieri</i> , <i>Leuc. mesenteroides</i> , <i>Candida</i> sp., <i>Acetobacter</i> sp.	Nigeria	Odufa and Oyewole (1997), Kolawole et al. (2013)
<i>Busa</i>	Maize, sorghum, millet	Submerged	<i>Sacch. cerevisiae</i> , <i>Schizosacchromyces pombe</i> , <i>Lb. plantarum</i> , <i>Lb. helveticus</i> , <i>Lb. salivarius</i> , <i>Lb. casei</i> , <i>Lb. brevis</i> , <i>Lb. buchneri</i> , <i>Leuc. mesenteroides</i> , <i>Ped. damnosus</i>	East Africa, Kenya	Odufa and Oyewole (1997), Blandino et al. (2003)

(Continued)

**Table 1.2 (Continued) Some Common and Uncommon Ethnic Fermented Cereal Foods of the World**

Product	Raw Material/ Substrate	Sensory Property and Nature	Microorganisms	Country	References
<i>Ben-saalga</i>	Pearl millet	Weaning food	<i>Lactobacillus</i> sp., <i>Pediococcus</i> sp., <i>Leuconostoc</i> sp., <i>Weissela</i> sp., yeasts	Burkina Faso, Ghana	Tou et al. (2007)
<i>Chilra</i>	Wheat, barley, buckwheat	Staple	LAB, <i>Sacch. cerevisiae</i>	India	Thakur et al. (2004)
<i>Dégué</i>	Millet	Condiment	<i>Lb. gasserii</i> , <i>Lb. fermentum</i> , <i>Lb. brevis</i> , <i>Lb. casei</i> , <i>Enterococcus</i> sp.	Burkina Faso	Abriouel et al. (2006)
<i>Dosa</i>	Rice and black gram	Thin, crisp pancake, Shallow-fried, staple	<i>Leuc. mesenteroides</i> , <i>Ent. faecalis</i> , <i>Tor. candida</i> , <i>Trichosporon pullulans</i>	India, Sri Lanka, Malaysia, Singapore	Soni et al. (1986)
<i>Enjera/Injera</i>	Tef flour, wheat	Acidic, sourdough, leavened, pancake- like bread, staple	<i>Lb. pontis</i> , <i>Lb. plantarum</i> , <i>Leuc.</i> <i>mesenteroides</i> , <i>Ped. cerevisiae</i> , <i>Sacch. cerevisiae</i> , <i>Cand. glabrata</i>	Ethiopia	Olasupo et al. (2010)
<i>Gowé</i>	Maize	Intermediate product used to prepare beverages, porridges	<i>Lb. fermentum</i> , <i>Lb. reuteri</i> , <i>Lb. brevis</i> , <i>Lb. confusus</i> , <i>Lb. curvatus</i> , <i>Lb. buchneri</i> , <i>Lb. salivarius</i> , <i>Lact. lactis</i> , <i>Ped. pentosaceus</i> , <i>Ped. acidilactici</i> , <i>Leuc. mesenteroides</i> ; <i>Candida</i> <i>tropicalis</i> , <i>C. krusei</i> , <i>Kluyveromyces</i> <i>marxianus</i>	Benin	Vieira-Dalodé et al. (2007), Greppi et al. (2013a)
<i>Hakua</i>	Rice	Strong off-flavor, therapeutic uses	Unknown	Nepal, India	Tamang (2005)
<i>Hopper</i>	Rice, coconut water	Steak-baked, pancake, staple	<i>Sacch. cerevisiae</i> , LAB	Sri Lanka	Steinkraus (1996)
<i>Hussuwa</i>	Sorghum	Cooked dough	<i>Lb. fermentum</i> , <i>Ped. acidilactici</i> , <i>Ped. pentosaceus</i> , Yeasts	Sudan	Yousif et al. (2010)

<i>Huitlacoche</i> or 'maize mushroom'	cobs of pre-harvest maize	Large fruiting body edible, condiment	<i>Ustilago maydis</i>	Mexico	Alexandraki et al. (2013)
<i>Hulumur</i>	Sorghum, rice, millet	Nonalcoholic drink	<i>Leuc. mesenteroides</i> , <i>Lb. Plantarum</i> , <i>Lactobacillus</i> sp.	Sudan, Turkey	Campbell-Platt (1987)
<i>Idli</i>	Rice, blackgram dhal or other dehusked pulses	Mild-acidic, soft, moist, spongy pudding; staple, breakfast	<i>Leuc. mesenteroides</i> , <i>Lb. delbrueckii</i> , <i>Lb. fermenti</i> , <i>Lb. coryniformis</i> , <i>Ped. acidilactis</i> , <i>Ped. cerevisiae</i> , <i>Streptococcus</i> sp., <i>Ent. faecalis</i> , <i>Lact. lactis</i> , <i>B. amyloliquefaciens</i> , <i>Cand. cacaioi</i> , <i>Cand. fragicola</i> , <i>Cand. glabrata</i> , <i>Cand. kefir</i> , <i>Cand. pseudotropicalis</i> , <i>Cand. sake</i> , <i>Cand. tropicalis</i> , <i>Deb. hansenii</i> , <i>Deb. tamarii</i> , <i>Issatchenkia terricola</i> , <i>Rhiz. graminis</i> , <i>Sacch. cerevisiae</i> , <i>Tor. candida</i> , <i>Tor. holmii</i>	India, Sri Lanka, Malaysia, Singapore	Steinkraus et al. (1967), Sridevi et al. (2010)
<i>Jalebi</i>	Wheat flour	Crispy sweet, donut-like, deep-fried, snacks	<i>Sacch. Bayanus</i> , <i>Lb. fermentum</i> , <i>Lb. buchneri</i> , <i>Lact. lactis</i> , <i>Ent. faecalis</i> , <i>Sacch. cerevisiae</i>	India, Nepal, Pakistan	FAO (1999)
<i>Kenkey</i>	Maize	Acidic, solid, steamed dumpling, staple	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Ent. cloacae</i> , <i>Acinetobacter</i> sp., <i>Sacch. cerevisiae</i> , <i>Cand. mycoderma</i>	Ghana	Odufa and Oyewole (1997), Oguntoyinbo et al. (2011)
<i>Khanom-jeen</i>	Rice	Noodle, staple	<i>Lactobacillus</i> sp., <i>Streptococcus</i> sp., <i>Rhizopus</i> sp., <i>Mucor</i> sp.	Thailand	Blandino et al. (2003)
<i>Khamak</i> ( <i>Kao-mak</i> )	Glutinous rice, <i>Look-pang</i> (starter)	Dessert	<i>Rhizopus</i> sp., <i>Mucor</i> sp., <i>Penicillium</i> sp., <i>Aspergillus</i> sp., <i>Endomycopsis</i> sp., <i>Hansenula</i> sp., <i>Saccharomyces</i> sp.	Thailand	Alexandraki et al. (2013)

(Continued)

Table 1.2 (Continued) Some Common and Uncommon Ethnic Fermented Cereal Foods of the World

Product	Raw Material/ Substrate	Sensory Property and Nature	Microorganisms	Country	References
<i>Kichudok</i>	Rice	Steamed cake, side dish	<i>Leuc. mesenteroides</i> , <i>Ent. faecalis</i> , <i>Saccharomyces</i> sp.	Korea	Von Mollendorff (2008)
<i>Kunu-zaki</i>	Maize, sorghum, millet	Mild-acidic, viscous, porridge, staple	<i>Lb. plantarum</i> , <i>Lb. pantheris</i> , <i>Lb. vaccinostercus</i> , <i>Corynebacterium</i> sp., <i>Aerobacter</i> sp., <i>Cand. mycoderma</i> , <i>Sacch. cerevisiae</i> , <i>Rhodotorula</i> sp., <i>Cephalosporium</i> sp., <i>Fusarium</i> sp., <i>Aspergillus</i> sp., <i>Penicillium</i> sp.	Nigeria	Oguntoyinbo et al. (2011)
<i>Kisra</i>	Sorghum	Thin pancake bread, staple	<i>Ped. pentosaceus</i> , <i>Lb. confusus</i> , <i>Lb. brevis</i> , <i>Lactobacillus</i> sp., <i>Erwinia ananas</i> , <i>Klebsiella pneumoniae</i> , <i>Ent. cloacae</i> , <i>Cand. intermedia</i> , <i>Deb. hansenii</i> , <i>Aspergillus</i> sp., <i>Penicillium</i> sp., <i>Fusarium</i> sp., <i>Rhizopus</i> sp.	Sudan	Mohammed et al. (1991), Hamad et al. (1997)
<i>Kishk</i>	Wheat, milk	Refreshing beverage	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. casei</i> , <i>B. subtilis</i> , Yeasts	Egypt	Blandino et al. (2003)
<i>Koko</i>	Maize	Porridge	<i>Ent. clocae</i> , <i>Acinetobacter</i> sp., <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Sacch. cerevisiae</i> , <i>Cand. mycoderma</i>	Ghana	Blandino et al. (2003)
<i>Kunu-zaki</i>	Maize (white and yellow), red sorghum	Mild, sour liquid/porridge, staple	<i>W. confusa</i> , <i>Strep. lutetiensis</i> , <i>Strep. gallolyticus</i> subsp. <i>macedonicus</i> ,	West Africa, Nigeria	Olasupo et al. (2010)
<i>Lao-chao</i>	Rice	Paste, soft, juicy, glurinous desert	<i>Rhiz. oryzae</i> , <i>Rhiz. Chinensis</i> , <i>Chlamydomucor oryzae</i> , <i>Sacchromycopsis</i> sp.	China	Blandino et al. (2003)

<i>Maheu</i>	Maize, sorghum, millet	Refreshing beverage	<i>Lb. delbrueckii</i>	South Africa	Steinkraus (2004)
<i>Mahewu</i>	Maize	Refreshing beverage	<i>Lb. delbruchi, Lact. lactis</i>	South Africa	Blandino et al. (2003)
<i>Mawè</i>	Maize	Intermediate product used to prepare beverages, porridges	<i>Lb. fermentum, Lb. reuteri, Lb. brevis, Lb. confusus, Lb. curvatus, Lb. buchneri, Lb. salivarius, Lact. lactis, Ped. pentosaceus, Ped. acidilactici, Leuc. mesenteroides; Candida glabrata, Sacch. cerevisiae, Kluyveromyces marxianus, Clavispora lusitaniae</i>	Benin, Togo	Hounhouigan et al. (1993), Greppi et al. (2013a,b)
<i>Masvusvu</i>	Maize	Refreshing beverage	LAB	Zimbabwe	Alexandraki et al. (2013)
<i>Marchu</i>	Wheat flour	Baked bread	Unknown	India, Pakistan	Tamang (2010a)
<i>Mbege</i>	Maize, sorghum, millet	Submerged	<i>Sacch. cerevisiae, Schizosaccharomyces pombe, Lb. plantarum, Leuc. mesenteroides</i>	Tanzania	Odufa and Oyewole (1997)
<i>Me</i>	Rice	Acidic, sour, condiment	LAB	Vietnam	Alexandraki et al. (2013)
<i>Minchin</i>	Wheat gluten	Solid, condiment	<i>Paceilomyces sp., Aspergillus sp., Cladosporium sp., Fusarium sp., Syncephalastum sp., Penicillium sp., Trichothecium sp.</i>	China	Blandino et al. (2003)
<i>Mungbean starch</i>	Mungbean	Fermented noodle	<i>Leuc. mesenteroides</i>	Thailand	Alexandraki et al. (2013)

(Continued)

Table 1.2 (Continued) Some Common and Uncommon Ethnic Fermented Cereal Foods of the World

Product	Raw Material/ Substrate	Sensory Property and Nature	Microorganisms	Country	References
Naan	Wheat flour	Leaved bread, baked	<i>Sacch. cerevisiae</i> , LAB	India, Pakistan, Afghanistan	Batra (1986)
Ogi	Maize, sorghum, millet	Mild-acidic, viscous, porridge, staple	<i>Lb. plantarum</i> , <i>Lb. pantheris</i> , <i>Lb. vaccinostercus</i> , <i>Corynebacterium</i> sp., <i>Aerobacter</i> sp., <i>Candida krusei</i> , <i>Clavispora lusitaniae</i> , <i>Sacch. cerevisiae</i> , <i>Rhodotorula</i> sp., <i>Cephalosporium</i> sp., <i>Fusarium</i> sp., <i>Aspergillus</i> sp., <i>Penicillium</i> sp.	Nigeria	Odufa and Oyewole (1997), Greppi et al. (2013a)
Perkarnaya	Rye	Acidic, aerated bread	Yeasts, LAB	Russia	Alexandraki et al. (2013)
Pito	Maize, sorghum, millet	Submerged	<i>Geotrichum candidum</i> , <i>Lactobacillus</i> sp., <i>Candida</i> sp.	West Africa	Odufa and Oyewole (1997)
Poto poto (Gruel)	Maize	Slurry	<i>Lb. gasserii</i> , <i>Lb. plantarum</i> / <i>paraplantarum</i> , <i>Lb. acidophilus</i> , <i>Lb. delbrueckii</i> , <i>Lb. reuteri</i> , <i>Lb. casei</i> , <i>Bacillus</i> sp., <i>Enterococcus</i> sp., yeasts	Congo	Abriouel et al. (2006)
Pozol	Maize	Mild-acidic, thick viscous, porridge, staple	<i>Lactobacillus</i> sp., <i>Leuconostoc</i> sp., <i>Candida</i> sp., <i>Enterobacteriaceae</i> , <i>B. cereus</i> , <i>Paracolobactrum aerogenoides</i> , <i>Agrobacterium azotophilum</i> , <i>Alkaligenes</i> <i>pozolis</i> , <i>E. coli</i> var. <i>napolitana</i> , <i>Pseudomonas mexicana</i> , <i>Kleb.</i> <i>pneumoniae</i> , <i>Saccharomyces</i> sp., molds	Mexico	FAO (1998)
Pumpernickel	Rye	Acidic, full-grain, aerated bread; long shelf-life	Yeasts, LAB, as for rye sourdough	Switzerland, Germany	Alexandraki et al. (2013)



<i>Puto</i>	Rice	Steamed cake, breakfast	<i>Leuc. mesenteroides</i> , <i>Ent. faecalis</i> , <i>Ped. cerevisiae</i> , yeasts	Philippines	Steinkraus (1996)
<i>Rabadi</i>	Buffalo or cow milk and cereals, pulses	Mild-acidic, thick slurry-like product	<i>Ped. acidilactici</i> , <i>Bacillus</i> sp., <i>Micrococcus</i> sp.; Yeasts	India, Pakistan	Ramakrishnan (1979), Gupta et al. (1992)
<i>Sourdough bread</i>	Rye	Sandwich, bread	<i>Lb. pontis</i> and <i>Lb. panis</i> , <i>Lb. amylovorus</i> , <i>Lb. acidophilus</i> , <i>Lb. crispatus</i> , <i>Lb. delbrueckii</i> , <i>Lb. fermentum</i> , <i>Lb. reuteri</i> , <i>Sacch. cerevisiae</i> , <i>Issatchenkia orientalis</i>	Germany, Northern Europe	Iacumin et al. (2009)
<i>San Francisco</i>	(Rye), mainly wheat	Mild-acidic, leavened bread	<i>Lb. sanfranciscensis</i> , <i>Lb. alimentarius</i> , <i>Lb. brevis</i> , <i>Lb. fructivorans</i> , <i>Lb. paralimentarius</i> , <i>Lb. pentosus</i> , <i>Lb. plantarum</i> , <i>Lb. pontis</i> , <i>Lb. spicheri</i> , <i>Leuc. mesenteroides</i> , <i>W. confusa</i>	USA	Gänzle et al. (1998)
<i>Seera</i>	Wheat grains	Dried, sweet dish	Unknown	India, Pakistan	Thakur et al. (2004)
<i>Selroti</i>	Rice-wheat flour-milk	Pretzel-like, deep-fried bread, staple	<i>Leuc. mesenteroides</i> , <i>Ent. faecium</i> , <i>Ped. pentosaceus</i> and <i>Lb. curvatus</i> , <i>Sacch. cerevisiae</i> , <i>Sacch. kluyveri</i> , <i>Deb. hansenii</i> , <i>P. burtonii</i> , <i>Zygosaccharomyces rouxii</i>	India, Nepal, Bhutan	Yonzan and Tamang (2010, 2013)
<i>Siddu</i>	Wheat flour, opium seeds, walnut	Steamed bread, oval-shaped, staple	<i>Sacch. cerevisiae</i> , <i>Cand. valida</i>	India	Thakur et al. (2004)
<i>Sourdough bread</i>	Rye, wheat	Mild-acidic, leavened bread	<i>Lb. sanfranciscensis</i> , <i>Lb. alimentarius</i> , <i>Lb. buchneri</i> , <i>Lb. casei</i> , <i>Lb. delbrueckii</i> , <i>Lb. fructivorans</i> , <i>Lb. plantarum</i> , <i>Lb. reuteri</i> , <i>Lb. johnsonii</i> , <i>Cand. humili</i> , <i>Issatchenkia orientalis</i>	America, Europe, Australia	de Vuyst et al. (2009)

(Continued)

**Table 1.2 (Continued) Some Common and Uncommon Ethnic Fermented Cereal Foods of the World**

<i>Product</i>	<i>Raw Material/ Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Tapai Pulut</i>	Glutinous rice, <i>Ragi</i>		<i>Chlamydomucor</i> sp., <i>Endomycopsis</i> sp., <i>Hansenula</i> sp.	Malaysia	Steinkraus (1996)
<i>Tape Ketan</i>	Glutinous rice, <i>Ragi</i>		<i>Thizopus</i> sp., <i>Chlamydomucor</i> sp., <i>Candida</i> sp., <i>Endomycopsis</i> sp., <i>Saccharomyces</i> sp.	Indonesia	Steinkraus (1996)
<i>Tepache</i>	Maize, pineapple, apple or orange		<i>B. subtilis</i> , <i>B. graveolus</i> and the yeasts, <i>Tor. inconspicna</i> , <i>Sacch. cerevisiae</i> and <i>Cand. querehana</i>	Mexico	FAO (1998)
<i>Ting</i>	Sorghum	Sour taste	LAB	Botswana	Sekwati-Monang and Gänzle (2011)
<i>Togwa</i>	Cassava, maize, sorghum, millet	Fermented gruel or beverage	<i>Lb. brevis</i> , <i>Lb. cellobiosus</i> , <i>Lb. fermentum</i> , <i>Lb. plantarum</i> and <i>Ped. pentosaceus</i> , <i>Candida</i> <i>pelliculosa</i> , <i>C. tropicalis</i> , <i>Issatchenkia</i> <i>orientalis</i> , <i>Sacch. cerevisiae</i>	Tanzania	Mugula et al. (2003)
<i>Tarhana</i>	Sheep milk, wheat	Mild-acidic, sweet- sour, soup or biscuit	<i>Lb. bulgaricus</i> , <i>Strep. thermophilus</i> , yeasts	Cyprus, Greece, Turkey	Karagozlu et al. (2008)
<i>Taotjo</i>	Wheat, rice, soybeans	Semi-solid food, condiment	<i>Asp. oryzae</i>	East Indies	Blandino et al. (2003)
<i>Uji</i>	Maize, sorghum, millet, cassava flour	Acidic, sour, porridge, staple	<i>Leuc. mesenteroides</i> , <i>Lb. plantarum</i>	Kenya, Uganda, Tanzania	Odufa and Oyewole (1997)

*Sacch. cerevisiae*, *Sacch. exiguus*, *Candida humilis*, *C. milleri*, *Issatchenkia orientalis* were isolated from sourdough (Iacumin et al. 2009, Weckx et al. 2010). Gluten-free sourdough was prepared from buckwheat and/or teff flours using *Ped. pentosaceus*, *Leuc. holzapfelii*, *Lb. gallinarum*, *Lb. vaginalis*, *Lb. sakei*, *Lb. graminis*, *W. cibaria* (Moroni et al. 2011).

### 1.5.3 Fermented Vegetable Products

People eat plants, both domesticated and wild, preparing them according to a variety of recipes. Perishable and seasonal leafy vegetables, radish, cucumbers including young edible tender bamboo shoots are traditionally fermented into edible products using the indigenous knowledge of biopreservation. Mostly species of *Lactobacillus* and *Pediococcus*, followed by *Leuconostoc*, *Weissella*, *Tetragenococcus*, and *Lactococcus* (Watanabe et al. 2009a, Savadogo et al. 2011) have been isolated from various fermented vegetable foods of the world (Table 1.3). Species of LAB present in Korean *kimchi* are *Lc. lactis*, *Lb. brevis*, *Lb. curvatus*, *Lb. plantarum*, *Lb. sakei* subsp. *sakei*, *Luec. citreum*, *Leuc. gasicomitatum*, *Leuc. gelidum*, *Leuc. kimchii*, *Leuc. mesenteroides* subsp. *mesenteroides*, *Ped. pentosaceus*, *Weissella confusa*, *W. kimchii*, and *W. koreensis* (Shin et al. 2008, Nam et al. 2009, Park et al. 2010, Jung et al. 2011). A few species of non-LAB and yeasts were also reported from *kimchi* which included species of *Halococcus*, *Haloterrigena*, *Candida*, *Kluyveromyces*, *Lodderomyces*, *Natrialba*, *Natronococcus*, *Pichia*, *Saccharomyces*, *Sporisorium*, and *Trichosporon* (Chang et al. 2008). The species of LAB reported from *sauerkraut* are *Lc. lactis* subsp. *lactis*, *Lb. brevis*, *Lb. curvatus*, *Lb. plantarum*, *Lb. sakei*, *Leuc. fallax*, *Leuc. mesenteroides* and *Ped. pentosaceus* (Johanningsmeier et al. 2007, Plengvidhya et al. 2007). *Lb. brevis*, *Lb. casei*, *Lb. casei* subsp. *pseudopplantarum*, *Lb. fermentum*, *Lb. plantarum*, *Leuc. fallax*, *Ped. pentosaceus* constitute the native lactic flora in the Himalayan fermented vegetables such as *gundruk*, *sinki*, *goyang*, *khalti*, *inziangsang* (Karki et al. 1983, Tamang et al. 2005, Tamang and Tamang 2007, 2010). *Lb. brevis*, *Lb. lactis*, *Lb. fermentum*, *Lb. pentosus*, *Lb. plantarum*, *Leuc. mesenteroides* and *Ped. pentosaceus* are the functional LAB in *pao cai* or *suan cai*, the naturally fermented vegetable products of China and Taiwan (Yan et al. 2008, Huang et al. 2009). A complex microbial community in the brines of fermented olives based on a culture-independent study consisted of LAB (*Lb. pentosus*/*Lb. plantarum*, *Lb. paracoloinoides*, *Lb. vaccinostercus*/*Lb. suebicus* and *Pediococcus* sp.), both cultivable and uncultivable non-lactics (*Gordonia* sp./*Pseudomonas* sp., *Halorubrum orientalis*, *Halosarcina pallid*, *Sphingomonas* sp./*Sphingobium* sp./*Sphingopyxis* sp., *Thalassomonas agarivorans*) and yeasts (*Candida* cf. *apicola*, *Pichia* sp., *Pic. manshurica*/*Pic. galeiformis*, *Sacch. cerevisiae*) (Abriouel et al. 2011).

*Sunki* is an ethnic nonsalted and fermented vegetable product of Japan prepared from the leaves and stems of the red turnip (Wacher et al. 2010). *Lb. plantarum*, *Lb. brevis*, *Lb. buchneri*, *Lb. kisonensis*, *Lb. otakiensis*, *Lb. rapi*, *Lb. sunkii*, *E. faecalis*, *B. coagulans*, and *P. pentosaceus* have been isolated from *sunki* (Endo et al. 2008, Watanabe et al. 2009a). *Fu-tsai* and *suan-tsai* are the ethnic fermented mustard products of Taiwan prepared by the Hakka tribes eaten as soup, fried with shredded meat, or stewed with meat (Chao et al. 2009). *Ped. pentosaceus* and *Tetragenococcus halophilus* (Chen et al. 2006), *Lb. farciminis*, *Leuc. mesenteroides*, *Leuc. pseudomesenteroides*, *W. cibaria*, and *W. paramesenteroides* (Chao et al. 2009), *Lb. futsaii* (Chao et al. 2012) are isolated from *fu-tsai* and *suan-tsai*.

*Ent. durans*, *Lb. brevis*, *Lb. coryniformis*, *Lb. curvatus*, *Lb. delbrueckii*, *Lb. plantarum*, *Lb. xylo-sus*, *Leuc. citreum*, *Leuc. fallax*, *Leuc. lactis*, *Leuc. mesenteroides*, *Ped. pentosaceus*, *Tetra. halophilus* were reported from Indian fermented bamboo shoots (Tamang and Sarkar 1996, Tamang et al. 2008, Tamang and Tamang 2009, Tamang et al. 2012, Sonar and Halami 2014). Species of *B. cereus*, *B. pumilus*, *B. subtilis* and *Pseudomonas fluorescens* along with LAB were also isolated

**Table 1.3 Some Common and Uncommon Ethnic Fermented Vegetable Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Anishi</i>	Taro leaves	Acidic, wet	LAB	India	Tamang (2010a)
<i>Bastanga</i>	Bamboo shoot	Acidic, soft	LAB	India	Tamang (2010a)
<i>Burong mustala</i>	Mustard	Acidic, wet	<i>Lb. brevis</i> , <i>Ped. cerevisiae</i>	Philippines	Rhee et al. (2011)
<i>Cucumbers (pickles)</i>	Cucumbers	Acidic, wet, pickle	<i>Leuc. mesenteroides</i> , <i>Ped. cerevisiae</i> , <i>Ped. acidilactici</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i>	Europe, USA, Canada	Vaughn (1985)
<i>Dha muoi</i>	Mustard and beet ( <i>dha muoi</i> ), eggplant ( <i>ca muoi</i> )	Acidic, wet	<i>Lb. fermentum</i> , <i>Lb. pentosus</i> , <i>Lb. plantarum</i> , <i>Ped. pentosaceus</i> , <i>Lb. brevis</i> , <i>Lb. paracasei</i> , <i>Lb. pantheris</i> , <i>Ped. acidilactici</i>	Vietnam	Nguyen et al. (2013b)
<i>Dakguadong</i>	Mustard leaf	Salad	<i>Lb. plantarum</i>	Thailand	Rhee et al. (2011)
<i>Ekung</i>	Bamboo shoot	Acidic, sour, soft, curry	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. casei</i> , <i>Tor. halophilus</i>	India	Tamang and Tamang (2009)
<i>Eup</i>	Bamboo shoot	Acidic, sour, dry, curry	<i>Lb. plantarum</i> , <i>Lb. fermentum</i> , <i>Lb. brevis</i> , <i>Lb. curvatus</i> , <i>Ped. pentosaceus</i> , <i>Leuc. mesenteroides</i> , <i>Leuc. fallax</i> , <i>Leuc. lactis</i> , <i>Leuc. citreum</i> , <i>Ent. durans</i>	India	Tamang and Tamang (2009)
<i>Fu-tsai</i>	Mustard	Acidic, sour	<i>Ent. faecalis</i> , <i>Lb. alimentarius</i> , <i>Lb. brevis</i> , <i>Lb. coryniformis</i> , <i>Lb. farciminis</i> , <i>Lb. plantarum</i> , <i>Lb. versmoldensis</i> , <i>Leuc. citreum</i> , <i>Leuc. mesenteroides</i> , <i>Leuc. pseudomesenteroides</i> , <i>Ped. pentosaceus</i> , <i>W. cibaria</i> , <i>W. paramesenteroides</i>	Taiwan	Chao et al. (2009, 2012)

<i>Goyang</i>	Wild vegetable	Acidic, sour, wet, soup	<i>Lb. plantarum</i> , <i>L. brevis</i> , <i>Lact. lactis</i> , <i>Ent. faecium</i> , <i>Ped. pentosaceus</i> , <i>Candida</i> sp.	India, Nepal	Tamang and Tamang (2007)
<i>Gundruk</i>	Leafy vegetable	Acidic, sour, dry, soup, side-dish	<i>Lb. fermentum</i> , <i>Lb. plantarum</i> , <i>Lb. casei</i> , <i>Lb. casei</i> subsp. <i>pseudoplantarum</i> , <i>Ped. pentosaceus</i> ,	India, Nepal, Bhutan	Karki et al. (1983), Tamang et al. (2005)
<i>Hirring</i>	Bamboo shoot tips	Acidic, sour, wet, pickle	<i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Lb. curvatus</i> , <i>Ped. pentosaceus</i> , <i>Leuc. mesenteroides</i> , <i>Leuc. fallax</i> , <i>Leuc. lactis</i> , <i>Leuc. citreum</i> , <i>Ent. durans</i> , <i>Lact. lactis</i>	India	Tamang and Tamang (2009)
<i>Hom-dong</i>	Red onion	Fermented red onion	<i>Leuc. mesenteroides</i> , <i>Ped. cerevisiae</i> , <i>Lb. plantarum</i> , <i>Lb. fermentum</i> , <i>Lb. buchneri</i>	Thailand	Phithakpol et al. (1995)
<i>Hum-choy</i>	Gai-choy	Chinese sauerkraut	<i>Pediococcus</i> sp., <i>Streptococcus</i> sp.,	China	Phithakpol et al. (1995)
<i>Inziang-sang</i>	Mustard leaves	Acidic, sour, dry, soup	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Ped. acidilactici</i>	India	Tamang et al. (2005)
<i>Jeruk</i>	Fruits and vegetables	Acidic, wet	LAB	Malaysia	Merican (1996)
<i>Jiang-sun</i>	Bamboo shoot, salt, sugar, <i>douchi</i> (fermented soybeans)	Fermented bamboo, side dish	<i>Lb. plantarum</i> , <i>Ent. faecium</i> , <i>Lc. lactis</i> subsp. <i>lactis</i>	Taiwan	Chen et al. (2010)
<i>Khalpi</i>	Cucumber	Acidic, sour, wet, pickle	<i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Ped.</i> <i>pentosaceus</i> , <i>Ped. Acidilactici</i> , <i>Leuc. Fallax</i>	India, Nepal	Tamang et al. (2005), Tamang and Tamang (2010)

(Continued)

**Table 1.3 (Continued) Some Common and Uncommon Ethnic Fermented Vegetable Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Kimchi (beachoo)</i>	Cabbage, green onion, hot pepper, ginger	Acidic, mild-sour, wet, side-dish	<i>Leuc. mesenteroides, Leuc. citreum, Leuc. gasicomitatum, Leuc. kimchii, Leuc. inhae, W. koreensis, W. cibaria, Lb. plantarum, Lb. sakei, Lb. delbrueckii, Lb. buchneri, Lb. brevis, Lb. fermentum, Ped. acidilactici, Ped. pentosaceus</i>	Korea	Nam et al. (2009), Jung (2012)
<i>Kimchi (Dongchimi)</i>	Radish, salt, water	Acidic, mild-sour, wet, soup, side-dish	<i>Leuc. mesenteroides, Lb. plantarum, Lb. brevis, Ped. cerevisiae</i>	Korea	Nam et al. (2009), Jung (2012)
<i>Kimchi (Kakdugi)</i>	Radish, salt, garlic, green onion, hot pepper, ginger	Acidic, mild-sour, wet, side-dish	<i>Leuc. mesenteroides, Lb. plantarum, Lb. brevis, Ped. cerevisiae</i>	Korea	Nam et al. (2009), Jung (2012)
<i>Lung-siej</i>	Bamboo shoot	Sour-acidic, soft	<i>Lb. brevis, Lb. plantarum, Lb. curvatus, Ped. pentosaceus, Leuc. mesenteroides, Leuc. fallax, Leuc. lactis, Leuc. citreum, Ent. durans</i>	India	Tamang (2010a)
<i>Naw-mai-dong</i>	Bamboo shoots	Acidic, wet	<i>Leuc. mesenteroides, Ped. cerevisiae, Lb. plantarum, Lb. brevis, Lb. fermentum, Lb. buchneri</i>	Thailand	Phithakpol et al. (1995)
<i>Mesu</i>	Bamboo shoot	Acidic, sour, wet	<i>Lb. plantarum, Lb. brevis, Lb. curvatus, Leu, citreum, Ped. pentosaceus</i>	India, Nepal, Bhutan	Tamang and Sarkar (1996), Tamang et al. (2008)
<i>Oiji</i>	Cucumber, salt, water	Fermented cucumber	<i>Leuc. mesenteroides, Lb. brevis, Lb. plantarum, Ped. cerevisiae</i>	Korea	Alexandraki et al. (2013)



<i>Olives</i> (fermented)	Olive	Acidic, wet, Salad, side dish	<i>Leuc. mesenteroides</i> , <i>Ped. pentosaceus</i> ; <i>Lb. plantarum</i> <i>Lb. pentosus/Lb. plantarum</i> , <i>Lb. paracollinoides</i> , <i>Lb. vaccinostercus/Lb. suebicus</i> and <i>Pediococcus</i> sp. non-lactics ( <i>Gordonia</i> sp./ <i>Pseudomonas</i> sp., <i>Halorubrum orientalis</i> , <i>Halosarcina pallid</i> , <i>Sphingomonas</i> sp./ <i>Sphingobium</i> sp./ <i>Sphingopyxis</i> sp., <i>Thalassomonas agarivorans</i> ) and yeasts ( <i>Candida</i> cf. <i>apicola</i> , <i>Pichia</i> sp., <i>Pic. manshurica/Pic. galeiformis</i> , <i>Sacch. cerevisiae</i> )	USA, Spain, Portugal, Peru, Chile	Abriouel et al. (2011)
<i>Pak-gard-dong</i>	Leafy vegetable, salt, boiled rice	Acidic, wet, side dish	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Ped. cerevisiae</i>	Thailand	Phithakpol et al. (1995)
<i>Pak-sian-dong</i>	Leaves of <i>Gynandropis pentaphylla</i>	Acidic, wet, side dish	<i>Leuc. mesenteroides</i> , <i>Ped. cerevisiae</i> , <i>Lb. plantarum</i> , <i>Lb. germentum</i> , <i>Lb. buchneri</i>	Thailand	Phithakpol et al. (1995)
<i>Poi</i>	Taro corms	Acidic, semi-solid	LAB, yeasts	Hawaii	Alexandraki et al. (2013)
<i>Pao cai</i>	Cabbage	Sweet and sour rather than spicy, breakfast	<i>Lb. pentosus</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. lactis</i> , <i>Lb. fermentum</i> , and <i>Leuc. mesenteroides</i> , and <i>Ped. pentosaceus</i>	China	Yan et al. (2008)
<i>Sauerkraut</i>	Cabbage	Acidic, sour, wet, salad, side dish	<i>Leuc. mesenteroides</i> , <i>Ped. pentosaceus</i> ; <i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Lb. sakei</i>	Europe, USA, Canada, Australia	Johanningsmeier et al. (2007)

(Continued)

**Table 1.3 (Continued) Some Common and Uncommon Ethnic Fermented Vegetable Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Sayur asin</i>	Mustard leaves, cabbage, salt, coconut	Acidic, sour, wet, salad	<i>Leuc. mesenteroides, Lb. plantarum, Lb. brevis, Lb. confuses, Ped. pentosaceus</i>	Indonesia	Puspito and Fleet (1985)
<i>Sinnamani</i>	Radish	Acidic, sour, wet	LAB	Nepal	Tamang (2010a)
<i>Soibum</i>	Bamboo shoot	Acidic, sour, soft, curry	<i>Lb. plantarum, Lb. brevis, Lb. coryniformis, Lb. delbrueckii, Leuc. fallax, Leuc. Lact. lactis, Leuc. mesenteroides, Ent. durans, Strep. lactis, B. subtilis, B. lichniformis, B. coagulans, B. cereus, B. pumilus Pseudomonas fluorescens, Saccharomyces sp., Torulopsis sp.</i>	India	Jeyaram et al. (2009), Tamang et al. (2012)
<i>Soidon</i>	Bamboo shoot tips	Acidic, sour, soft, curry	<i>Lb. brevis, Leuc. fallax, Lact. lactis</i>	India	Tamang et al. (2008)
<i>Soijim</i>	Bamboo shoot	Acidic, liquid, condiment	<i>Lb. brevis, Lb. plantarum, Lb. curvatus, Ped. pentosaceus, Leuc. mesenteroides, Leuc. fallax, Leuc. lactis, Leuc. citreum, Ent. durans</i>	India	Tamang et al. (2008)
<i>Sinki</i>	Radish tap-root	Acidic, sour, dry, soup, pickle	<i>Lb. plantarum, Lb. brevis, Lb. casei, Leuc. fallax</i>	India, Nepal, Bhutan	Tamang and Sarkar (1993), Tamang et al. (2005)
<i>Suan-cai</i>	Vegetables	Acidic, sour, wet	<i>Ped. pentosaceus, Tetrigenococcus halophilus</i>	China	Chen et al. (2006)

<i>Suan-tsai</i>	Mustard	Acidic, sour, dry	<i>Ent. faecalis</i> , <i>Lb. alimentarius</i> , <i>Lb. brevis</i> , <i>Lb. coryniformis</i> , <i>Lb. farciminis</i> , <i>Lb. plantarum</i> , <i>Lb. versmoldensis</i> , <i>Leuc. citreum</i> , <i>Leuc. mesenteroides</i> , <i>Leuc. pseudomesenteroides</i> , <i>Ped. pentosaceus</i> , <i>W. cibaria</i> , <i>W. paramesenteroides</i>	Taiwan	Chao et al. (2009)
<i>Sunki</i>	Turnip	Acidic, sour, wet	<i>Lb. plantarum</i> , <i>Lb. fermentum</i> , <i>Lb. delbrueckii</i> , <i>Lb. parabuchneri</i> , <i>Lb. kisonensis</i> , <i>Lb. otakiensis</i> , <i>Lb. rapi</i> , <i>Lb. sunkii</i>	Japan	Endo et al. (2008), Watanabe et al. (2009a)
<i>Takuanzuke</i>	Japanese radish, salt, sugar, <i>Shochu</i>	Pickle radish	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Leuc. mesenteroides</i> , <i>Streptococcus</i> sp., <i>Pediococcus</i> sp., yeasts	Japan	Alexandraki et al. (2013)
<i>Takanazuke</i>	Broad leaved mustard, red pepper, salt, turmeric	Vegetable pickle Takuanzuke	<i>Ped. halophilus</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i>	Japan	Alexandraki et al. (2013)
<i>Tuaitur</i>	Bamboo shoot	Solid, wet, sour, curry	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Ped. pentosaceus</i> , <i>Lc. lactis</i> , <i>Bacillus circulans</i> , <i>B. firmus</i> , <i>B. sphaericus</i> , <i>B. subtilis</i>	India	Chakrabarty et al. (2014)
<i>Tuairoi</i>	Bamboo shoot	Solid, dry, sour, curry	<i>Lb. plantarum</i> , <i>Ent. faecium</i> , <i>Ped. pentosaceus</i> , <i>Leuc. mesenteroides</i> , <i>B. laterosporus</i> , <i>B. circulans</i> , <i>B. stearothermophilus</i> , <i>B. firmus</i> , <i>B. cereus</i>	India	Chakrabarty et al. (2014)
<i>Yan-Jiang</i>	Ginger	Pickle	LAB	Taiwan	Chang et al. (2011)

from the fermented bamboo shoots of India (Jeyaram et al. 2010). *Jiang-sun* is a traditional fermented bamboo shoot food in Taiwan (Chen et al. 2010).

### 1.5.4 Fermented Legumes

Among the fermented legumes (Table 1.4), most of the products are of soybean origin. Two types of fermented soybean foods are produced: soybeans which are naturally fermented by *Bacillus* spp. (mostly *B. subtilis*) with characteristic stickiness, and soybeans which are fermented by filamentous molds (mostly *Aspergillus*, *Mucor*, *Rhizopus*). The *Bacillus*-fermented sticky soybean foods of Asia are *natto* of Japan (Kubo et al. 2011), *chungkokjang* of Korea (Shin et al. 2012), *kinema* of India, Nepal, and Bhutan (Tamang 2001), *aakhune*, *bekang*, *hawaijar*, *perayaan*, and *tungrymbai* of India (Tamang et al. 2009), *thua nao* of Thailand (Dajanta et al. 2011), *pepok* of Myanmar (Tamang 2010b) and *sieng* of Cambodia and Laos (Tamang 2010b). Mold-fermented soybean products are *miso* and *shoyu* of Japan (Sugawara 2010), *tempeh* of Indonesia (Nout and Kiers 2005), *douchi* of China (Zhang et al. 2007), *sufu* of China (Han et al. 2001), *doenjang* of Korea (Kim et al. 2009). Among the common non-soybean fermented legumes of the world are *bikalga*, *dawadawa*, *iru*, *soumbala*, *ugba* of Africa (Parkouda et al. 2009, Ouoba et al. 2010), *dhokla*, *papad*, and *wari* of India (Nagai and Tamang 2010), *ontjom* of Indonesia (Nagai and Tamang 2010), and *maseura* of India and Nepal (Chettri and Tamang 2008).

The species of *Bacillus* isolated from *kinema* include *B. cereus*, *B. circulans*, *B. licheniformis*, *B. sphaericus*, *B. subtilis*, and *B. thuringiensis* (Sarkar et al. 1994, 2002, Tamang 2003), however, *B. subtilis* is the dominant functional bacterium in *kinema* (Sarkar and Tamang 1994, Tamang and Nikkuni 1996). *Ent. faecium* is also present in *kinema* (Sarkar et al. 1994). Based on molecular identification tools using ARDRA, ITS-PCR, and RAPD-PCR techniques, the species of *Bacillus* were isolated from *tungrymbai* (or *turangbai*) and *bekang*, naturally fermented soybean foods of the states of Meghalaya and Mizoram in North East India and were identified as *B. subtilis*, *B. pumilus*, *B. licheniformis*, *B. cereus*, *B. coagulans*, *B. circulans*, *B. brevis*, and *Lysinibacillus fusiformis* (Chettri 2013). *B. subtilis*, *B. licheniformis*, *B. cereus*, *Staphylococcus aureus*, *Staph. sciuri*, *Alkaligenes* spp., *Providencia rettgeri* were isolated from *hawaijar* of the state of Manipur in India (Jeyaram et al. 2008a).

Species of *Bacillus* isolated from naturally fermented *chungkokjang* are *B. amyloliquefaciens*, *B. licheniformis*, and *B. subtilis* (Tamang et al. 2002, Choi et al. 2007, Kwon et al. 2009), *B. subtilis* subsp. *chungkokjang* (Park et al. 2005), *B. megaterium* (Shon et al. 2007). *Ent. faecium* is also present in *chungkokjang* (Yoon et al. 2008). Nam et al. (2012) analyzed 12,697 bacterial pyrosequences in *chungkokjang* and found that almost all the bacteria were members of the phylum *Firmicutes* (>95%), with only a small portion belonging to *Proteobacteria* (<5%). In various samples, specific unclassified *Bacillus* species and LAB existed as the dominant microbes of *chungkokjang* (Nam et al. 2012).

Japanese *natto* is the only *Bacillus*-fermented soybean food which is now produced by commercial mono-culture starter *B. natto*, first isolated from naturally fermented *natto* by Sawamura (1906). *B. natto* differs from *B. subtilis* on account of biotin requirement, production of polyglutamate, possession of 5.7-kb and 60-kb plasmids (Hara et al. 1983, Nagai et al. 1997), and insertion sequences (Nagai et al. 2000, Kimura and Itoh 2007). *Thua nao*, an ethnic fermented nonsalty sun-dried wafer-type soybean food of Thailand, is used as a seasoning. *B. subtilis* is a functional bacterium in *thua nao* (Chunhachart et al. 2006, Inatsu et al. 2006).

*Rhizopus microsporus* is a functional mold for fermentation of *tempeh* with varieties *Rhi. microsporus*, *Rhi. oligosporus*, *Rhi. rhizopodiformis*, *Rhi. tuberosus*, and *Rhi. chinensis*

**Table 1.4 Some Common and Uncommon Ethnic Fermented Legume (Soybeans and Non-Soybean) Products of the World**

<i>Product</i>	<i>Substrate/Raw Material</i>	<i>Sensory Features and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Aakhone</i>	Soybean	Alkaline, sticky, paste	<i>B. subtilis, Proteus mirabilis</i>	India	Singh et al. (2014)
<i>Bekang</i>	Soybean	Alkaline, sticky, Paste, curry	<i>B. subtilis, B. pumilus, B. licheniformis, B. sphaericus, B. brevis, B. coagulans, B. circulans, B. amyloliquefaciens, Ent. faecium, Ent. durans, Ent. hirae, Ent. Raffinossus, Ent. cecorum, Proteus mirabilis Sacch. cerevisiae, Debaryomyces hansenii, Pic. burtonii.</i>	India	Chettri (2013), Singh et al. (2014)
<i>Bhalla</i>	Black gram	Mild acidic, side dish	<i>B. subtilis, Candida curvata, C. famata, C. membranaefaciens, C. variovaarai, Cryptococcus humicoius, Deb. hansenii, Geotrichum candidum, Hansenula anomala, H. polymorpha, Kl. marxianus, Lb. fermentum, Leuc. mesenteroides, Ped. membranaefaciens, Rhiz. marina, Sacch. cerevisiae, Ent. faecalis, Trichosporon beigeli, Trichosporon pullulans, Wingea robertsii</i>	India	Rani and Soni (2007)
<i>Bikalga</i>	Roselle ( <i>Hibiscus sabdariffa</i> )	Condiment	<i>B. subtilis, B. licheniformis, B. megaterium, B. pumilus</i>	Burkina Faso	Ouoba et al. (2007a,b)
<i>Ce-lew</i>	Soybean, corn flour, rice flour, salt	Soya sauce	<i>Ped. halopholus, Bacillus sp., Asp. oryzae, Asp. flavus</i>	Thailand	Alexandraki et al. (2013)
<i>Chee-fan</i>	Soybean whey curd	Cheese-like, solid	<i>Mucor sp., Asp. glaucus</i>	China	Blandino et al. (2003)

(Continued)

**Table 1.4 (Continued) Some Common and Uncommon Ethnic Fermented Legume (Soybeans and Non-Soybean) Products of the World**

Product	Substrate/Raw Material	Sensory Features and Nature	Microorganisms	Country	References
<i>Chungkokjang</i> (or <i>jeonkukjang</i> , <i>cheonggukjang</i> )	Soybean	Alkaline, sticky, soup	<i>B. subtilis</i> , <i>B. amyloliquefaciens</i> , <i>B. licheniformis</i> , <i>B. cereus</i> , <i>Pantoea agglomerans</i> , <i>Pantoea ananatis</i> , <i>Enterococcus</i> sp., <i>Pseudomonas</i> sp., <i>Rhodococcus</i> sp.	Korea	Hong et al. (2012), Shin et al. (2012)
<i>Dage</i>	Coconut press cake, <i>Ragi</i>	Solid, side dish	<i>Rhizopus</i> sp.,	Indonesia	Alexandraki et al. (2013)
<i>Douchi</i>	Soybean	Alkaline, paste	<i>B. amyloliquefaciens</i> , <i>B. subtilis</i> , <i>Asp. oryzae</i>	China, Taiwan	Wang et al. (2006), Zhang et al. (2007)
<i>Dawadawa</i>	Locust bean	Alkaline, sticky	<i>B. pumilus</i> , <i>B. licheniformis</i> , <i>B. subtilis</i> , <i>B. firmus</i> , <i>B. atrophaeus</i> , <i>B. amyloliquefaciens</i> , <i>B. mojavensis</i> , <i>Lysinibacillus sphaericus</i> .	Ghana, Nigeria	Amoa-Awua et al. (2006), Meerak et al. (2008)
<i>Dhokla</i>	Bengal gram	Mild acidic, spongy, Steamed, snack	<i>Leuc. mesenteroides</i> , <i>Lb. fermenti</i> , <i>Ent. faecalis</i> , <i>Tor. candida</i> , <i>Tor. pullulans</i>	India	Blandino et al. (2003)
<i>Doenjang</i>	Soybean	Alkaline, paste	<i>B. subtilis</i> , <i>B. licheniformis</i> , <i>B. pumilis</i> , <i>Mu. plumbeus</i> , <i>Asp. oryzae</i> , <i>Deb. hansenii</i> , <i>Leuc. mesenteroides</i> , <i>Tor. halophilus</i> , <i>Ent. faecium</i> , <i>Lactobacillus</i> sp.	Korea	Kim et al. (2009), Shin et al. (2012)
<i>Dosa</i>	Rice, blackgram dhal ( <i>Phaseolus mango</i> )	Fermented fan cake	<i>Leuc. mesenteroides</i> , <i>Lb. delbrueckii</i> , <i>Lb. fermenti</i> , <i>Ent. faecalis</i> , <i>B. amyloliquefaciens</i> , <i>Cand. boidini</i> , <i>Cand. glabrata</i> , <i>Cand. sake</i> , <i>Deb. hansenii</i> , <i>Hansenula polymorpha</i> , <i>Issatchenkia terricola</i> , <i>Rhiz. graminis</i>	India, Sri Lanka	Soni et al. (1986)

<i>Furu</i>	Soybean curd	Mild acidic	<i>B. pumilus</i> , <i>B. megaterium</i> , <i>B. stearothersophilus</i> , <i>B. firmus</i> , <i>Staph. hominis</i>	China	Sumino et al. (2003)
<i>Gochujang</i>	Soybean, red pepper	Hot-flavored seasoning	<i>B. velezensis</i> , <i>B. amyloliquefacios</i> , <i>B. subtilis</i> , <i>B. liqueformis</i> , species of <i>Oceanobacillus</i> , <i>Zygosaccharomyces</i> , <i>Candida lactis</i> , <i>Zygorouxii</i> , <i>Aspergillus</i> , <i>Penicillium</i> , <i>Rhizopus</i>	Korea	Nam et al. (2012), Kim et al. (2013).
<i>Hawaijar</i>	Soybean	Alkaline, sticky	<i>B. subtilis</i> , <i>B. licheniformis</i> , <i>B. amyloliquefaciens</i> , <i>B. cereus</i> , <i>Staph. aureus</i> , <i>Staph. sciuri</i> , <i>Alkaligenes</i> sp., <i>Providencia rettgers</i> , <i>Proteus mirabilis</i>	India	Jeyaram et al. (2008a), Singh et al. (2014)
<i>Hishiho-Miso</i>	Soybean, barley or wheat, salt, vegetables, Mizuame, sugar, shoyu	Sweetend Miso	<i>Asp. oryzae</i> , <i>Ped. halophilus</i> , <i>Sacch. Rouxii</i> , <i>Streptococcus</i> sp.	Japan	Sugawara (2010)
<i>Iru</i>	Locust bean	Alkaline, sticky	<i>B. subtilis</i> , <i>B. pumilus</i> , <i>B. licheniformis</i> , <i>B. megaterium</i> , <i>B. fumus</i> , <i>B. atrophaeus</i> , <i>B. amyloliquefaciens</i> , <i>B. mojavensis</i> , <i>Lysinibacillus sphaericus</i> , <i>Staph. saprophyticus</i>	Nigeria, Benin	Odunfa and Oyewole (1997), Meerak et al. (2008)
<i>Kanjang</i>	Soybean, meju, salt, water	Soya sauce	<i>Asp. oryzae</i> , <i>B. subtilis</i> , <i>B. pumillus</i> , <i>B. citreus</i> , <i>Sarcina mazima</i> , <i>Sacch. Rouxii</i>	Korea	Shin et al. (2012)
<i>Kawal</i>	Leaves of legume ( <i>Cassia</i> sp.)	Alkaline, strong flavored, dried balls	<i>B. subtilis</i> , <i>propionibacterium</i> sp., <i>Lb. plantarum</i> , <i>Staph. sciuri</i> , Yeasts	Sudan	Dirar et al. (2006)
<i>Kecap</i>	Soybean, wheat	Liquid	<i>Rhiz. oligosporus</i> , <i>Rhiz. oryzae</i> , <i>Asp. oryzae</i> , <i>Ped. halophilus</i> , <i>Staphylococcus</i> sp., <i>Candida</i> sp., <i>Debaromyces</i> sp., <i>Sterigmatomyces</i> sp.	Indonesia	Alexandraki et al. (2013)

(Continued)



Table 1.4 (Continued) Some Common and Uncommon Ethnic Fermented Legume (Soybeans and Non-Soybean) Products of the World

Product	Substrate/Raw Material	Sensory Features and Nature	Microorganisms	Country	References
<i>Ketjap</i>	Soybean (black)	Syrup	<i>Asp. oryzae</i> , <i>Asp. flavus</i> , <i>Rhiz. oligosporus</i> , <i>Rhiz. arrhizus</i>	Indonesia	Alexandraki et al. (2013)
<i>Kinda</i>	Locust bean	Alkaline, sticky	<i>B. pumilus</i> , <i>B. licheniformis</i> , <i>B. subtilis</i> , <i>B. atrophaeus</i> , <i>B. amyloliquefaciens</i> , <i>B. mojavensis</i> , <i>Lysinibacillus sphaericus</i> .	Sierra Leone	Meerak et al. (2008)
<i>Kinema</i>	Soybean	Alkaline, sticky	<i>B. subtilis</i> , <i>B. licheniformis</i> , <i>B. cereus</i> , <i>B. circulans</i> , <i>B. thuringiensis</i> , <i>B. sphaericus</i> , <i>Ent. faecium</i> , <i>Cand. parapsilosis</i> , <i>Geotrichum candidum</i>	India, Nepal, Bhutan	Sarkar et al. (1994), Tamang (2003)
<i>Khaman</i>	Bengal gram	Mild acidic, spongy	<i>Leuc. mesenteroides</i> , <i>Lb. fermentum</i> , <i>Lact. lactis</i> , <i>Ped. acidilactici</i> , <i>Bacillus</i> sp.	India	Ramakrishnan (1979)
<i>Koikuchi Shoyu</i>	Defatted soybean flake, wheat, brine, tane-koji	Soy sauce	<i>Aspergillus sojae</i> , <i>Asp. oryzae</i> , <i>Sacch. rouxii</i> , <i>Tor. versatilis</i> , <i>Tor. echellsii</i> , <i>Ped. halophilus</i> , <i>Sacch. halomembransis</i> , <i>Ent. faecalis</i> , <i>Bacillus</i> sp.	Japan	Sugawara (2010)
<i>Maseura</i>	Black gram	Dry, ball-like, brittle, condiment	<i>B. subtilis</i> , <i>B. mycoides</i> , <i>B. pumilus</i> , <i>B. laterosporus</i> , <i>Ped. acidilactici</i> , <i>Ped. pentosaceus</i> , <i>Ent. durans</i> , <i>Lb. fermentum</i> , <i>Lb. salivarius</i> , <i>Sacch. cerevisiae</i> , <i>Pic. burtonii</i> , <i>Cand. castellii</i>	Nepal, India	Chettri and Tamang (2008)
<i>Meitauza</i>	Soybean	Liquid	<i>B. subtilis</i> , <i>Asp. oryzae</i> , <i>Rhiz. oligosporus</i> , <i>Mu. meitauza</i> , <i>Actinomucor elegans</i>	China, Taiwan	Zhu et al. (2008)

<i>Meju</i>	Soybean	Alkaline, paste	<i>Asp. flavus, Asp. fumigatus, Asp. niger, Asp. oryzae, Asp. retricus, Asp. spinosa, Asp. terreus, Asp. Wentii, Botrytis cinerea Mu. adundans, Mu. circinelloides, Mu. griseocyanus, Mu. hiemalis, Mu. jasseni, Mu. Racemosus, Pen. citrinum, Pen. griseopurpureum, Pen. griesotula, Pen. kaupscinskii, Pen. lanosum, Pen. thomii, Pen. turalense, Rhi. chinensis, Rhi. nigricans, Rhi. oryzae, Rhi. Sotronifer; Candida edax, Can. incommenis, Can. utilis Hansenula anomala, Han. capsulata, Han. Holstii, Rhodotorula flava, Rho. glutinis, Sacch. exiguus, Sacch. cerevisiae, Sacch. kluyveri, Zygosaccharomyces japonicus, Zyg. rouxii; Bacillus citreus, B. circulans, B. licheniformis, B. megaterium, B. mesentericus, B. subtilis, B. pumilis, Lactobacillus sp., Ped. acidilactici</i>	Korea	Choi et al. (1995)
<i>Miso</i>	Soybean	Alkaline, paste	<i>Ped. acidilactici, Leuc. paramesenteroides, Micrococcus halobius, Zygosaccharomyces rouxii, Asp. oryzae</i>	Japan	Asahara et al. (2006)
<i>Miso (Hishiho)</i>	Soybean, barley or wheat, salt, vegetables, Mizuame (dextrose syrup), sugar, shoyu	Sweet miso	<i>Asp. oryzae, Ped. halophilus, Sacch. rouxii, Streptococcus sp.</i>	Japan	Sugawara (2010)

(Continued)

Table 1.4 (Continued) Some Common and Uncommon Ethnic Fermented Legume (Soybeans and Non-Soybean) Products of the World

Product	Substrate/Raw Material	Sensory Features and Nature	Microorganisms	Country	References
Miso (Kome Ama)	Rice, soybean, salt, <i>tane-koji</i>	Sweet rice miso	<i>Asp. oryzae</i> , <i>Streptococcus</i> sp., <i>Pediococcus</i> sp., <i>Sacch. rouxii</i>	Japan	Sugawara (2010)
Miso (Kome Kara)	Rice, soybean, salt, <i>tane-koji</i> , salt	Salt rice miso	<i>Asp. oryzae</i> , <i>Sacch. rouxii</i> , <i>Ped. halophilus</i> , <i>Tor. versalis</i> , <i>Tor. echellsii</i> , <i>Bacillus</i> sp.	Japan	Sugawara (2010)
Miso (Mame)	Cereal, soybean, salt		<i>Asp. oryzae</i> , <i>Asp. sojae</i> , <i>Ent. faecalis</i> , <i>Tor. versatilis</i> , <i>Bacillus</i> sp.	Japan	Sugawara (2010)
Miso (Mugi)	Barley, soybean, salt, <i>koji</i>	Barley Miso	<i>Asp. oryzae</i> , <i>Sacch. rouxii</i> , <i>Ped. halophilus</i> , <i>Ent. faecalis</i> , <i>Tor. versatilis</i> , <i>Tor. echellsii</i> , <i>Bacillus</i> sp.	Japan	Sugawara (2010)
Moromi	Soybean		<i>Aspergillus</i> sp., <i>Sacch. rouxii</i>	Japan	FAO (1998, 1999)
Natto	Soybean	Alkaline, sticky	<i>B. subtilis</i> ( <i>natto</i> )	Japan	Nagai and Tamang (2010)
Oncom Hitam (Black Oncom) And Oncom Merah (Orange Oncom)	Peanut press cake, tapioca, soybean curd starter	Fermented peanut press cake, roasted or fried	<i>Neurospora intermedia</i> , <i>N. crassa</i> , <i>N. sitophila</i> (from red oncom), <i>Rhi. oligosporus</i> (from black oncom)	Indonesia	Ho (1986)
Ogiri/Ogili	Melon seeds, castor oil seeds, pumpkin bean, sesame		<i>B. subtilis</i> , <i>B. pumilus</i> , <i>B. licheniformis</i> , <i>B. megaterium</i> , <i>B. rimus</i> , <i>Pediococcus</i> sp., <i>Staph. saprophyticus</i> , <i>Lb. plantarum</i>	West, East and Central Africa	Odunfa and Oyewole (1997)

<i>Okpehe</i>	Seeds from <i>Prosopis africana</i>	Alkaline, sticky	<i>B. subtilis</i> , <i>B. amyloliquefaciens</i> , <i>B. cereus</i> , <i>B. licheniformis</i>	Nigeria	Oguntoyinbo et al. (2010)
<i>Owoh</i>	Cotton seed		<i>B. subtilis</i> , <i>B. pumilus</i> , <i>B. licheniformis</i> , <i>Staph. saprophyticus</i>	West Africa	Odunfa and Oyewole (1997)
<i>Papad</i>	Black gram	Circular wafers	<i>Cand. krusei</i> , <i>Deb. hansenii</i> , <i>Lb. fermentum</i> , <i>Leuc. mesenteroides</i> , <i>P. membranaefaciens</i> , <i>Sacch. cerevisiae</i> , <i>Ent. faecalis</i> , <i>Trichosporon beigelii</i>	India, Nepal	Rani and Soni (2007)
<i>Pepok</i>	Soybean	Alkaline, sticky	<i>Bacillus</i> sp.	Myanmar	Nagai and Tamang (2010)
<i>Perayaan</i>	Soybean	Alkaline, sticky	<i>B. subtilis</i> , <i>B. amyloliquefaciens</i> , <i>Vagococcus lutrae</i> , <i>Ped. acidilactici</i> , <i>Ent. faecalis</i>	India	Singh et al. (2014)
<i>Sieng</i>	Soybean	Alkaline, sticky	<i>Bacillus</i> sp.	Cambodia, Laos	Nagai and Tamang (2010)
<i>Soumbala</i>	Locust bean	Alkaline, sticky	<i>B. pumilus</i> , <i>B. atrophaeus</i> , <i>B. amyloliquefaciens</i> , <i>B. mojavensis</i> , <i>Lysinibacillus sphaericus</i> , <i>B. subtilis</i> , <i>B. thuringiensis</i> , <i>B. licheniformis</i> , <i>B. cereus</i> , <i>B.adius</i> , <i>B. firmus</i> , <i>B. megaterium</i> , <i>B. mycoides</i> , <i>B. sphaericus</i> , <i>Peaibacillus alvei</i> , <i>Peaibacillus larvae</i> , <i>Brevibacillus laterosporus</i>	Burkina Faso	Ouoba et al. (2003a,b, 2004)
Soy sauce	Soybean	Alkaline, liquid	<i>Asp. oryzae</i> , <i>Asp. niger</i> , <i>Sacch. rouxii</i> , <i>Ped. acidilactis</i> , <i>Ped. cerevisiae</i> , <i>Ped. halophilus</i> , <i>Lb. delbrueckii</i>	Worldwide	Sugawara (2010)
<i>Shoyu</i>	Soybean	Alkaline, liquid, Seasoning	<i>Asp. oryzae</i> or <i>Asp. sojae</i> , <i>Z. Rouxii</i> , <i>C. versatilis</i>	Japan, Korea, China	Sugawara (2010)

(Continued)

**Table 1.4 (Continued) Some Common and Uncommon Ethnic Fermented Legume (Soybeans and Non-Soybean) Products of the World**

<i>Product</i>	<i>Substrate/Raw Material</i>	<i>Sensory Features and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Sufu</i>	Soybean curd	Mild-acidic, soft	<i>Actinomucor elenans, Mu. silvatixus, Mu. corticolus, Mu. hiemalis, Mu. praini, Mu. racemosus, Mu. subtilissimus, Rhiz. chinensis</i>	China, Taiwan	Han et al. (2001)
<i>Tamari Shoyu</i>	Defatted soybean, salt, water, wheat	Soybean rich Shoyu	<i>Asp. sojae, Asp. oryzae, Sacch. rouxii, Tor. versatilis, Tor. echellsii, Ped. halophilus, Ent. faecalis, Bacillus sp.</i>	Japan	Alexandraki et al. (2013)
<i>Tauco</i>	Soybean	Alkaline, paste, use as flavoring agent	<i>Rhiz. oryzae, Rhiz. ologosporus, Asp. oryzae, Lb. delbrueckii, Zygosaccharomyces soyae</i>	Indonesia	Winarno et al. (1973)
<i>Tao-si</i>	Soybean, salt, rice bran, wheat flour	Fermented soybean curd	<i>Asp. oryzae</i>	Philippines	Blandino et al. (2003)
<i>Tempe</i>	Soybean	Alkaline, solid, fried cake, breakfast	<i>Asp. niger, Rhiz. oligosporus, Rhiz. arrhizus, Rhiz. oryzae, Rhiz. stolonifer, Citrobacter freundii, Enterobacter cloacae, K. pneumoniae, K. pneumoniae subsp. ozaenae, Pseudomas fluorescens as vitamin B<sub>12</sub>-producing bacteria, Lb. fermentum, Lb. lactis, Lb. plantarum, Lb. reuteri</i>	Indonesia (Origin), the Netherlands, Japan, USA	Nout and Kiers (2005), Jennessen et al. (2008)
<i>Tempe Benguk</i>	Velvet bean seeds, <i>Ragi</i> Tempe	Alkaline, solid, fried cake, breakfast	<i>Rhizopus sp., Rhiz. oligosporus, Rhiz. arrhizus</i>	Indonesia	Nagai and Tamang (2010)

<i>Tempe Gembus</i>	Solid residue of soybean curd, tapioca, <i>Ragi</i> tempe	Alkaline, solid, fried cake, breakfast	<i>Rhizopus</i> sp. <i>Rhiz. oryzae</i> , <i>Rhiz. oligosporus</i>	Indonesia	Alexandraki et al. (2013)
<i>Tempe Kecipir</i>	Winged bean seed, <i>Ragi</i> , old Tempe	Alkaline, solid, fried cake, breakfast	<i>Rhiz. oryzae</i> , <i>Rhiz. arrhizus</i> , <i>Rhiz. oligosporus</i> , <i>Rhiz. achlamydosporus</i>	Indonesia	Alexandraki et al. (2013)
<i>Tempe Kedelai</i>	Soybean, tapioca flour, maize, young papaya, cassava, coconut press cake, starter	Alkaline, solid, fried cake, breakfast	<i>Rhizopus</i> sp., <i>Rhiz. oryzae</i> , <i>Rhiz. oligosporus</i>	Indonesia	Alexandraki et al. (2013)
<i>Tempe Koro Pedang</i>	Jack bean seed ( <i>Canavalia ensiformis</i> ), <i>ragi</i> , old tempe	Alkaline, solid, fried cake, breakfast	<i>Rhiz. oryzae</i> , <i>Rhiz. arrhizus</i> , <i>Rhiz. achlamydosporus</i>	Indonesia	Alexandraki et al. (2013)
<i>Tempe Lamtoro</i>	Wied Tamarind ( <i>Leucaena Leucocephala</i> )	Alkaline, solid, fried cake, breakfast	<i>Rhizopus</i> sp., <i>Rhiz. oryzae</i>	Indonesia	Alexandraki et al. (2013)
<i>Thua nao</i>	Soybean	Alkaline, paste, dry, side dish	<i>B. subtilis</i> , <i>B. pumilus</i> , <i>Lactobacillus</i> sp.	Thailand	Chunhachart et al. (2006)

(Continued)

**Table 1.4 (Continued) Some Common and Uncommon Ethnic Fermented Legume (Soybeans and Non-Soybean) Products of the World**

<i>Product</i>	<i>Substrate/Raw Material</i>	<i>Sensory Features and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Tofu (stinky tofu)</i>	Soybean	Alkaline, liquid	<i>Bacillus</i> sp., <i>Ent. hermanniensis</i> , <i>Lb. agilis</i> , <i>Lb. brevis</i> , <i>Lb. buchneri</i> , <i>Lb. crispatus</i> , <i>Lb. curvatus</i> , <i>Lb. delbrueckii</i> , <i>Lb. farciminis</i> , <i>Lb. fermentum</i> , <i>Lb. pantheris</i> , <i>Lb. salivarius</i> , <i>Lb. vaccinostercus</i> , <i>Lc. lactis</i> , <i>Lactococcus</i> sp., <i>Leuc. camosum</i> , <i>Leuc. citreum</i> , <i>Leuc. fallax</i> , <i>Leuc. lactis</i> , <i>Leuc. mesenteroides</i> , <i>Leuc. pseudomesenteroides</i> , <i>Ped. acidilactici</i> , <i>Strep. bovis</i> , <i>Strep. macedonicus</i> , <i>W. cibaria</i> , <i>W. confusa</i> , <i>W. paramesenteroides</i> , <i>W. soli</i>	China, Japan	Chao et al. (2008)
<i>Toyo</i>	Soybean, salt, brown sugar, wheat starter	Cowpea sauce	<i>Asp. oryzae</i> , <i>Hansenula anomala</i> , <i>Hansenula subpelliculosa</i> , <i>Lb. delbrueckii</i>	Philippines	Alexandraki et al. (2013)
<i>Tungrymbai/turangbai</i>	Soybean	Alkaline, sticky, curry, soup	<i>B. subtilis</i> , <i>B. pumilus</i> , <i>B. licheniformis</i> , <i>B. amyloliquefaciens</i> , <i>Lb. brevis</i> , <i>Ent. faecium</i> , <i>Ent. durans</i> , <i>Ent. hirae</i> , <i>Ent. Raffinossus</i> , <i>Ent. cecorum</i> , <i>Vagococcus carniphilus</i> Sacch. <i>cerevisiae</i> , <i>Debaryomyces hansenii</i> , <i>Pic. burtonii</i> ,	India	Chettri (2013), Singh et al. (2014)



<i>Tuong</i>	Rice, maize, salt	Staple	<i>Asp. oryzae</i> , <i>Sacch. rouxii</i> , <i>Ped. halophilus</i>	Vietnam	Alexandraki et al. (2013)
<i>Ugba</i>	African oil bean ( <i>Pentaclethra macrophylla</i> )	Alkaline, flat, glossy, brown in color	<i>B. subtilis</i> , <i>B. pumilus</i> , <i>B. licheniformis</i> , <i>Staph. saprophyticus</i>	Nigeria	Odunfa and Oyewole (1997)
<i>Uri</i>	Locust bean	Alkaline, sticky, condiment, soup	<i>Bacillus</i> spp.	West Africa	Alexandraki et al. (2013)
<i>Usukuchi Shoyu</i>	Soybean, wheat, Tane-Koji, Amasake	Soy sauce, seasoning	<i>Asp. oryzae</i> , <i>Sacch. rouxii</i> , <i>tor. versatilis</i> , <i>Tor. echellsii</i> , <i>Ped. halophilus</i> , <i>Sacch. halomembransii</i> , <i>Ent. faecalis</i> , <i>Bacillus</i> sp.	Japan	Alexandraki et al. (2013)
<i>Vadai</i>	Black gram	Paste, side dish	<i>Pediococcus</i> sp., <i>Streptococcus</i> sp., <i>Leuconostoc</i> sp.	India	Blandino et al. (2003)
<i>Wari</i>	Black gram	Ball-like, brittle, side dish	<i>B. subtilis</i> , <i>Cand. curvata</i> , <i>Cand. famata</i> , <i>Cand. krusei</i> , <i>Cand. parapsilosis</i> , <i>Cand. vartiovaarai</i> , <i>Cryptococcus humicolus</i> , <i>Deb. hansenii</i> , <i>Deb. tamaraii</i> , <i>Geotrichum candidum</i> , <i>Hansenula anomala</i> , <i>Kl. marxianus</i> , <i>Sacch. cerevisiae</i> , <i>Rhiz. lactosa</i> , <i>Ent. faecalis</i> , <i>Wingea robetsii</i> , <i>Trichosporon beigeli</i>	India	Rani and Soni (2007)

(Nout and Kiers 2005, Jennessen et al. 2008). *Citrobacter freundii*, *Enterobacter cloacae*, *Kl. pneumoniae*, *Kl. pneumoniae* subsp. *ozaenae*, *Pseudomonas fluorescens* as vitamin B<sub>12</sub>-producing bacteria, and LAB—*Lb. fermentum*, *Lb. lactis*, *Lb. plantarum*, *Lb. reuteri* are important microorganisms in naturally fermented *tempeh* (Denter and Bisping 1994, Feng et al. 2005). Four types of *douchi* are produced in China: *Mucor*-fermented *douchi*, *Aspergillus*-fermented *douchi*, *Rhizopus*-fermented *douchi*, and *Bacillus*-fermented *douchi* (Zhang et al. 2007). *Aspergillus awamori*, *Asp. kawachii*, *Asp. oryzae*, *Asp. Shirousamii*, and *Asp. sojae* have been widely used as the starter in preparation of *koji* in Japan for production of *miso* and *shoyu* (Kitamoto 2002, Matsushita et al. 2009). *B. subtilis*, *B. licheniformis*, *Ent. faecium*, *Leuc. mesenteroides*, *Tetra. halophilus*, and *Asp. oryzae*, *Debaryomyces hansenii*, and *Mucor plumbeus* were isolated from *doenjang* (Kim et al. 2009). Using next-generation sequencing, Nam et al. (2011) have analysed the microbial community of traditional Korean soybean pastes, and derived 17,675 bacterial sequences from nine local and two commercial brands of *doenjang* samples.

*Bikalga*, *dawadawa*, *iru*, *mbodi*, *ntoba* and *soumbala* are the ethnic nonsalted fermented locust bean (*Parkia biglobosa*) foods of Africa. Microorganisms involved are *B. amyloliquefaciens*, *B. licheniformis*, *B. megaterium*, *B. pumilus*, *B. subtilis*, *Enterococcus avium*, *Ent. casseliflavus*, *Ent. faecalis*, *Ent. faecium*, *Ent. hirae*, *Lb. plantarum*, *Micrococcus* spp., *Ped. acidilactici*, *Ped. pentosaceus*, *Staphylococcus* (*Staph.*) *hominis*, *Staph. saprophyticus*, *Staph. xylosus*, *Weissella cibaria*, and *W. confusa* (Amoa-Awua et al. 2006, Azokpota et al. 2006, Meerak et al. 2008, Ouoba et al. 2010). Although *B. subtilis* has been reported as the dominant bacterium involved in the fermentation, other *Bacillus* spp. such as *B. amyloliquefaciens*, *B. cereus*, and *B. licheniformis* have also been detected, for example, in *okpehe* produced in Nigeria from *Prosopis africana* (legume) seeds, thus suggesting the inter-species diversity in these kinds of products (Oguntoyinbo et al. 2010). A study by Ouoba et al. (2008) reflects the interspecies diversity characterizing this fermentation, comprising *B. subtilis* as the predominant organism followed by *B. licheniformis*, while strains of *B. cereus*, *B. pumilus*, *B. badius*, *Brevibacillus bortelensis*, *B. sphaericus*, and *B. fusiformis* were also detected. A number of studies have been undertaken to develop starter cultures for the fermentation of indigenous legumes, such as the seeds of *Prosopis africana* (Oguntoyinbo et al. 2007), and of the African locust bean (*Parkia biglobosa*) (Ouoba et al. 2003a,b, 2004, 2005, 2007a,b).

### 1.5.5 Fermented Root Crop and Tuber Products

Cassava (*Manihot esculenta*) root is traditionally fermented into staple foods such as *gari* of Nigeria, *fufu* of Togo, Burkina Faso, Benin, Nigeria and Ghana, *agbelima* of Ghana, *chikawgue* of Zaire, *kivunde* of Tanzania, and *kocho* of Ethiopia (Table 1.5). *Gari* making involves several stages including fermentation, dextrinization, partial gelatinization, and retrogradation (Abimbola 2007). In initial stage of fermentation of cassava is dominated by *Corynebacterium manihot* (Oyewole et al. 2004). *Lb. acidophilus*, *Lb. casei*, *Lb. fermentum*, *Lb. pentosus*, *Lb. plantarum* are present in *gari* (Oguntoyinbo and Dodd 2010). *Lb. plantarum*, *Leuconostoc* sp., and *Streptococcus* spp. play a major role in detoxification of the cyanogenic glucosides during *gari* fermentation (Ngaba and Lee 1979). *Geotrichum candidum* is the dominant yeast responsible for the characteristic taste and aroma of *gari* (Okafor and Ejiofor 1990).

Cassava root is also traditionally fermented into a sweet dessert such as *tapé* in Indonesia (Ardhana and Fleet 1989). A mixed culture of *Streptococcus*, *Rhizopus*, and *Saccharomycopsis* produces the aroma in the *tapé*, whereas *Sm. fibuligera* produces  $\alpha$ -amylase and *Rhizopus* sp. produces glucoamylase (Suprianto et al. 1989). *Simal tarul ko jaanr* is a mild-alcoholic food beverage prepared from cassava root in Nepal and India (Tamang et al. 1996).

**Table 1.5 Some Ethnic Fermented Root Crop Products of the World**

<i>Product</i>	<i>Substrate/ Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Chikwangu</i>	Cassava	Solid state, staple	Species of <i>Corynebacterium</i> , <i>Bacillus</i> , <i>Lactobacillus</i> , <i>Micrococcus</i> , <i>Pseudomonas</i> , <i>Acinetobacter</i> , <i>Moraxella</i>	Central Africa, Zaire	Odunfa and Oyewole (1997)
<i>Cingwada</i>	Cassava	Solid state	Species of <i>Corynebacterium</i> , <i>Bacillus</i> ., <i>Lactobacillus</i> , <i>Micrococcus</i>	East and Central Africa	Odunfa and Oyewole (1997)
<i>Dage</i>	Coconut press cake, <i>ragi</i>		<i>Rhizopus</i> sp.	Indonesia	Alexandraki et al. (2013)
<i>Fufu</i>	Cassava	Submerged, staple	<i>Bacillus</i> sp., <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> , <i>Lb. cellobiosus</i> , <i>Lb. brevis</i> ; <i>Lb. coprophilus</i> , <i>Lact. lactis</i> ; <i>Leuc. lactis</i> , <i>Lb. bulgaricus</i> , <i>Klebsiella</i> sp., <i>Leuconostoc</i> sp., <i>Corynebacterium</i> sp., <i>Candida</i> sp.	West Africa	Odunfa and Oyewole (1997)
<i>lape Ketela</i>	Cassava, <i>ragi</i>		Species of <i>Rhizopus</i> , <i>Chlamydomucor</i> , <i>Candida</i> , <i>Saccharomyces</i> , <i>Endomycopsis</i>	Indonesia	Alexandraki et al. (2013)
<i>Gari</i>	Cassava	Solid state, staple	<i>Corynebacterium mannihot</i> , <i>Geotrichum</i> sp., <i>Lb. plantarium</i> , <i>Lb. buchneri</i> , <i>Leuconostoc</i> sp., <i>Streptococcus</i> sp.	West and Central Africa	Oyewole et al. (2004)
<i>Lafun/ Konkonte</i>	Cassava	Submerged, staple	<i>Bacillus</i> sp., <i>Klebsiella</i> sp., <i>Candida</i> sp., <i>Aspergillus</i> sp., <i>Leuc. mesenteroides</i> , <i>Corynebacterium manihot</i> , <i>Lb. plantarum</i> , <i>Micrococcus luteus</i> , <i>Geotrichum candidum</i>	West Africa	Odunfa and Oyewole (1997)

(Continued)

**Table 1.5 (Continued) Some Ethnic Fermented Root Crop Products of the World**

<i>Product</i>	<i>Substrate/ Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Peujeum</i>	Cassava roots	Acidic, solid, eaten after baking	Yeasts, mold	Indonesia	Alexandraki et al. (2013)
<i>Simal tarul to jaanr</i>	Cassava	Sweet, mild-alcoholic food, staple	<i>Mu. circinelloides</i> , <i>Rhi. Chinensis</i> ,- <i>Pichia anomala</i> , <i>Sacch. cerevisiae</i> , <i>Candida glabrata</i> , <i>Saccharomycopsis fibuligera</i> , <i>Ped. pentosaceus</i> , <i>Lb. bifermentans</i>	India and Nepal	Tamang (2005)
<i>Tapé</i>	Cassava	Sweet dessert	<i>Streptococcus</i> sp., <i>Rhizopus</i> sp., <i>Saccharomycopsis fibuligera</i>	Indonesia	Suprianto et al. (1989)
<i>Tapai Ubi</i>	Cassava, Ragi	Sweet dessert	<i>Saccharomycopsis fibuligera</i> , <i>Amylomyces rouxii</i> , <i>Mu. circinelloides</i> , <i>Mu. javanicus</i> , <i>Hansenula</i> spp., <i>Rhi. arrhizus</i> , <i>Rhi. oryzae</i> , <i>Rhi. chinensis</i>	Malaysia	Merican and Yeoh (1989)

### 1.5.6 Fermented Fruit Products

Some fermented fruit pickles are *atchara* (green unripe papaya), *burong mangga* (green unripe mango), *burong prutas* (local fruits) of the Philippines, *achar* of India and Nepal, and so on (Table 1.6). *Ca muoi* is a fermented fruit of Vietnam, and *tempoyak* is a fermented durian fruit of Malaysia. *Sacch. cerevisiae*, *Schizosaccharomyces pombe*, *Lb. plantarum*, and *Leuc. mesenteroides* were isolated from *palm winelemu*, a fermented palm fruit of West Africa (Odunfa and Oyewole 1997). *Lb. plantarum* strains were isolated from Thai fermented fruits (Tanganurat et al. 2009).

### 1.5.7 Fermented Meat Products

Traditionally preserved and fermented meat products (Table 1.7) of many countries are the *salami* of Europe (Toldra 2007), *albeira* of Portugal (Albano et al. 2009), *androlla* of Spain (Garcia-Fontan et al. 2007), *nham* of Thailand (Chokesajjawatee et al. 2009), *kargyong*, *satchu*, and *suka ko masu* of India and Nepal (Rai et al. 2009, 2010), *arjia*, *chartayshya* and *jamma* of India (Oki et al. 2011), and *nem chua* of Vietnam (Khanh et al. 2011, Nguyen et al. 2011). The major microorganisms involved in meat fermentation are a species of LAB and coagulase-negative cocci, however, yeasts and enterococci are also present in some meat products (Rantsiou and Cocolin 2006). Identification based on the culture-independent approach using the DGGE-method has revealed *Lb. curvatus* and *Lb. sakei* as the main species of LAB involved in the transformation process, accompanied by coagulase-negative cocci *Staphy. xylosus* during meat fermentation and ripening (Cocolin et al. 2011). *Lb. curvatus*, *Lb. paraplantarum*, *Lb. plantarum*, *Lb. sakei*, *Lb. brevis*, *Lb. carnis*, *Lb. casei*, *Lb. curvatus*, *Lb. divergens*, *Lb. sanfransiscensis*, *Leuc. carnosum*, *Leuc. gelidium*, *Leuc. pseudomesenteroides*, *Leuc. citreum*, *Leuc. mesenteroides*, *Ped. acidilactici*, *Ped. pentosaceus*, *W. cibaria*, *W. viridescens*, *B. lentus*, *B. licheniformis*, *B. mycoides*, *B. subtilis*, *B. thuringiensis*, *E. cecorum*, *E. durans*, *E. faecalis*, *E. faecium*, *E. hirae* are the dominant LAB in fermented meats (Albano et al. 2009, Rai et al. 2010, Cocolin et al. 2011, Oki et al. 2011, Nguyen et al. 2013a); and also coagulase-negative staphylococci, micrococci, Enterobacteriaceae in fermented meats (Marty et al. 2011). The species of yeasts present in Spanish fermented sausages are *C. intermedia/curvata*, *C. parapsilosis*, *C. zeylanoides*, *Citeromyces matritensis*, *Trichosporon ovoides*, and *Yarrowia lipolytica* (Encinas et al. 2000).

### 1.5.8 Fermented, Dried and Smoked Fish Products

Preservation of fish through fermentation, sun drying, smoking and salting is traditionally performed by people living in coastal regions, or near lakes and rivers, and such preserved/fermented fish products are consumed as seasoning, condiments, curry and on the side (Salampessy et al. 2010). Some ethnic fermented fish products of the world (Table 1.8) are *hentak*, *ngari*, and *tungtap* of India (Thapa et al. 2004), and *bordia*, *karati*, and *lashim* of India (Thapa et al. 2007), *jeotgal* or *jeot* or *saeu-jeot* of Korea (Guan et al. 2011, Jung et al. 2013), *plaa-som* of Thailand (Saithong et al. 2010), *shiokara* of Japan (Fujii et al. 1999), *patis* of the Philippines (Steinkraus 1996), *surströmming* of Sweden (Kobayashi et al. 2000a), and sun-dried or smoked fish products such as *gnuchi*, *sidra*, *sukuti* of India, Nepal, and Bhutan (Thapa et al. 2006), *Ent. faecalis*, *Lb. plantarum*, *Lb. reuteri*, *Strep. salivarius*, species of *Bacillus*, *Micrococcus*, *Pediococcus* and yeasts including species of *Candida* and *Saccharomyces* are reported from fermented fish products of Thailand (Saithong et al. 2010, Hwanhlem et al. 2011). *Micrococcus* and *Staphylococcus* are dominant bacterial genera during ripening of *shiokara* (Wu et al. 2000). *Haloanaerobium praevalens* has been

**Table 1.6 Some Ethnic Fermented Fruit Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Achar/ chatney</i>	Fruits, vegetables, oil, salt	Acidic, hot and sour, pickle	LAB	India, Nepal, Pakistan, Bangladesh	Tamang (2010b)
<i>Atchara</i>	Green unripe papaya, onion, red pepper, garlic, ginger, salt	Unripe papaya pickle	<i>Leuc. mesenteroides</i> , <i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Strep. faecalis</i> , <i>Ped. cerevisiae</i>	Philippines	Alexandraki et al. (2013)
<i>Burong Mangga</i>	Green unripe mango, salt	Pickled green mango		Philippines	Alexandraki et al. (2013)
<i>Burong Prutas</i>	Fruits, salt, sugar	Picked fruits	<i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i>	Philippines	Alexandraki et al. (2013)
<i>Chuk</i>	Fruits	Sour, dark-brown paste, therapeutic uses	unknown	Nepal, India	Tamang (2005)
<i>Ogiri</i>	Melon seeds	Alkaline, condiment	LAB, <i>Bacillus</i> spp.	Nigeria	Tamang (2010b)
<i>Owoh</i>	Cotton seeds	Alkaline, condiment	LAB, <i>Bacillus</i>	Nigeria	Tamang (2010b)
<i>Tempoyak</i>	Durian fruit	Fermented durian fruit	Yeast, <i>Bacillus</i> sp., <i>Acetobacter</i> sp., <i>Lactobacillus</i> sp.	Malaysia	Merican (1996)

**Table 1.7 Some Common and Uncommon Ethnic Fermented Meat Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Alheira</i>	Pork or beef, bread chopped fat, spices, salt	Dry/ semi-dry, sausage	LAB, staphylococci, micrococci, yeast	Portugal	Albano et al. (2009)
<i>Androlla</i>	Ground lean pork	Dry, sausage	LAB, micrococci, yeast	Spain	Garcia-Fontan et al. (2007)
<i>Arjia</i>	Large intestine of chevon	Sausage, curry	<i>Ent. faecalis</i> , <i>Ent. faecium</i> , <i>Ent. hirae</i> , <i>Leuc. citreum</i> , <i>Leuc. mesenteroides</i> , <i>Ped. pentosaceus</i> , <i>Weissella cibaria</i>	India, Nepal	Oki et al. (2011)
Bacon	Cured pork, beef	Dry, semi-dry, staple	LAB, yeast, micrococci	Germany, Belgium, Spain	Tanaka et al. (1980)
<i>Chartayshya</i>	Chevon	Dried, smoked meat, curry	<i>Ent. faecalis</i> , <i>Ent. faecium</i> , <i>Ent. hirae</i> , <i>Leuc. citreum</i> , <i>Leuc. mesenteroides</i> , <i>Ped. pentosaceus</i> , <i>Weissella cibaria</i>	India	Oki et al. (2012)
<i>Chorizo</i>	Pork, coarse chopped, spices, salt	Dry	<i>Lb. sake</i> , <i>Lb. curvatus</i> , <i>Lb. plantarum</i>	Spain	Garcia-Varona et al. (2000)
<i>Chilu</i>	Yak, beef, sheep fat	Hard, oily, edible oil	LAB	India, China (Tibet), Bhutan	Tamang et al. (2009)
Ham	Cured pork	Semi-dry, breakfast	LAB, yeasts, micrococci	Spain, Italy	Simoncini et al. (2007)
<i>Honoheigrain</i>	Pig/boar meat	Rough, hard, dried meat; curry	<i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> , <i>E. faecium</i> , <i>B. cereus</i> , <i>B. pumilus</i> , <i>B. firmus</i> , <i>B. circulans</i> , <i>B. stearothermophilus</i> , <i>Micrococcus</i> , <i>Staphylococcus</i> ; <i>Debaryomyces hansenii</i> , <i>Sacch. cerevisiae</i>	India	Chakrabarty et al. (2014)

(Continued)



**Table 1.7 (Continued) Some Common and Uncommon Ethnic Fermented Meat Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Kargyong</i>	Yak, beef, pork, crushed garlic, ginger, salt	Sausage like meat product, curry	<i>Lb. sakei, Lb. divergens, Lb. carnis, Lb. sanfrancensis, Lb. curvatus, Leuc. mesenteroides, Ent. faecium, B. subtilis, B. mycoides, B. thuringiensis, Staph. aureus, Micrococcus sp., Deb. hansenii, Pic. anomala</i>	India	Rai et al. (2010)
<i>Kheuri</i>	Yak, beef	Chopped intestine of yak, curry	LAB	India, China (Tibet), Bhutan	Tamang et al. (2009)
Jerky	Beef	Dry, semi-dry, side dish	LAB, yeast, molds, micrococci	South America	Delong (1992)
<i>Longanisa</i>	Pork lean, pork backfat, ground pork, salt, sugar, soysauce, vinegar, anisado wine, potassium nitrate		LAB	Philippines	Alexandraki et al. (2013)
<i>Mortadello</i>	Pork	Unsmoked chopped meat, sausage	LAB, micrococci	Italy, France, USA	
<i>Nham (Musom)</i>	Pork meat, pork skin, salt, rice, garlic	Fermented pork	<i>Ped. cerevisiae, Lb. plantarum, Lb. brevis</i>	Thailand	Chokesajjawatee et al. (2009)

<i>Nem-chua</i>	Pork, salt, cooked rice	Fermented sausage	<i>Lb. pentosus</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. paracasei</i> , <i>Lb. fermentum</i> , <i>Lb. acidipiscis</i> , <i>Lb. farciminis</i> , <i>Lb. rossiae</i> , <i>Lb. fuchuensis</i> , <i>Lb. namurensis</i> , <i>Lc. lactis</i> , <i>Leuc. citreum</i> , <i>Leuc. fallax</i> , <i>Ped. acidilactici</i> , <i>Ped. pentosaceus</i> , <i>Ped. stilesii</i> , <i>Weissella cibaria</i> , <i>W. paramesenteroides</i> .	Vietnam	Nguyen et al. (2011)
<i>Pastirma</i>	Chopped beef lean meat with lamb fat, not smoked, heavily seasoned	Dry/semi-dry, sausage	<i>Lb. plantarum</i> , <i>Lb. sake</i> , <i>Pediococcus</i> , <i>Micrococcus</i> , <i>Staph. xylosum</i> , <i>Staph. carnosus</i>	Turkey, Iraq	Aksu et al. (2005)
<i>Peperoni</i>	Pork, beef	Dried meat, smoked, sausage	LAB, micrococci	Europe, America, Australia	Adams (2010)
<i>Salami</i>	Pork	Sausage	LAB, micrococci	Europe	Adams (2010)
<i>Sai-krok-prieo</i>	Pork, rice	Sausage	LAB	Thailand	Phithakpol et al. (1995)
<i>Soppressata</i>	Chopped lean pork meat, NaCl and spices	Dry/ semi-dry, sausage	LAB, yeast, staphylococci, micrococci, enterobacteriaceae	Italy	Parente et al. (1994)
<i>Salchichon</i>	Pork or beef meat, fat, NaCl, spices	Dry, sausage	LAB, Yeast, micrococcaceae, enterobacteriaceae, molds	Spain	Fernandez-Lopez et al. (2008)
<i>Salsiccia</i>	Chopped pork meat, spices, NaCl	Dry/ semi-dry, sausage	LAB, yeast, enterobacteriaceae staphylococci, micrococci	Italy	Parente et al. (2001a,b)
<i>Sucuk</i>	Chopped meat, pork or beef, curing salts and various spices	Dry, sausage	LAB, micrococci, staphylococci, enterobacteriaceae	Turkey	Genccelep et al. (2008)

(Continued)

**Table 1.7 (Continued) Some Common and Uncommon Ethnic Fermented Meat Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Sai-krok-prieo</i>	Pork, rice, garlic, salt	Fermented sausage	<i>Lb. plantarum</i> , <i>Lb. salivarius</i> , <i>Ped. pentosacuns</i>	Thailand	Adams (2010)
<i>Satchu</i>	Beef, yak, port, tumeric powder, edible oil, butter, salt	Ethnic dried meat, curry	<i>Ped. pentosaceuous</i> , <i>Lb. casei</i> , <i>Lb. carnis</i> , <i>Ent. faecium</i> , <i>B. subtilis</i> , <i>B. mycoides</i> , <i>B. lentus</i> , <i>Staph. aureus</i> , <i>Micrococcus</i> sp., <i>Deb. hansenii</i> , <i>Pic. anomala</i>	India	Tamang et al. (2012)
<i>Suka ko masu</i>	Goat, buffalo meat, tumeric powder, mustard oil, salt	Dried or smoked meat, curry	<i>Lb. carnis</i> , <i>Ent. faecium</i> , <i>Lb. plantarum</i> , <i>B. subtilis</i> , <i>B. mycoides</i> , <i>B. thuringiensis</i> , <i>Staph. aureus</i> , <i>Micrococcus</i> sp., <i>Debaromyces hansenii</i> , <i>Pic. burtonii</i>	India	Rai et al. (2010)
<i>Sukula</i>	Buffalo	Dried, smoked, curry	LAB	Nepal	Tamang (2010a)
<i>Tocino</i>	Pork, salt, sugar, potassium nitrate	Fermented cured pork	<i>Ped. cerevisiae</i> , <i>Lb. brevis</i> , <i>Leuc. mesenteroides</i>	Philippines	Alexandraki et al. (2013)

**Table 1.8 Some Common and Uncommon Ethnic Fermented Fish Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Ayaiba</i>	Fish	Smoked fish, pickle, curry	unknown	India	Tamang (2010a)
<i>Bagoong Alamang (Bagoong Isda, Bagoong)</i>	Fish/shrimp, salt	Fish/shrimp paste, condiment	<i>Bacillus</i> sp., <i>Pediococcus</i> sp.	Philippines	Alexandraki et al. (2013)
<i>Balao-balao (Burong Hipon Tagbilao)</i>	Shrimp, rice, salt.	Fermented rice shrimp, condiment	<i>Leuc. mesenteroides</i> , <i>Ped. cerevisiae</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Ent. faecalis</i>	Philippines	Alexandraki et al. (2013)
<i>Balao-balao (Burong Hipon Tagbieao)</i>	Shrimp, rice, salt	Fermented fish-rice, condiment	<i>Leuc. mesenteroides</i> , <i>Ped. cerevisiae</i>	Philippines	Arroyo et al. (1978)
<i>Belacan (Blacan)</i>	Shrimp, salt	Shrimp paste, condiment	<i>Bacillus</i> , <i>Pediococcus</i> , <i>Lactobacillus</i> , <i>Micrococcus</i> , <i>Sarcina</i> , <i>Clostridium</i> , <i>Brevibacterium</i> , <i>Flavobacterium</i> , <i>Corynebacteria</i>	Malaysia	Salampessy et al. (2010)
<i>Bagoong</i>	Fish	Fish paste; condiment	<i>Bacillus</i> sp., <i>Micrococcus</i> sp., <i>Moraxella</i> sp.	Philippines	Mabesa and Babaan (1993)
<i>Bagoong alamang</i>	Shrimp	Shrimp paste; condiment	LAB	Philippines	Mabesa and Babaan (1993)
<i>Bakasang</i>	Fish, shrimp	Fish or shrimp paste, condiment	<i>Pseudomonas</i> , <i>Enterobacter</i> , <i>Moraxella</i> , <i>Micrococcus</i> , <i>Streptococcus</i> , <i>Lactobacillus</i> , <i>Pseudomonas</i> , <i>Moraxella</i> , <i>Staphylococcus</i> , <i>Pediococcus</i> spp.	Indonesia	Ijong and Ohta (1996)

(Continued)

**Table 1.8 (Continued) Some Common and Uncommon Ethnic Fermented Fish Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Burong Bangus</i>	Milkfish, rice, salt, vinegar	Fermented milkfish, sauce	<i>Leuc. mesenteroides, Lb. plantarum, W. confusus</i>	Philippines	Alexandraki et al. (2013)
<i>Burong Isda</i>	Fish, rice, salt	Fermented fish, sauce	<i>Leuc. mesenteroides, Ped. cerevisiae, Lb. plantarum, Strep. faecalis, Micrococcus sp.</i>	Philippines	Sakai et al. (1983a,b)
<i>Budu</i>	Marine fishes, salt, sugar	Fish sauce	<i>Ped. halophilus, Staph. aureus, Staph. epidermidis, B. subtilis, B. laterosporus, Proteus sp., Micrococcus sp., Sarcina sp., Corynebacterium sp.</i>	Thailand, Malaysia	Merican (1977), Phithakpol et al. (1995)
<i>Gnuchi</i>	Fish ( <i>Schizothorax richardsonii</i> ), salt, tumeric powder	Eat as curry	<i>Lb. plantarum, Lact. lactis, Leuc. mesenteroides, Ent. faecium, Ent. faecalis, Ped. pentosaceus, Cand. chiropterorum, Cand. bombicola, Saccharomycopsis sp.</i>	India	Tamang et al. (2012)
<i>Gulbi</i>	Shell-fish	Salted and dried, side dish	<i>Bacillus licheniformis, Staphylococcus sp., Aspergillus sp., Candida sp.</i>	Korea	Kim et al. (1993)
<i>Hákarl</i>	Shark flesh	Fermented, side dish	LAB	Iceland	Alexandraki et al. (2013)
<i>Hentak</i>	Finger sized fish ( <i>Esomus danricus</i> )	Condiment	<i>Lact. lactis, Lb. plantarum, Lb. fructosus, Lb. amylophilus, Lb. coryniformis, Ent. faecium, B. subtilis, B. pumilus, Micrococcus sp., Candida sp., Saccharomycopsis sp.</i>	India	Thapa et al. (2004)

<i>Hoi-malaeng pu-dong</i>	Mussel ( <i>Mytilus smaragdinus</i> ), salt	Fermented mussel	<i>Ped. halophilus</i> , <i>Staph. aureus</i> , <i>Staph. epidermidis</i>	Thailand	Phithakpol et al. (1995)
<i>Ika-Shiokara</i>	Squid, salt	Fermented squid	<i>Micrococcus</i> sp., <i>Staphylococcus</i> sp., <i>Debaryomyces</i> sp.	Japan	Alexandraki et al. (2013)
<i>Jaadi</i>	Fish, salt	Salted fish, curry, condiment	LAB	Sri Lanka	Alexandraki et al. (2013)
<i>Jeot kal</i>	Fish	High-salt fermented, staple	LAB	Korea	Guan et al. (2011)
<i>Kapi</i>	Small fish	Paste, condiment	Micrococci, LAB	Thailand	Phithakpol (1993)
<i>Karati, Bordia, Lashim</i>	Fish ( <i>Gudushia chapra</i> , <i>Pseudeutropius atherinoides</i> , <i>Cirrhinus reba</i> ), salt	Dried, salted, side dish	<i>Lact. lactis</i> , <i>Leuc. mesenteroides</i> , <i>Lb. plantarum</i> , <i>B. subtilis</i> , <i>B. pumilus</i> , <i>Candida</i> sp.	India	Thapa et al. (2007)
<i>Kung-chom</i>	Shrimp, salt, sweetened rice	Fermented fish-rice	<i>Ped. cerevisiae</i>	Thailand	Phithakpol (1993)
<i>Kung chom</i>	Shrimp ( <i>Macrobrachum lanchesteri</i> ), salt, garlic, rice	Fermented shrimp	<i>Ped. halophilus</i> , <i>Staph. aureus</i> , <i>Staph. epidermidis</i>	Thailand	Phithakpol (1993)
<i>Kusaya</i>	Horse mackerel, salt	Fermented dried fish	<i>Corynebacterium kusaya</i> , <i>Spirillum</i> sp., <i>C. bifermentans</i> , <i>Penicillium</i> sp.	Japan	Alexandraki et al. (2013)
<i>Mehiawah</i>	Marine fish	Fermented paste; side-dish		Middle-East	Al-Jedah et al. (1999)

(Continued)

**Table 1.8 (Continued) Some Common and Uncommon Ethnic Fermented Fish Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Myulchijeot</i>	Small sardine, salt	Fermented small sardine	<i>Ped. cerevisiae</i> , <i>Staphylococcus</i> sp., <i>Bacillus</i> sp., <i>Micrococcus</i> sp.	Korea	Alexandraki et al. (2013)
<i>Narezushi</i>	Sea water fish, cooked millet, salt	Fermented fish-rice	<i>Leuc. mesenteroides</i> , <i>Lb. plantarum</i>	Japan	Alexandraki et al. (2013)
<i>Nam pla</i> ( <i>Nampla-dee</i> , <i>Nampla-sod</i> )	<i>Solephorus</i> sp., <i>Ristelliger</i> sp., <i>Cirrhinus</i> sp., water, brackish water, marine fish, salt	Fish sauce	<i>Micrococcus</i> sp., <i>Pediococcus</i> sp., <i>Staphylococcus</i> sp., <i>Streptococcus</i> sp., <i>Sarcina</i> sp., <i>Bacillus</i> sp., <i>Lactobacillus</i> sp., <i>Corynebacterium</i> sp., <i>Pseudomonas</i> sp., <i>Halococcus</i> sp., <i>Halobacterium</i> sp.	Thailand	Saisithi (1987), Wongkhalaung (2004)
<i>Ngari</i>	Fish ( <i>puntius sophore</i> ), salt	Fermented fish	<i>Lact. lactis</i> , <i>Lb. plantarum</i> , <i>Ent. faecium</i> , <i>Lb. fructosus</i> , <i>Lb. amylophilus</i> , <i>Lb. coryniformis</i> , <i>B. subtilis</i> , <i>B. pumilus</i> , <i>Micrococcus</i> sp., <i>Candida</i> sp., <i>Saccharomycopsis</i> sp.	India	Thapa et al. (2004)
<i>Nga pi</i>	Fish	Fermented paste, condiment	LAB	Myanmar	Tyn (1993)
<i>Ngan pyaye</i>	Fish	Fish sauce, condiment	LAB	Myanmar	Tyn (1993)
<i>Nuoc mam</i>	Marine fish	Fish sauce, condiment	<i>Bacillus</i> sp., <i>Pseudomonas</i> sp., <i>Micrococcus</i> sp., <i>Staphylococcus</i> sp., <i>Halococcus</i> sp., <i>Halobacterium salinarium</i> , <i>H. cutirubrum</i>	Vietnam	Lopetcharat et al. (2001)



<i>Patis</i>	<i>Stolephorus</i> sp., <i>Clupea</i> sp., <i>Decapterus</i> sp., <i>Leionathus</i> sp., fish, salt, food color-optional	Fish sauce	<i>Ped. halophilus</i> , <i>Micrococcus</i> sp., <i>Halobacterium</i> sp., <i>Halococcus</i> sp., <i>Bacillus</i> sp.	Philippines, Indonesia	Baens-Arega (1977)
<i>Pla-ra</i>	Fresh water fish, salt, roasted rice	Fermented fish-rice	<i>Pediococcus</i> sp.	Thailand	Phithakpol (1993)
<i>Pla-chao</i> ( <i>Pla-Khaomak</i> )	Fresh water fish, salt, Khaomak	Thai sweetened fish	<i>Ped. cerevisiae</i> , <i>Staphylococcus</i> sp., <i>Bacillus</i> sp., <i>Micrococcus</i> sp.	Thailand	Phithakpol et al. (1995)
<i>Pla-chom</i> ( <i>Pla-khoa-kour</i> )	Fresh water or marine anchovy, boiled rice, salt, garlic, roasted rice flour	Fermented fish, Thai anchovy	<i>Ped. cerevisiae</i> , <i>Lb. brevis</i> , <i>Bacillus</i> sp.	Thailand	Phithakpol et al. (1995)
<i>Pekasam</i>	Freshwater fish-rice	Fermented fish, side dish	LAB	Malaysia	Karim (1993)
<i>Pindang</i>	Fish	Dried, salted, side dish	LAB	Indonesia	Putro (1993)
<i>Pla-paeng-daeng</i>	Marine fish, red molds rice ( <i>Ang- kak</i> ), salt	Red fermented fish	<i>Pediococcus</i> sp., <i>Ped. halophilus</i> , <i>Staph. aureus</i> , <i>Staph. epidermidis</i> ,	Thailand	Phithakpol et al. (1995)
<i>Pla ra</i> ( <i>Pla-dag</i> , <i>Pla-ha</i> , <i>Ra</i> )	Fresh water fish, brackish water fish, marine fish	Fermented fish	<i>Ped. cerevisiae</i> , <i>Lb. brevis</i> , <i>Staphylococcus</i> sp., <i>Bacillus</i> sp.	Thailand	Phithakpol et al. (1995)

(Continued)

**Table 1.8 (Continued) Some Common and Uncommon Ethnic Fermented Fish Products of the World**

<i>Product</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Pla-som (Pla-khao-sug)</i>	Marine fish, salt, boiled rice, garlic	Fermented fish	<i>Ped. cerevisiae</i> , <i>Lb. brevis</i> , <i>Staphylococcus</i> sp., <i>Bacillus</i> sp.	Thailand	Saithong et al. (2010)
<i>Saeoo Jeot (Jeotkal)</i>	Shrimp ( <i>Acetes chinensis</i> ), salt	Fermented shrimp	<i>Halobacterium</i> sp., <i>Pediococcus</i> sp.	Korea	Guan et al. (2011)
<i>Som-fug (Som-dog, Pla-fu, Pla-muig, Fug-som)</i>	Fresh fish, boiled rice, salt, garlic	Thai fermented fish, condiment	<i>Ped. cerevisiae</i> , <i>Lb. brevis</i> , <i>Staphylococcus</i> sp., <i>Bacillus</i> sp.	Thailand	Phithakpol et al. (1995)
<i>Shottsuru</i>	Anchovy, opossum shrimp, salt	Fish sauce, condiment	<i>Halobacterium</i> sp., <i>Aerococcus viridians (Ped. homari)</i> , halotolerant and halophilic yeasts	Japan	Itoh et al. (1993)
<i>Sidra</i>	Fish ( <i>Punitus sarana</i> )	Dried fish, curry	<i>Lact. lactis</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> , <i>Ent. faecium</i> , <i>Ent. faecalis</i> , <i>Ped. pentosaceus</i> , <i>W. confusus</i> , <i>Cand. chiropterorum</i> , <i>Cand. bombicola</i> , <i>Saccharomycopsis</i> sp.	India	Thapa et al. (2006)
<i>Sikhae</i>	Sea water fish, cooked millet, salt	Fermented fish-rice, sauce	<i>Leuc. mesenteroides</i> , <i>Lb. plantarum</i>	Korea	Lee (1993)
<i>Shiokara</i>	Squid	Fermented; side-dish	LAB	Japan	Fujii et al. (1999)

<i>Suka ko maacha</i>	River fish ( <i>Schizothorax richardsoni</i> ), salt, turmeric powder	Smoked, dried, curry	<i>Lact. lactis</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> , <i>Ent. faecium</i> , <i>Ent. faecalis</i> , <i>Ped. pentosaceus</i> , <i>Cand. chiropterorum</i> , <i>Cand. bombicola</i> , <i>Saccharomycopsis</i> sp.	India	Thapa et al. (2006)
<i>Sukuti</i>	Fish ( <i>Harpodon nehereus</i> )	Pickle, soup, and curry	<i>Lact. lactis</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> , <i>Ent. faecium</i> , <i>Ent. faecalis</i> , <i>Ped. pentosaceus</i> , <i>Cand. chiropterorum</i> , <i>Cand. bombicola</i> , <i>Saccharomycopsis</i> sp.	India	Thapa et al. (2006)
<i>Surströmming</i>	Fish	Fermented herrings	<i>Haloanaerobium praevalens</i>	Sweden	Kobayashi et al. (2000a)
<i>Tai-pla</i>	Fresh water fish, brackish water fish, marine fishes, salt	Fermented fish, condiment	<i>Pediococcus</i> sp., <i>Ped. halophilus</i> , <i>Staph. aureus</i> , <i>Staph. epidermidis</i>	Thailand	Phithakpol et al. (1995)
<i>Trassi</i>	Shrimps/fish	Fermented paste; side-dish	LAB, micrococci	Indonesia	Van Veen (1965)
<i>Tungtap</i>	Fish	Fermented fish, paste, pickle	<i>Lc. lactis</i> subsp. <i>cremoris</i> , <i>Lc. plantarum</i> , <i>Ent. faecium</i> , <i>Lb. fructosus</i> , <i>Lb. amylophilus</i> , <i>Lb. coryniformis</i> subsp. <i>Torquens</i> , <i>Lb. plantarum</i> , <i>B. subtilis</i> , <i>B. pumilus</i> , <i>Micrococcus</i> , yeasts-species of <i>Candida</i> , <i>Saccharomycopsis</i> .	India	Thapa et al. (2004)
<i>Yu lu</i>	Small fish like sardine or anchovies	Fish sauce	LAB, micrococci	China	Jiang et al. (2007)

reported from *surströmming*, the fermented herrings of Sweden and *Haloanaerobium fermentans*, *Tetra. muriaticus* and *Tetra. halophilus* from the Japanese puffer fish ovaries (Kobayashi et al. 2000b,c). *B. subtilis*, *B. pumilus*, *E. faecalis*, *E. faecium*, *Lc. lactis* subsp. *cremoris*, *Lc. lactis* subsp. *lactis*, *Lc. plantarum*, *Lb. amylophilus*, *Lb. fructosus*, *Lb. confusus*, *Lb. coryniformis* subsp. *torquens*, *Lb. plantarum*, *Leuc. mesenteroides*, *P. pentosaceus*, *Micrococcus*; yeasts—*Candida bombicola*, *C. chiropterorum*, and *Saccharomycopsis* spp. were isolated from Indian fermented and sun-dried fish products (Thapa et al. 2004, 2006, 2007).

### 1.5.9 Vinegar

Vinegar (Table 1.9) is one of the most popular condiments in the world and is prepared from any sugar containing substrates and hydrolyzed starchy materials by acetic acid fermentation. *Acetobacter aceti* subsp. *aceti*, *A. oryzae*, *A. pasteurianus*, *A. polyxygenes*, *A. xylinum*, *A. malorum*, *A. pomorum* are the dominant bacteria for vinegar fermentation (Haruta et al. 2006, Bourdichon et al. 2012). Yeast species in vinegar fermentation are *Candida lactis-condensi*, *C. stellata*, *Hanseniaspora valbyensis*, *H. osmophila*, *Saccharomycodes ludwigii*, *Sac. cerevisiae*, *Zygosaccharomyces bailii*, *Z. bisporus*, *Z. lentus*, *Z. mellis*, *Z. pseudorouxii*, and *Z. rouxii* (Solieri and Giudici 2008).

### 1.5.10 Ethnic Fermented Tea

Tea, the second most popular beverage in the world after water, originated in China and two common species of tea are *Camellia sinensis* var. *sinensis* and *Camellia sinensis* var. *assamica* (Schillinger et al. 2010). Though normal black tea is drunk everywhere, however, some ethnic Asian communities have special fermented tea such as *miang* of Thailand, *puer* tea and *fuzhuan brick* of China, and *kombucha*. Fermented tea (Table 1.10) is *puer tea*, *kombucha*, and *fuzuan brick* tea of China (Mo et al. 2008), and *miang* of Thailand (Tanasupawat et al. 2007).

*Aspergillus niger* is the predominant fungus in *puer tea*; *Blastobotrys adeninivorans*, *A. glaucus*, species of *Penicillium*, *Rhizopus* and *Saccharomyces*, and the bacterial species *Actinoplanes* and *Streptomyces* were also isolated from *puer tea* (Jeng et al. 2007, Abe et al. 2008). *Brettanomyces bruxelensis*, *Candida stellata*, *Rhodotorula mucilaginosa*, *Saccharomyces* spp., *Schizosaccharomyces pombe*, *Torulasporea delbrueckii*, *Zygosaccharomyces bailii*, *Z. bisporus*, *Z. kombuchaensis*, and *Z. microellipsoides* were isolated from *kombucha* (Kurtzman et al. 2001, Teoh et al. 2004). The major bacterial genera present in *kombucha* were *Gluconacetobacter* (>85%), *Acetobacter* (<2%), *Lactobacillus* (up to 30%), and the yeast populations were found to be dominated by *Zygosaccharomyces* (>95%) (Marsh et al. 2014). *Lb. thailandensis*, *Lb. camelliae*, *Lb. plantarum*, *Lb. pentosus*, *Lb. vaccinostercus*, *Lb. pantheris*, *Lb. fermentum*, *Lb. suebicus*, *Ped. siamensis*, *E. casseliflavus*, and *E. camelliae* are involved in the fermentation of *miang* production (Sukontasing et al. 2007, Tanasupawat et al. 2007). Species of *Aspergillus*, *Penicillium* and *Eurotium* are major fungi for fermentation of *fuzhuan brick* tea (Mo et al. 2008).

### 1.5.11 Bacterial Cellulose

*Nata* or bacterial cellulose (Table 1.10) produced by *Acetobacter xylinum* is a candied delicacy of the Philippines (Kozaki 1976, Jagannath et al. 2010, Adams 2014). Two types of *nata* are well known: *nata de piña*, produced from the juice from pineapple trimmings, and *nata de coco*, produced from coconut water or coconut skim milk. Species identification using 16S rDNA sequencing revealed that the two strains belong to two different species of

**Table 1.9 Some Common and Uncommon Vinegar Products of the World**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Cuka Aren</i>	Sap from flower stalk of Aren	Vinegar, seasoning	<i>Acetobacter</i> sp.	Indonesia	Alexandraki et al. (2013)
<i>Cuka Nipah</i>	Sap from inflorescence stalk of Nipa fruiticans	Vinegar, seasoning	<i>Acetobacter</i> sp.	Malaysia	Alexandraki et al. (2013)
<i>Sirca</i>	Gur of molasses or grains	Vinegar, seasoning	<i>Sacch. cerevisiae, Acetobacter</i> sp.	Pakistan	Alexandraki et al. (2013)
<i>Sirka</i>	Fruit juices or sugar cane juices	Vinegar, seasoning	<i>Acetobacter</i> sp.	Bangladesh	Alexandraki et al. (2013)
<i>Suka</i>	Coconut water or fruits or sugar or palm sap or rice washings	Vinegar, seasoning	<i>Lb. fermentum, Lb. plantarum, Lb. panis, Lb. pontis, W. cibaria, Acetobacter pomoum, Actobacter ghanensis, Acetobacter orientalis, Acetobacter pasteurianus</i>	Philippines	Dalmacio et al. (2011)
<i>Vinegar</i>	Sugar containing substrates	Acetic acid flavored, liquid, condiment, seasoning	<i>Acetobacter aceti</i> subsp. <i>aceti, A. oryzae, A. pasteurianus, A. polyxygenes, A. xylinum, A. malorum, A. pomorum; Candida lactis-condensi, C. stellata, Hanseniaspora valbyensis, H. osmophila, Saccharomycodes ludwigii, Sacch. cerevisiae, Zygosaccharomyces bailii, Z. bisporus, Z. lentus, Z. mellis, Z. pseudorouxii, Z. rouxii</i>	Worldwide	Solieri and Giudici (2008), Sengum and Karabiyikli (2011)

**Table 1.10 Some Miscellaneous Fermented Products of the World**

<i>Fermented Products</i>	<i>Substrate/Raw Materials</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<b>Fermented Tea</b>					
<i>Fuzhuan brick</i>	Tea	Fermented tea, drink	<i>Aspergillus, Penicillium, Eurotium</i>	China	Mo et al. (2008)
<i>Kombucha or Tea fungus</i>	Tea liquor	Flavored, drink	<i>Acetobacter xylinum, Zygosaccharomyces kombuchaensis, Z. bailii, Z. bisporus, Z. microellipsoides, Brettanomyces, Saccharomyces, Schizosaccharomyces pombe, Torulaspora delbrueckii, Rhodotorula mucilaginosa, Candida stellata, Brettanomyces bruxellensis</i>	China (Tibet), India	Schillinger et al. (2010)
<i>Miang</i>	Tea	Fermented tea, flavored, drink	<i>Lb. thailandensis, Lb. camelliae, Lb. plantarum, Lb. pentosus, Lb. vaccinostercus, Lb. pantheris, Lb. fermentum, Lb. suebicus, Pediococcus siamensis, E. casseliflavus and E. camelliae</i>	Thailand	Tanasupawat et al. (2007)
<i>Puer</i>	Tea	Fermented tea, brownish red, and a fragrance produced, drink	<i>Asp. glaucus, species of Penicillium, and Rhizopus, Blastobotrys adeninivorans, and Saccharomyces Actinoplanes, Streptomyces</i>	China	Jeng et al. (2007), Abe et al. (2008)
<b>Bacterial Cellulose</b>					
<i>Nata de coco</i>	Coconut water or coconut skim milk	Thick white or cream-colored, candied, ice cream, fruit salads	<i>Acetobacter xylinus</i> and <i>A. hansenii</i>	Philippines	Bernardo et al. (1998)

<i>Nata de piña</i>	Juice from pineapple	Insoluble gelatinous film of polysaccharides, ice cream, fruit salads	<i>Acetobacter xylinus</i> and <i>A. hansenii</i>	Philippines	Bernardo et al. (1998)
<b>Chocolate</b>					
<i>Cacao</i>	Cacao beans in pods of tree <i>Theobroma cacao</i>	Chocolate, confectionery	<i>Lb. fermentum</i> , <i>Acetobacter pasteurianus</i> , <i>A. senegalensis</i> , <i>Lb. ghanensis</i> , <i>Lb. plantarum</i> , <i>Lb. cacaonum</i> , <i>Lb. fabifermentans</i> , <i>Weissella fabaria</i> , <i>W. ghanensi</i> , <i>Fructobacillus pseudoficulneus</i> , <i>Tatumella ptyseos</i> , <i>Tatumella citrea</i> , <i>Bacillus coagulans</i> ; <i>Hanseniaspora uvarum</i> , <i>H. quilliermundii</i> , <i>Issatchenkia orientalis</i> ( <i>Candida krusei</i> ), <i>Pichia membranifaciens</i> , <i>Sacch. cerevisiae</i> , <i>Kluyveromyces</i> sp.	Worldwide	Papalexandratou et al. (2011)
<b>Coffee</b>					
<i>Coffee</i>	Coffee	Flavored coffee, refreshing drink	<i>Ent. cloacae</i> , <i>Klebsiella oxytoca</i> , <i>Hafnia alvei</i> , <i>Lactobacillus</i> , <i>Leuconostoc</i> , <i>Weissella</i> spp.	Worldwide	Holzappel and Müller (2007), Schillinger et al. (2008)
<b>Fermented egg</b>					
<i>Pidan</i>	Duck egg	Alkaline, side dish	<i>B. cereus</i> , <i>B. macerans</i> , <i>Staphylococcus cohnii</i> , <i>Staph. epidermidis</i> , <i>Staph. haemolyticus</i> , <i>Staph. warneri</i>	China	Wang and Fung (1996)

*Acetobacter*: *A. xylinus* and *A. hansenii* and may be a new subspecies under these species designation (Bernardo et al. 1998). Bacterial cellulose, a microbial polysaccharide, has significant potential as a food ingredient in view of its high purity, *in situ* change of flavor and color, and its ability to form various shapes and textures (Shi et al. 2014).

### 1.5.12 Cocoa/Chocolates

Chocolate (Table 1.10) is also a fermented product obtained from cocoa beans which require fermentation as one of the first stages. *Lb. fermentum* and *Acetobacter pasteurianus* were the predominating bacterial species during cocoa fermentation (Lefeber et al. 2010, Papalexandratou et al. 2011). Diverse LAB species appear to be typically associated with the fermentation of cocoa beans in Ghana, and, in fact, a number of new species have been described in recent years, for example, *Lb. ghanensis* (Nielsen et al. 2007), *Weissella ghanensis* (de Bruyne et al. 2008), *Lb. cacaonum*, *Lb. fabifermentans* (de Bruyne et al. 2009) and *Weissella fabaria* (de Bruyne et al. 2010). *Fructobacillus pseudoficulneus*, *Lb. plantarum*, and *Acetobacter senegalensis* were among the prevailing species during the initial phase of cocoa fermentation, and *Tatumella ptyseos* and *Tatumella citrea* were the prevailing enterobacterial species in the beginning of the fermentation (Papalexandratou et al. 2011). *Bacillus coagulans* are also recovered in vinegar (Bourdichon et al. 2012). Yeasts involved during spontaneous cocoa fermentation are *Hanseniaspora uvarum*, *H. quilliermundii*, *Issatchenkia orientalis* (*Candida krusei*), *Pichia membranifaciens*, *Sacch. cerevisiae*, and the *Kluyveromyces* species for flavor development (Ardhana and Fleet 1989).

### 1.5.13 Coffee Cherries

Coffee cherries are harvested from *Coffea arabica* trees and are processed by either wet or dry methods to remove the pulp and mucilaginous materials that surround the seeds (Silva et al. 2008). Species of yeasts and bacteria grow throughout these processes, producing pectinolytic, hemicellulolytic, and other enzymes that facilitate pulp and mucilage degradation (Masoud et al. 2004). *Ent. cloacae*, *Klebsiella oxytoca*, and *Hafnia alvei* have been isolated from coffee berries of Ethiopia (Holzapfel and Müller 2007), while LAB represented by *Lactobacillus*, *Leuconostoc*, and *Weissella* spp. seem to be typically associated with desirable fermentations of the coffee cherry (Schillinger et al. 2008).

### 1.5.14 Fermented Eggs

*Pidan* (Table 1.10), consumed by the Chinese, are preserved eggs prepared from alkali-treated fresh duck eggs, which have a strong smell of hydrogen sulfide and ammonia (Ganaseen and Bejakul 2010). The main alkaline chemical reagent used for making *pidan* is sodium hydroxide, which is produced by the reaction of sodium carbonate, water, and calcium oxide of pickle or coating mud. *B. cereus*, *B. macerans*, *Staphy. cohnii*, *Staph. epidermidis*, *Staph. haemolyticus*, and *Staph. warneri* are predominant in *pidan* (Wang and Fung 1996).

## 1.6 Types of Fermented Beverages

### 1.6.1 Amylolytic Mixed Starters

The production of amylolytic starters (Table 1.11) are a unique traditional technology of preservation of the essential consortia of microorganisms (filamentous molds, amylolytic and



**Table 1.11 Some Ethnic Amylolytic Starter Cultures of Asia**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Balan</i>	Wheat	Dry, ball-like starter	Molds, yeasts	India	Tamang (2010a)
<i>Bakhar</i>	Rice-herbs	Dry, ball-like starter	Yeasts	India	Tamang (2010a)
<i>Budod</i>	Rice, starter	Basi production	<i>Mu. circinelloides</i> , <i>Mu. griseocyanus</i> , <i>Rhi. cohnii</i> , <i>Sacch. cerevisiae</i> , <i>Saccharomycopsis fibuligera</i>	Philippines	Kozaki and Uchimura (1990)
<i>Binubudan (Binuburan, Purad)</i>	Milled rice + Budod	Basi production	<i>Debarymyces hansenni</i> , <i>Cand. parapsilosis</i> , <i>Trichosporon fennicum</i>	Philippines	Tanimura et al. (1978)
<i>Binokhok</i>	Roast rice	Starter	Unknown	Philippines	Alexandraki et al. (2013)
<i>Chiu-yueh</i>	Rice, wild herbs	Gray-white, dry ball to preparae <i>lao-chao</i>	Species of <i>Rhizopus</i> , <i>Amylomyces</i> , <i>Torulopsis</i> , <i>Hansenula</i>	China, Taiwan, Singapore	Wei and Jong (1983)
<i>Chou or chu or shi or qu</i>	Rice, wheat, sorghum or barley flour	Dry, ball, cake or brick shaped	Species of <i>Aspergillus</i> , <i>Candida</i> , <i>Weissella</i> , <i>Staphylococcus</i>	China	Yan et al. (2013)
<i>Chuzo</i>	Rice, wild herbs	Dry, ball-like starter	Molds, yeasts, LAB	Mongolia	Alexandraki et al. (2013)
<i>Dabai</i>	Rice, wild herbs	Dry, ball-like starter	Molds, yeasts, LAB	India	Sha et al. (2012)

(Continued)

Table 1.11 (Continued) Some Ethnic Amylolytic Starter Cultures of Asia

Product	Substrate	Sensory Property and Nature	Microorganisms	Country	References
<i>Hamei</i>	Rice	Starter to make <i>atingbai</i>	<i>Mucor</i> sp., <i>Rhizopus</i> sp., <i>Sacch. cerevisiae</i> , <i>Pic. anomala</i>	India	Jeyaram et al. (2008b)
<i>Hong qu (hóng qu)</i>	Rice	Starter to make Chinese wine	<i>Saccharomycopsis fibuligera</i> , <i>Sacch. cerevisiae</i> , <i>Pichia</i> , <i>Candida</i> , <i>Cryptococcus</i> , <i>Rhodotorula</i> , <i>Sporobolomyces</i> , <i>Rhodospiridium</i>	China	Lv et al. (2013)
<i>Humao</i>	Rice, barks of wild plants	Dry, flat, cake-like starter for <i>Judima</i> production	<i>Ped. Pentosaceous</i> , <i>B. polymyxa</i> , <i>B. licheniformis</i> , <i>B. stearothersophilus</i> , <i>D. hansenii</i> , <i>Sacch. cerevisiae</i> , <i>Rhizopus</i> , <i>Mucor</i>	India	Chakrabarty et al. (2014)
<i>Jui paing</i>	Rice, wild herbs	Dry, ball-like starter to prepare <i>tapai</i>	Molds, yeasts, LAB	Malaysia	Alexandraki et al. (2013)
<i>Ipoh/Siye</i>	Rice, wild herbs	Dry, mixed starter	Molds, yeasts, LAB	India	Tamang (2010a)
<i>Koji</i>	Rice, wheat	Dry, black-yellow colored, mold-culture to produce <i>saké</i> , <i>miso</i> , <i>shoyu</i>	<i>Asp. awamori</i> , <i>Asp. kawachii</i> , <i>Asp. oryzae</i> , <i>Asp. shirousamii</i> , <i>Asp. Sojae</i> , yeasts	Japan	Lee et al. (2007), Suganuma et al. (2007)
<i>Khekhrii</i>	Germinated rice	Dry starter to make <i>Zutho</i>	Yeasts, LAB	India	Tamang (2010a)
<i>Loogpang</i>	Rice flour, powder of Kha root, spices	Starter, cake	<i>Rhizopus</i> sp., <i>Mucor</i> sp., <i>Chlamydomucor</i> sp., <i>Penicillium</i> sp., <i>Aspergillus</i> sp., <i>Asp. niger</i> , <i>Asp. flavus</i> , <i>Endomycopsis</i> sp., <i>Hansenula</i> sp., <i>Saccharomyces</i> sp.	Thailand	Tamang (2012b)

<i>Marcha</i>	Glutinous rice, roots, wild herbs, ginger, red dry chili	Starter	<i>Mu. circinelloides</i> , <i>Mu. hiemalis</i> , <i>Rhiz. chinensis</i> , <i>Rhiz. stolonifer</i> , <i>Saccharomycopsis fibuligera</i> , <i>Saccharomycopsis capsularis</i> , <i>Pichia anomala</i> , <i>Pichia burtonii</i> , <i>Sacch. cerevisiae</i> , <i>Sacch. bayanus</i> , <i>Cand. glabrata</i> , <i>Ped. pentosaceus</i> , <i>Lb. bif fermentans</i> , <i>Lb. brevis</i>	India	Tsuyoshi et al. (2005)
<i>Mana</i>	Wheat, herbs	Dry, granulated starter to produce alcoholic drinks	<i>Asp. oryzae</i> , <i>Rhizopus</i> spp.	Nepal	Nikkuni et al. (1996)
<i>Manapu</i>	Rice-wheat, herbs	Dry, mixed starter to produce <i>poko</i>	Molds, yeasts	Nepal	Shrestha et al. (2002)
<i>Men</i>	Rice, wild herbs, spices	Dry, ball-like starter to produce <i>Ruou</i>	<i>Rhi. oryzae</i> , <i>Rhi. microsporus</i> , <i>Absidia corymbifera</i> , <i>Amylomyces rouxii</i> , <i>Saccharomycopsis fibuligera</i> , <i>Sacch. cerevisiae</i> , <i>Issatchenkia</i> sp., <i>Pic. anomala</i> , <i>Pic. ranongensis</i> , <i>Candida tropicalis</i> , <i>Clavisporea lusitaniae</i> , <i>Xeromyces bisporus</i> , <i>Botryobasidium subcoronatum</i> ; <i>Ped. pentosaceus</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Weissella confusa</i> , <i>W. paramesenteroides</i> , <i>Bacillus subtilis</i> , <i>B. circulans</i> , <i>B. amyloliquefaciens</i> , <i>B. sporothermodurans</i> , <i>Acetobacter orientalis</i> , <i>A. pasteurianus</i> , <i>Burkholderia ubonensis</i> , <i>Ralstonia solanacearum</i> , <i>Pelomonas puraquae</i>	Vietnam	Dung et al. (2006), Thanh et al. (2008)
<i>Meju</i>	Soybean	Fermented soybean starter	<i>Asp. oryzae</i> , <i>B. subtilis</i>	Korea	Alexandraki et al. (2013)

(Continued)

**Table 1.11 (Continued) Some Ethnic Amylolytic Starter Cultures of Asia**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Nuruk (Kokja)</i>	Wheat	Starter to produce <i>Takju</i> , <i>sojo</i> , <i>yakju</i>	<i>Asp. oryzae</i> , <i>Candida</i> sp., <i>Asp. niger</i> , <i>Rhizopus</i> sp., <i>Penicillium</i> sp., <i>Mucor</i> sp., <i>Hansenula anomala</i> , <i>Leuc. mesenteroides</i> , <i>B. subtilis</i>	Korea	Jung et al. (2012)
<i>Phab</i>	Wheat, wild herbs	Dry, mixed starter to produce <i>chyang</i>	Molds, yeasts, LAB	India, China (Tibet), Bhutan, Nepal	Tamang (2010a)
<i>Poo</i>	Rice, herbs	Dry, mixed starter to produce <i>chyang</i>	Molds, yeasts, LAB	Bhutan	Tamang (2012b)
<i>Ragi</i>	Rice flour, spices	Starter to produce <i>tape</i>	<i>Rhizopus</i> , <i>Mucor</i> , <i>Amylomyces rouxii</i> , <i>Aspergillus</i> , <i>Saccharomycopsis</i> , <i>Candida parapsilosis</i> , <i>C. melinii</i> , <i>C. lactosa</i> , <i>C. pelliculosa</i> , <i>Sacch. cerevisiae</i> , <i>Hansenula subpelliculosa</i> , <i>H. anomala</i> and <i>H. malanga</i> , <i>Enterococcus faecalis</i> , <i>Lb. plantarum</i> and <i>Ped. pentosaceus</i> , <i>B. coagulans</i> , <i>B. brevis</i> , <i>B. stearrowthermophilus</i>	Indonesia	Hesseltine et al. (1988), Hesseltine and Ray (1988)

<i>Samac</i>	Sugar cane	Basi production	Yeast, bacteria, molds	Philippines	Alexandraki et al. (2013)
<i>Thiat</i>	Rice-herbs	Dry, mixed starter to produce <i>kiad-lieh</i>	Molds, yeasts, LAB	India	Tamang (2010a)
<i>Torami</i>	Rice	Fermented rice gruel	<i>Hansenlu anomala</i> , <i>Candida guilliermondii</i> , <i>C. tropicalis</i> , <i>Geotrichium candidum</i>	India	Batra and Millner (1974)
<i>Yao Qu</i> (yào qu)	Rice	Starter to make Chinese wine	<i>Saccharomycopsis fibuligera</i> , <i>Sacch. Cerevisiae</i> , <i>Pichia</i> , <i>Candida</i> , <i>Cryptococcus</i> , <i>Rhodotorula</i> , <i>Sporobolomyces</i> , <i>Rhodosporidium</i>	China	Lv et al. (2013)

alcohol-producing yeasts, and LAB) with rice or wheat as the base in the form of dry, flat or round balls, for the production of alcoholic beverages in South East Asia including the Himalayan regions of India, Nepal, and Bhutan (Tamang 2010a). Three types of dry, amylolytic and mixed cultures or inocula are traditionally used in Asia as starters to convert cereal starch to sugars and subsequently to alcohol and organic acids (Tamang and Fleet 2009):

1. A consortium of mycelial or filamentous molds, amylolytic and alcohol-producing yeasts and LAB with rice or wheat as the base in the form of dry, flat or round balls of various sizes. The starter is inoculated with a previous starter. This mixed flora is allowed to develop for a short time, then dried, and used to make either alcohol or fermented foods from starchy materials, for example, *marcha*, *ragi*, *bubod*, *loogpang*, *nuruk*, *men*, and so on (Tamang 2010c), which are used as starters for a number of fermentations based on rice and cassava or other cereals in Asia (Table 1.12).
2. A combination of *Aspergillus oryzae* and *A. sojae* is used in the form of a starter called *koji* in Japan to produce alcoholic beverages including *saké*. *Koji* also produces amylases that convert starch into fermentable sugars, which are then used for the second stage yeast fermentation to make nonalcoholic fermented soybean products called *miso* and *shoyu*, while proteases are formed to break down the soybean protein.
3. Whole-wheat flour is moistened and made into large compact cakes, which are incubated to culture yeasts and filamentous molds and are used to ferment starchy material to produce alcohol in China.

Asian amylolytic starters have different vernacular names such as *marcha* in India and Nepal, *hamei*, *humao*, *phab* in India (Tamang et al. 1996, 2012; Shrestha et al. 2002), *mana* and *manapu* in Nepal (Nikkuni et al. 1996), *men* in Vietnam (Dung et al. 2007), *ragi* in Indonesia (Uchimura et al. 1991), *bubod* in the Philippines (Hesseltine and Kurtzman 1990), *chiu/chu* in China and Taiwan (Steinkraus 1996), *loogpang* in Thailand (Vachanavinich et al. 1994), and *nuruk* in Korea (Steinkraus 1996).

Microbial profiles of Indian amylolytic starters are filamentous molds—*Mucor circinelloides* forma *circinelloides*, *Mu. hiemalis*, *Rhi. chinensis*, and *Rhi. stolonifer* variety *lyococcus* (Tamang et al. 1988), yeasts—*Sacch. cerevisiae*, *Sacch. bayanus*, *Saccharomycopsis (Sm.) fibuligera*, *Sm. capsularis*, *Pichia anomala*, *Pic. burtonii*, and *Candida glabrata* (Tamang and Sarkar 1995, Tsuyoshi et al. 2005, Tamang et al. 2007, Jeyaram et al. 2008b, 2011), and LAB—*Ped. pentosaceus*, *Lb. bifermentans*, and *Lb. brevis* (Hesseltine and Ray 1988, Tamang and Sarkar 1995, Tamang et al. 2007). Microorganisms in *men* of Vietnam include amylase producers (*Rhi. oryzae*, *Rhi. microsporus*, *Absidia corymbifera*, *Amylomyces rouxii*, *Saccharomycopsis fibuligera*), ethanol producers (*Sacch. cerevisiae*, *Issatchenkia* sp., *Pic. anomala*, *Pic. ranongensis*, *Candida tropicalis*, *Clavispora lusitaniae*), yeasts contaminants (*Xeromyces bisporus*, *Botryobasidium subcoronatum*); LAB (*Ped. pentosaceus*, *Lb. plantarum*, *Lb. brevis*, *Weissella confusa*, *W. paramesenteroides*), amylase-producing bacilli (*Bacillus subtilis*, *B. circulans*, *B. amyloliquefaciens*, *B. sporothermodurans*), acetic acid bacteria (*Acetobacter orientalis*, *A. pasteurianus*), and environmental contaminants (*Burkholderia ubonensis*, *Ralstonia solanacearum*, *Pelomonas puraquae*) (Dung et al. 2006, 2007, Thanh et al. 2008).

A combination of *Asp. oryzae* and *Asp. sojae* are used in *koji* in Japan to produce alcoholic beverages including *saké* (Zhu and Trampe 2013). *Koji* (Chinese *chu*, *shi*, or *qu*) also produces amylases that convert starch into fermentable sugars, which are then used for the second stage yeast fermentation to make nonalcoholic fermented soybean *miso* and *shoyu* (Sugawara 2010). *Asp. awamori*, *Asp. kawachii*, *Asp. oryzae*, *Asp. shirousamii*, and *Asp. sojae* have been widely used as the

**Table 1.12 Some Common and Uncommon Ethnic Fermented Beverages and Alcohol Drinks of the World**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Aara</i>	Cereals	Clear distilled liquor	Unknown	India	Tamang (2010a)
<i>Aarak</i>	Barley, millet, <i>phab</i>	Distilled from <i>chyang</i> , clear liquor	Unknown	India, China (Tibet), Bhutan	Tamang (2010a)
<i>Arrakku</i>	Palm sap, sugar	Palm wine	<i>Sacch. cerevisiae</i>	Sri Lanka	
<i>Atingba</i>	Rice, <i>hamei</i>	Mild-alcoholic, sweet-sour		India	Jeyaram et al. (2009)
<i>Apong</i>	Rice, <i>phab</i>	Mild-alcoholic		India	Chakrabarty et al. (2014)
<i>Bantu beer</i>	Sorghum, millet	Opaque appearance, sour flavor, beer	LAB, yeasts	South Africa	Kutyauripo et al. (2009)
<i>Basi</i>	Sugar cane, <i>bubod</i>	Clear or cloudy liquid	Yeasts, molds	Philippines	Tanimura et al. (1978)
<i>Bhaati Jaanr</i>	Glutinous rice, <i>marcha</i>	Fermented rice beverage	<i>Mu. circinelloides</i> , <i>Rhiz. chinensis</i> , <i>Sm. fibuligera</i> , <i>Pic. anomala</i> , <i>Sacch. cerevisiae</i> , <i>Cand. glabrata</i> , <i>Ped. pentosaceus</i> , <i>Lb. bifermentans</i>	India	Tamang and Thapa (2006)
<i>Bhang-chyang</i>	Maize-rice/barley, <i>phab</i>	Extract of <i>mingri</i> , alcoholic beverage	Yeasts, molds	India	Thakur
<i>Brem</i>	Glutinous rice, ragi	Alcoholic drink	<i>Sacch. cerevisiae</i> , <i>Cand. glabrata</i> , <i>Cand. parapsilosis</i> , <i>P. anomala</i> , <i>Issatchenkia orientalis</i> , <i>Lactobacillus</i> sp., <i>Acetobacter</i> sp. <i>Mu. indicus</i> ,	Indonesia	Sujaya et al. (2004)

(Continued)

**Table 1.12 (Continued) Some Common and Uncommon Ethnic Fermented Beverages and Alcohol Drinks of the World**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Bouza</i>	Wheat, malt	Alcoholic thin gruel, alcoholic drink	LAB	Egypt	Steinkraus (1996)
<i>Boza</i>	Wheat, rye, millet, maize	Cooked slurry, food beverage	LAB, yeasts	Bulgaria, Romania, Turkey, Albania	Steinkraus (1996)
<i>Bussa</i>	Maize, sorghum, finger millet	Alcoholic thin gruel, refreshing drink	Yeasts, LAB	Kenya	Steinkraus (1996)
<i>Bushera</i>	Sorghum, millet	Slurry, food beverage	Yeasts, LAB	Uganda	Steinkraus (1996)
<i>Bupju</i>	Rice, glutinous rice, water, starter (Nuruk)	Alcoholic beverage	<i>Saccharomyces</i> sp.	Korea	Jung et al. (2012)
<i>Cauim</i>	Cassava, rice, peanuts, pumpkin, cotton seed, maize	Alcoholic beverage	<i>Lb. plantarum</i> , <i>Lb. fermentum</i> , <i>Lb. paracasei</i> , <i>Lb. brevis</i> ; <i>Pic. guilliermondii</i> , <i>K. lactis</i> , <i>Candida</i> sp, <i>Rhi. toruloides</i> , <i>Sacch. cerevisiae</i>	Brazil	Ramos et al. (2010)
<i>Chyang/Chee</i>	Finger millet/barley, <i>phab</i>	Mild-alcoholic, slightly sweet-acidic		China (Tibet), Bhutan, Nepal, India	Tamang (2010a)
<i>Chicha</i>	Maize, human saliva	Alcoholic drink	<i>Sacch. cerevisiae</i> , <i>Sacch. apiculata</i> , <i>Sacch. pastorianus</i> , species of <i>Lactobacillus</i> , <i>Acetobacter</i>	Peru	Vallejo et al. (2013)
<i>Chulli</i>	Apricot	Filtrate, clear, alcoholic drink	Yeast	India	Thakur et al. (2004)



<i>Daru</i>	Cereal	Alcoholic beverages; filtrate, jiggery	Yeast, LAB	India	Thakur et al. (2004)
<i>Emu</i>	Palm	Palm wine, submerged	<i>Sacch. cerevisiae</i> , <i>Schizosaccharomyces pombe</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i>	West Africa	Odufa and Oyewole (1997)
<i>Ennog</i>	Rice, paddy husk	Black rice beer	Yeast, LAB	India	Tamang (2010a)
<i>Ewhaju</i>	Rice, <i>nuruk</i>	Nondistilled, filtered and clarified, clear liquor	Yeast, LAB	Korea	Jung et al. (2012)
<i>Faapar ko jaanr</i>	Buck wheat, <i>marcha</i>	Mild-acidic, alcoholic beverage		India, Nepal	Tamang (2010a)
<i>Feni</i>	Cashew apple	Distilled wine from cashew apples, strong flavor	<i>Sacch. cerevisiae</i>	World-wide	Tamang (2010c)
<i>Gahoon ko jaanr</i>	Wheat, <i>marcha</i>	Mild-acidic, alcoholic beverage		India, Nepal	Tamang (2010a)
<i>Haria</i>	Rice, <i>dabai</i>	Alcoholic beverage	<i>Sacch. cerevisiae</i> , <i>Sacch. boulardii</i> , <i>Zygosaccharomyces cidri</i> , <i>Candida tropicalis</i> , <i>C. musae</i> , <i>C. nitratophila</i> , <i>Issatchenkia</i> sp., <i>Pediococcus</i> sp., <i>Lactobacillus</i> sp.	India	Sha et al. (2012)
<i>Jao ko jaanr</i>	Barley, <i>marcha</i>	Mild-acidic, alcoholic beverage	Yeast, molds, <i>Pediococcus</i>	India, Nepal	Tamang (2010a)
<i>Jou</i>	Rice	Mild-alcoholic beverage	Yeasts, LAB	India	Tamang (2010a)

(Continued)

**Table 1.12 (Continued) Some Common and Uncommon Ethnic Fermented Beverages and Alcohol Drinks of the World**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Kaffir beer</i> (same as <i>Bantu beer</i> )	Sorghum, millet	Opaque appearance, sour flavor, beer	LAB, yeasts	South Africa	Tamang (2010c)
<i>Kanji</i>	Carrot/beet roots, <i>Torani</i>	Strong flavored, alcoholic drink	<i>Hansenlu anomala</i> , <i>Candida guilliermondii</i> , <i>C. tropicalis</i> , <i>Geotrichium candidum</i> , <i>Leuc. mesenteroides</i> , <i>Pediococcus</i> spp., <i>Lb. dextranicum</i> , <i>Lb. paraplantarum</i> , <i>Lb. pentosus</i>	India	Sura et al. (2001), Kingston et al. (2010)
<i>Khao maak</i>	Rice, <i>loogpang</i>	Juicy, white colored, sweet taste, mild alcoholic dessert	LAB, yeasts	Thailand	Phithakpol et al. (1995)
<i>Kiad lieh</i>	Rice, <i>thiat</i>	Distilled liquor, clear, alcoholic drink	LAB, yeasts	India	Tamang (2010a)
<i>Krachae</i>	Rice, <i>loogpang</i>	Nondistilled and filtered liquor	Molds, yeasts	Thailand	Vachanavinich et al. (1994)
<i>Kodo Ko Jaanr</i>	Millet	Alcoholic liquor	<i>Mu. circinelloides</i> , <i>Rhiz. Chinensis</i> , <i>Sm. fibuligera</i> , <i>P. anomala</i> , <i>Sacch. cerevisiae</i> , <i>Cand. glabrata</i> , <i>Ped. pentosaceus</i> , <i>Lb. bifermentans</i>	India, Nepal	Thapa and Tamang (2004)
<i>Judima</i>	Rice	Alcoholic beverage	<i>Ped. pentosaceous</i> , <i>B. circulans</i> , <i>B. laterosporus</i> , <i>B. pumilus</i> , <i>B. firmus</i> ; <i>D. hansenii</i> , <i>Sacch. cerevisiae</i>	India	Chakrabarty et al. (2014)
<i>Lohpani</i>	Maize-rice, barley	Alcoholic liquor, beverage	Unknown	India	Tamang (2010a)

<i>Lugri</i>	Barley, <i>phab</i>	Sweet-sour, mild alcoholic, thick liquid, alcoholic beverage	Yeasts, molds	India, China (Tibet)	Thakur et al. (2004)
<i>Madhu</i>	Rice	Distilled liquor, alcoholic drink	Yeasts, molds	India	Tamang (2010a)
<i>Makai ko jaanr</i>	Maize, <i>marcha</i>	Mild-alcoholic, sweet-sour beverage	Yeast, molds, <i>Pediococcus</i>	India, Nepal	Tamang (2010a)
<i>Makgeolli</i>	Rice, <i>nuruk</i>	Mild-alcoholic, sweet-sour beverage	Family-Saccharomycetaceae $\gamma$ -Proteobacteria to Firmicutes	Korea	Jung et al. (2012)
<i>Mangisi</i>	Maize	Liquor, alcoholic drink	Yeast, LAB	Zimbabwe	Tamang (2010c)
<i>Mbege</i>	Malted millet	Acidic, mild-alcoholic, drink	Yeast, LAB	Tanzania	Tamang (2010c)
<i>Merrisa</i>	Millet, cassava	Turbid drink, beer	Yeasts, LAB	Sudan	Tamang (2010c)
<i>Mingri</i>	Maize-rice/barley	Sweet, mild alcoholic, thick, beverage	<i>Phab</i>	India	Thakur et al. (2004)
<i>Nam khao</i>	Rice, <i>loogpang</i>	Distilled liquor, alcoholic drink		Thailand	Phithakpol et al. (1995)
<i>Nareli</i>	Coconut palm	Sweet, milky, effervescent, mild alcoholic beverage	Yeasts, LAB	India	Tamang (2010a)
<i>Nchiangne</i>	Red rice, <i>khekhrii</i>	Distilled liquor, alcoholic drink	Yeasts, LAB	India	Tamang (2010a)
<i>Oh</i>	Rice-millet	Soft, mild-alcoholic beverage	Unknown	India	Tamang (2010a)
<i>Ou</i>	Rice, <i>loogpang</i>	Distilled liquor, alcoholic drink	Yeasts, LAB	Thailand	Phithakpol et al. (1995)

(Continued)

**Table 1.12 (Continued) Some Common and Uncommon Ethnic Fermented Beverages and Alcohol Drinks of the World**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Toddy/tari</i>	Palm sap	Palm wine, sweet, milky, effervescent and mild alcoholic beverage	<i>Sacch. cerevisiae</i> , <i>Schizosaccharomyces pombe</i> , <i>Acetobacter aceti</i> , <i>A. rancens</i> , <i>A. suboxydans</i> , <i>Leuc. dextranicum</i> , <i>Micrococcus</i> sp., <i>Pediococcus</i> sp., <i>Bacillus</i> sp., <i>Sarcina</i> sp.	India	Shamala and Sreekantiah (1988)
<i>Poko</i>	Rice, <i>manapu</i>	Sweet-acidic, mild-alcoholic beverage	Yeasts, molds, LAB	Nepal	Shrestha et al. (2002)
<i>Pona</i>	Rice	Mild-alcoholic, sweet-sour, paste	Molds, yeast, LAB	India	Tamang (2010a)
<i>Pulque</i>	Agave juice	White, viscous, acidic-alcoholic, refreshing drink	<i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lb. acetotolerans</i> , <i>Lb. acidophilus</i> , <i>Lb. hilgardii</i> , <i>Lb. kefir</i> , <i>Lb. plantarum</i> , <i>Leuc. citreum</i> , <i>Leuc. kimchi</i> , <i>Leuc. mesenteroides</i> , <i>Leuc. pseudomesenteroides</i> , <i>Erwinia rhapontici</i> , <i>Enterobacter</i> spp., <i>Acinetobacter radioresistens</i> , <i>Zymomonas mobilis</i> , <i>Acetobacter malorum</i> , <i>A. pomorium</i> , <i>Microbacterium arborescens</i> , <i>Flavobacterium johnsoniae</i> , <i>Gluconobacter oxydans</i> , <i>Hafnia alvei</i> , <i>Sacch. bayanus</i> , <i>Sacch. cerevisiae</i> , <i>Sacch. paradoxus</i> , <i>Candida</i> spp., <i>C. parapsilosis</i> , <i>Clavispora lusitaniae</i> , <i>Hanseniaspora uvarum</i> , <i>Kl. lactis</i> , <i>Kl. marxianus</i> , <i>Pic. membranifaciens</i> , <i>Pic. spp.</i> , <i>Torulaspora delbrueckii</i>	Mexico	Escalante et al. (2008), Lappe-Oliveras et al. (2008)

<i>Raksi</i>	Cereals, <i>marcha</i>	Clear distilled liquor , alcoholic drink	Molds, LAB	India, Nepal	Kozaki et al. (2000)
<i>Ruou de</i>	Rice, <i>men</i>	Distilled liquor, clear, alcoholic drink	Molds, LAB	Vietnam	Dung (2004)
<i>Ruou nep</i>	Rice, <i>men</i>	Distilled liquor, clear, alcoholic drink	Molds, LAB	Vietnam	Dung (2004)
<i>Ruou nep than</i>	Rice (purple), <i>men</i>	Nondistilled, viscous, thick food beverage	Molds, LAB	Vietnam	Dung (2004)
<i>Ruou nep chan</i>	Rice, maize, cassava, <i>men</i>	Nondistilled, viscous, thick; or distilled, alcoholic drink	Molds, LAB	Vietnam	Dung (2004)
<i>Ruhi</i>	Rice	Distilled liquor, alcoholic beverage	Yeasts	India	Tamang (2010a)
<i>Roselle wine</i>	Roselle fruit, water, sugar	Roselle wine	<i>Sacch. ellipsoideus</i> var. <i>montrachet</i>	Philippine	Tamang (2010c)
<i>Saké</i>	Polished rice, glucose, <i>koji</i>	Nondistilled, clarified, and filtered liquor, alcoholic drink	<i>Asp. oryzae</i> , <i>Sacch. cerevisiae</i> , <i>Lb. sakei</i> , <i>Leuc. mesenteroides</i>	Japan	Kotaka et al. (2008)
<i>Sato</i>	Rice, <i>loogpang</i>	Distilled liquor, alcoholic drink	Yeasts, molds	Thailand	Phithakpol et al. (1995)
<i>Seketch</i>	Maize	Alcoholic beverage	<i>Sacch. cerevisiae</i> , <i>Sacch. chevalieri</i> , <i>Sacch. elegans</i> , <i>Lb. plantarum</i> , <i>Lc. lactis</i> , <i>B. subtilis</i> , <i>Asp. niger</i> , <i>Asp. flavus</i> , <i>Mu. Rouxii</i>	Nigeria	Blandino et al. (2003)
<i>Shochu/soju</i>	Rice, sweet potato, barley, millet, corn, <i>koji</i>	Distilled spirit, alcoholic drink	<i>Asp. awamorii</i> , <i>Asp. kawachii</i> , <i>Sacch. cerevisiae</i>	Japan	Steinkraus (1996)

(Continued)

**Table 1.12 (Continued) Some Common and Uncommon Ethnic Fermented Beverages and Alcohol Drinks of the World**

<i>Product</i>	<i>Substrate</i>	<i>Sensory Property and Nature</i>	<i>Microorganisms</i>	<i>Country</i>	<i>References</i>
<i>Soju</i>	Rice, <i>nuruk</i>	Distilled liquor	Mold, yeasts	Korea	Steinkraus (1996)
<i>Sura</i>	Finger millet, <i>dheli</i>	Alcoholic, staple	Unknown	India	Thakur et al. (2004)
<i>Takju</i>	Rice or barley, wheat flour, sweet potato, starter ( <i>nuruk</i> )	Lower or diluted concentration of <i>yakju</i> is known as <i>takju</i> , alcoholic beverage	<i>Sacch. cerevisiae</i> , <i>Hansenula anomala</i> , <i>Bacillus</i> sp., <i>Lactobacillus</i> sp.	Korea	Lee and Rhee (1970)
<i>Tapai</i>	Rice, cassava, <i>juj-paing</i>	Alcoholic beverage	Molds, yeasts	Malayasia	Wong and Jakson (1977)
<i>Tapuy</i>	Rice, bubod	Alcoholic beverage	<i>Saccharomycopsis fibuligera</i> , <i>Rhodotorula glutinis</i> , <i>Debaromyces hansenii</i> , <i>Candida parapsilosis</i> , <i>Trichosporon fennicum</i> , <i>Leuconostoc</i>	Philippines	Tanimura et al. (1977)
<i>Tchoukoutou</i>	Red sorghum	Effervescent, sweet, beer	<i>Sacch. cerevisiae</i> , <i>Hanseniaspora uvarum</i> , <i>H. guillermondii</i> , <i>Kl. marxianus</i> , <i>Clavispora lusitaniae</i>	Benin	Greppi et al. (2013a,b)
<i>Tej</i>	Honey	Sweet, effervescent, cloudy alcoholic	<i>Sacch. cerevisiae</i> , <i>Kluyvermyces bulgaricus</i> , <i>K. veronae</i> , <i>Debaromyces phaffii</i> , <i>Lactobacillus</i> , <i>Streptococcus</i> , <i>Leuconostoc</i> , <i>Pediococcus</i> spp.	Mexico	Bahiru et al. (2006)

<i>Tequila</i>	Agave juice	Effervescent, sweet, distilled alcoholic drink	LAB, <i>Sacch. cerevisiae</i> , <i>Candida lusitanae</i> , <i>Kluyvermyces marxianus</i> , <i>Pichia fermentans</i>	Mexico	De León-Rodríguez et al. (2006)
<i>Themsing</i>	Finger millet, barley	Mild-alcoholic, sweet, alcoholic beverage	Molds, yeasts	India	Tamang (2010a)
<i>Tien-chiu-niang</i>	Rice, <i>chiu-yueh</i>	Mild-alcoholic, sweet, alcoholic beverage	Molds, yeasts	China, Taiwan	Tamang (2010c)
<i>Tari or Toddy</i>	Palmyra and date palm sap	Sweet, milky, effervescent and mild alcoholic, alcoholic beverage	<i>Sacch. cerevisiae</i> , <i>Schizosaccharomyces pombe</i> , <i>Acetobacter acetii</i> , <i>A. rancens</i> , <i>A. suboxydans</i> , <i>Leuc. dextranicum</i> , <i>Micrococcus</i> sp., <i>Pediococcus</i> sp., <i>Bacillus</i> sp., <i>Sarcina</i> sp.	India	Shamala and Sreekantiah (1988)
<i>Togwa</i>	Maize	Cooked slurry, alcoholic beverage	Yeasts, LAB	East Africa	Tamang (2010c)
<i>Yakju</i>	Rice, wheat, barley, maize, <i>nuruk</i>	Alcoholic beverage	<i>Sacch. cerevisiae</i> , <i>Hansenula</i> spp.	Korea	Kim and Kim (1993)
<i>Yu</i>	Rice, <i>hamei</i>	Distilled from <i>atingba</i> , alcoholic drink	Yeasts, LAB	India	Singh and Singh (2006)
<i>Zu</i>	Rice	Distilled from fermented rice; clear liquor	Yeasts, LAB	India	Tamang (2010a)
<i>Zutho/Zhuchu</i>	Rice, <i>khekhrii</i>	Milky white, sweet-sour, mild-alcoholic	<i>Sacch. cerevisiae</i>	India	Tamang et al. (2012)

starter in preparation of *koji* for production of *saké*, *shoyu*, *miso*, *shochu* (Lee et al. 2007, Suganuma et al. 2007). The predominant fungi in Chinese *koji* are *Aspergillus* and *Candida* species along with a few species of bacteria *Weissella* and *Staphylococcus* (Yan et al. 2013).

### 1.6.2 Alcoholic Beverages and Drinks

There are 10 major categories of alcoholic beverages consumed/drunk across the world (Tamang 2010c, Table 1.12):

1. Nondistilled and unfiltered alcoholic beverages produced by amyolytic starters
2. Nondistilled and filtered alcoholic beverages produced by amyolytic starters
3. Distilled alcoholic beverages produced by amyolytic starters
4. Alcoholic beverages produced by human saliva
5. Alcoholic beverages produced by mono-fermentation
6. Alcoholic beverages produced from honey
7. Alcoholic beverages produced from plants
8. Alcoholic beverages produced by malting (germination)
9. Alcoholic beverages prepared from fruits without distillation
10. Distilled alcoholic beverages prepared from fruits and cereals

### 1.6.3 Nondistilled and Unfiltered Alcoholic Beverages Produced by Amyolytic Starters

Common examples of nondistilled and unfiltered alcoholic beverages produced by mixed amyolytic starters are *lao-chao* of China, *tapé* of Indonesia, *makgeolli* of Korea, *bhaati jaanr* and *kodo ko jaanr* of India and Nepal (Tamang 2010c), *kanji* of India (Tamang 2012b). The biological process of liquefaction and saccharification of cereal starch by filamentous molds and yeasts, supplemented by amyolytic starters, under solid-state fermentation is one of the two major stages of production of alcoholic beverages in Asia. These alcoholic beverages are mostly considered as food beverages and eaten as a staple food with high calories in many parts of Asia, for example, *kodo ko jaanr*, a fermented finger millet beverage of the Himalayan regions in India, Nepal, Bhutan, and China (Tibet) with 5% alcohol content (Thapa and Tamang 2004). *Marcha* used as an amyolytic starter supplements functional microorganisms in *kodo ko jaanr* fermentation. Yeasts *Candida glabrata*, *Pic. anomala*, *Sacch. cerevisiae*, *Sm. fibuligera*, and LAB *Lb. bifermentans* and *Ped. pentosaceus* have been recovered in *kodo ko jaanr* samples (Thapa and Tamang 2004). Saccharifying activities are mostly shown by *Rhizopus* spp. and *Sm. fibuligera* whereas liquefying activities are shown by *Sm. fibuligera* and *Sac. cerevisiae* (Thapa and Tamang 2006). *Bhaati jaanr* is the Himalayan sweet-sour, mild alcoholic food beverage paste prepared from rice and consumed as a staple food (Tamang and Thapa 2006). *Rhizopus*, *Amylomyces*, *Torulopsis*, and *Hansenula* are present in *lao-chao*, a popular ethnic fermented rice beverage of China (Wei and Jong 1983). During fermentation of *makgeolli* (a traditional Korean alcoholic beverage prepared by the amyolytic starter *nuruk*) the proportion of family Saccharomycetaceae increased significantly, and the major bacterial phylum of the samples shifted from  $\gamma$ -*Proteobacteria* to *Firmicutes* (Jung et al. 2012).

*Kanji* is an ethnic Indian strong-flavored but mild alcoholic beverage prepared from beet and carrots by natural fermentation (Batra and Millner 1974). It is drunk as a mild-alcoholic refreshing drink in India. Alcohol content in *kanji* is 2.5% and pH is 4.0 showing the product as mild-alcoholic and acidic in taste (Sura et al. 2001). During its preparation, carrots or beet are washed,



shredded and mixed with salt and mustard seeds and placed in earthen pot and allowed to ferment naturally at 26–34°C for 4–7 days. Sometimes, the mixture is inoculated with a portion of a previous batch of *kanji*. After fermentation, a pink alcoholic liquor is drained off and bottled or drunk directly. In north India it is prepared with purple or occasionally orange cultivars of carrots plus beet and spices, whereas in south India *torami*, yeast-containing fermented rice gruel is used as a starter for *kanji* production. *Hansenlu anomala*, *Candida guilliermondii*, *C. tropicalis*, and *Geotrichium candidum* are involved in *kanji* fermentation (Batra and Millner 1974). *Leuc. mesenteroides*, *Pediococcus* spp. and *Lb. dextranicum* were isolated from *kanji* fermentation (Sura et al. 2001). Kingston et al. (2010) reported *Lb. paraplantarum* and *Lb. pentosus* from *kanji* based on rep-PCR identification method.

#### 1.6.4 Nondistilled and Filtered Alcoholic Beverages Produced by Amyolytic Starters

Nondistilled and filtered alcoholic beverages produced by amyolytic starters are *saké* of Japan, *krachae* or *nam-khaao* or *sato* of Thailand, *basi* of the Philippines, *yakju* and *takju* of Korea (Table 1.12). Alcoholic beverages produced by an amyolytic starter (*koji*) is not distilled but the extract of the fermented cereals is filtered into clarified high alcohol-content liquor, for example, *saké* which is the national drink of Japan and is one of the most popular traditional nondistilled alcoholic drinks in the world. It is prepared from rice using *koji* and is clear, pale yellow, containing 15%–20% alcohol (Tamang 2010c). Improved strains of *Asp. oryzae* are used for *saké* production on an industrial scale (Kotaka et al. 2008, Hirasawa et al. 2009).

#### 1.6.5 Distilled Alcoholic Beverages Produced by Amyolytic Starters

This category of alcoholic drinks are the clear, distillate part of high alcohol-content drinks prepared from fermented cereal beverages by using amyolytic starters, for example, *raksi* of India, Nepal, and Bhutan, *shochu* of Japan, *soju* of Korea. *Raksi* is an ethnic alcoholic (22%–27% v/v) drink of the Himalayas with a characteristic aroma, and distilled from traditional fermented cereal beverages (Kozaki et al. 2000).

#### 1.6.6 Alcoholic Beverages Produced by Human Saliva

Traditionally, saliva serves as the source of amylase for conversion of cereal starch to fermentation sugars. *Chicha* is a unique ethnic fermented alcoholic (2%–12% v/v) beverage of the Andes Indians of South America, mostly Peru, and is prepared from maize by the human salivation process (Escobar et al. 1996, Hayashida 2008). *Sacch. cerevisiae*, *Sacch. apiculata*, *Sacch. pastorianus*, and species of *Lactobacillus* and *Acetobacter* are present in *chicha* (Escobar 1977). *Sacch. cerevisiae* was isolated from *chicha* and identified using MALDI-TOF (Vallejo et al. 2013).

#### 1.6.7 Alcoholic Beverages Produced by Mono Fermentation

Beer, a fermented extract of malted barley with alcohol content of 2%–8%, is the most common example of mono fermentation (*Sacch. cerevisiae* strain). Strains within *Sacch. carlsbergensis* and *Sacch. uvarum* have been merged into either *Sacch. cerevisiae* or *Sacch. pastorianus* (Kurtzman and Robnet 2003). Species of *Dekkera* produce high levels of acetic acid and esters, *Pichia* and *Hansenula* species give excessive ester production in beer (Dufour et al. 2003).

### 1.6.8 Alcoholic Beverage Produced from Honey

Some alcoholic beverages are produced from honey, for example, *tej* of Ethiopia. It is a yellow, sweet, effervescent and cloudy alcoholic (7%–14% v/v) beverage (Steinkraus 1996). *Sacch. cerevisiae*, *Kluyvermyces bulgaricus*, *Debaromyces phaffi*, and *K. veronae*, and LAB *Lactobacillus*, *Streptococcus*, *Leuconostoc*, and *Pediococcus* species were isolated from *tej* fermentation (Bahiru et al. 2006).

### 1.6.9 Alcoholic Beverages Produced from Plants

*Pulque* is one of the oldest alcoholic beverages prepared from the juice of the cactus (*Agave*) plant of Mexico (Steinkraus 1996). Bacteria present during the fermentation of *pulque* were LAB—*Lc. lactis* subsp. *lactis*, *Lb. acetotolerans*, *Lb. acidophilus*, *Lb. hilgardii*, *Lb. kefir*, *Lb. plantarum*, *Leuc. citreum*, *Leuc. kimchi*, *Leuc. mesenteroides*, *Leuc. pseudomesenteroides*, the  $\gamma$ -Proteobacteria—*Erwinia rhapsontici*, *Enterobacter* spp. and *Acinetobacter radioresistens*, several  $\alpha$ -Proteobacteria—*Zymomonas mobilis*, *Acetobacter malorum*, *Acetobacter pomorium*, *Microbacterium arborescens*, *Flavobacterium johnsoniae*, *Gluconobacter oxydans*, and *Hafnia alvei* (Escalante et al. 2004, 2008). Yeasts isolated from *pulque* include *Saccharomyces* (*Sacch. bayanus*, *Sacch. cerevisiae*, *Sacch. paradoxus*) and non-*Saccharomyces* (*Candida* spp., *C. parapsilosis*, *Clavispora lusitaniae*, *Hanseniaspora uvarum*, *Kl. lactis*, *Kl. marxianus*, *Pichia membranifaciens*, *Pichia* spp., *Torulasporea delbrueckii*) (Lappe-Oliveras et al. 2008).

*Toddy* or *tari* is an ethnic alcoholic drink of India prepared from palm juice. There are three types of *toddy* (Batra and Millner 1974): (1) *sendi*, from the palm; (2) *tari*, from the *palmyra* and date palms; and (3) *nareli*, from the coconut palm. *Geotrichum*, *Saccharomyces* and *Schizosaccharomyces* spp. of yeast are responsible for fermentation (Batra and Millner 1974, 1976). Microorganisms that are responsible in fermenting *toddy* are *Sacch. cerevisiae*, *Schizosaccharomyces pombe*, *Acetobacter acetii*, *A. rancens*, *A. suboxydans*, *Leuc. dextranicum*, *Micrococcus* sp., *Pediococcus* sp., *Bacillus* sp., and *Sarcina* sp. (Shamala and Sreekantiah 1988).

### 1.6.10 Alcoholic Beverages Produced by Malting or Germination

Malting or germination process allows amylase to break down cereal starch to sugars, which are used as substrates for alcohol fermentation, for example, *Bantu* beer or sorghum beer of the Bantu tribes of South Africa (Taylor 2003). The major part of the sorghum crop (*Sorghum caffrorum* or *S. vulgare*) is malted and used for brewing beer (Kutyauripo et al. 2009).

### 1.6.11 Alcoholic Beverages Produced from Fruits without Distillation

Wine generally refers to the alcoholic fermentation of grape juice, or other fruits, without distillation. Until 75–100 years ago, most wines were produced by spontaneous or natural alcoholic fermentation of grape juice by indigenous yeast flora (Walker 2014). Wine fermentation is initiated by the growth of various species of *Saccharomyces* and non-*Saccharomyces* yeasts (e.g., *Candida colliculosa*, *C. stellata*, *Hanseniaspora uvarum*, *Kloeckera apiculata*, *Kl. thermotolerans*, *Torulasporea delbrueckii*, *Metschnikowia pulcherrima*) (Versavaud et al. 1995, Pretorius 2000, Moreira et al. 2005, Sun et al. 2014). Using mCOLD-PCR-DGGE method, *Candida* sp. and *Cladosporium* sp. were isolated from fermenting white wine, which were not detected by conventional PCR (Takahashi et al. 2014). *Sacch. cerevisiae* strains develop during wine fermentation which play an active role in the characteristics of wine (Vilanova and Siero 2006, Capece et al. 2013).

### 1.6.12 Distilled Alcoholic Beverages without Amyolytic Starters

Distillate high alcohol-content spirit from fermented cereal is whisky, fermented molasses is rum and fermented grape is brandy. Rum production from molasses fermentation may involve contributions from *Schizosaccharomyces pombe* and *Sacch. cerevisiae* (Fahrasame and Ganow-Parfeit 1998).

### 1.6.13 Recommendations

- Exchange of LAB strains/cultures from plant-origins with superior functional properties to ferment new animal products (milks, meat, sausages, etc.).
- Every country to establish microbial sequence banks and culture collection centres, of strains/cultures isolated from naturally fermented foods and beverages.
- Basic training on molecular microbial taxonomy and identification.
- Incorporation of published research materials on ethnic fermented foods and beverages into academic programs at master and doctoral level in Food Microbiology/Biotechnology/Food Science and Technology courses in universities following the references of the academic program on fermented foods in Sikkim University (India) and Wageningen University (the Netherlands).

## 1.7 Conclusion

The sustainable use of microorganisms in food fermentation is based on the interrelationship of indigenous knowledge of food fermentation, modern expertise and information, basic understanding of the microbial background of fermentation and of Good Hygienic Practices (GHP), some experience in handling of microbial strains or cultures, even under crude conditions such as back-slopping, and conservation of microbial strains. The diversity of functional microorganisms in fermented foods and beverages consists of bacteria, yeasts, and fungi. Microorganisms establish on relevant substrates for survival and produce bioactive compounds that enrich the human diet, thereby benefiting the health of consumers. Ethnic fermented foods of the world are considered to be a means to preserve microbial diversity *ex situ* and they are custodians of microbial diversity and play a key role in the storage and supply of authentic reference material for research and development. One of the challenges facing scientists in the future will undoubtedly be to allow the large-scale production of fermented foods without losing the unique flavor and other traits associated with the traditional products from which they are derived.

## References

- Aagaard, K., Ma, J., Antony, K.M., Ganu, R., and J. Versalovic. 2014. The placenta harbors a unique microbiome. *Science Translational Medicine* 6: 237–265.
- Abe, M., Takaoka, N., Idemoto, Y., Takagi, C., Imai, T., and K. Nakasaki. 2008. Characteristic fungi observed in the fermentation process for Puer tea. *International Journal of Food Microbiology* 124: 199–203.
- Abimbola, U. 2007. Degradation of dietary fibre from gari by faecal bacteria and bacteria extracellular polysaccharides. *Nigerian Food Journal* 25: 46–58.
- Abriouel, H., Omar, N.B., López, R.L., Martínez-Cañamero, M., Keleke, S., and A. Gálvez. 2006. Culture-independent analysis of the microbial composition of the African traditional fermented foods *poto poto* and *dégué* by using three different DNA extraction methods. *International Journal of Food Microbiology* 111: 228–233.

- Abriouel, H., Benomar, N., Lucas, R., and A. Gálvez. 2011. Culture-independent study of the diversity of microbial populations in brines during fermentation of naturally-fermented Aloreña green table olives. *International Journal of Food Microbiology* 144: 487–496.
- Adams, M.R. 2010. Fermented meat products, In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 309–322.
- Adams, M.R. 2014. Vinegar, In: *Encyclopaedia of Food Microbiology*, 2nd edition (Eds: Batt, C. and Tortorello, MA). Elsevier Ltd., Oxford, pp. 717–721.
- Adhikari R.R. and H. Ghimirey. 2000. *Nepalese Society and Culture*. Vidharthi Pushtak Bhandar, Kathmandu (Nepali).
- Aidoo, K.E. and M.J.R. Nout. 2010. Functional yeasts and molds in fermented foods and beverages, In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 127–148.
- Aidoo, K.E., Nout, M.J.R., and P.K. Sarkar. 2006. Occurrence and function of yeasts in Asian indigenous fermented foods. *FEMS Yeast Research* 6: 30–39.
- Akabanda, F., Owusu-Kwarteng, J., Tano-Debrah, K., Glover, R.L.K., Nielsen, D.S., and L. Jespersen. 2013. Taxonomic and molecular characterization of lactic acid bacteria and yeasts in *numu*, a Ghanaian fermented milk product. *Food Microbiology* 34(2): 277–283.
- Aksu, M.I., Kaya, M., and H.W. Ockerman. 2005. Effect of modified atmosphere packaging and temperature on the shelf life of sliced Pastirma produced from frozen/thawed meat. *Journal of Muscle Foods* 16(3): 192–206.
- Albano, H., van Reenen, C.A., Todorov, S.D., Cruz, D., Fraga, L., Hogg, T., Dicks, L.M., and P. Teixeira. 2009. Phenotypic and genetic heterogeneity of lactic acid bacteria isolated from “Alheira”, a traditional fermented sausage produced in Portugal. *Meat Science* 82: 389–398.
- Alegría, A., González, R., Díaz, M., and B. Mayo. 2011. Assessment of microbial populations dynamics in a blue cheese by culturing and denaturing gradient gel electrophoresis. *Current Microbiology* 62: 888–893.
- Alexandraki, V., Tsakalidou, E., Papadimitriou, K., and W.H. Holzapfel. 2013. Status and trends of the conservation and sustainable use of microorganisms in food processes. Commission on Genetic Resources for Food and Agriculture. FAO Background Study Paper No. 65.
- Al-Jedah, J.H., Ali, M.Z., and R.K. Robinson. 1999. Chemical and microbiological properties of *mehi-awah*—a popular fish sauce of golf. *Journal of Food Science and Technology* 36(6): 561–564.
- Amoa-Awua, W.K., Terlabie, N.N., and E. Sakyi-Dawson. 2006. Screening of 42 *Bacillus* isolates for ability to ferment soybeans into *dawadawa*. *International Journal of Food Microbiology* 106: 343–347.
- Aneja, R.P., Mathur, B.N., Chandan, R.C., and A.K. Banerjee. 2002. Cultured/fermented products. Technology of Indian milk products: *Handbook on Process Technology Modernization for Professionals, Entrepreneurs and Scientists*. Dairy India Yearbook.
- Angelakis, E., Million, M., Henry, M., and D. Raoult. 2011. Rapid and accurate bacterial identification in probiotics and yoghurts by MALDI-TOF mass spectrometry. *Journal of Food Science* 76: 568–572.
- Ardhana, M.M. and G.H. Fleet. 1989. The microbial ecology of tapé ketan fermentation. *International Journal of Food Microbiology* 9: 157–165.
- Arroyo, P.T., Ludovico-Pelayo, L.A., Solidium, H.T., Chiu, Y.N., Lero, M.C., and E.E. Alcantara. 1978. Studies on rice-shrimp fermentation, balao-balao. *Philippines Journal of Food Science and Technology* 2(1&2): 106–125.
- Asahara, N., Zhang, X.B., and Y. Ohta. 2006. Antimutagenicity and mutagen-binding activation of mutagenic pyrolyzates by microorganisms isolated from Japanese *miso*. *Journal of Science of Food and Agriculture* 58: 395–401.
- Astuli, M. 1999. History of the development of *tempeh*. In: *The Complete Handbook of Tempeh*, ed. J. Agranoff, 2–7. Singapore: American Soybean Association.
- Azokpota, P., Hounhouigan, D.J., and M.C. Nago. 2006. Microbiological and chemical changes during the fermentation of African locust bean (*Parkia biglobosa*) to produce afitin, iru, and sonru, three traditional condiments produced in Benin. *International Journal of Food Microbiology* 107: 304–309.
- Bahiru, B., Mehari, T., and M. Ashenafi. 2006. Yeast and lactic acid flora of *tej*, an indigenous Ethiopian honey wine: Variations within and between production units. *Food Microbiology* 23: 277–282.

- Baens-Arega, L. 1977. Patis, a traditional fermented fish sauce and condiment of the Philippines. In: *The Proceeding of the Symposium on Indigenous Fermented Foods*, November 21–27, 1977, GIAM-V, Bangkok.
- Batra, L.R. 1986. Microbiology of some fermented cereals and grains legumes of India and vicinity. In: *Indigenous Fermented Food of Non-Western Origin*, eds. C.W. Hesseltine and H.L. Wang, 85–104. Berlin: J. Cramer.
- Batra, L.R. and P.D. Millner. 1974. Some Asian fermented foods and beverages and associated fungi. *Mycologia* 66: 942–950.
- Batra, L.R. and P.D. Millner. 1976. Asian fermented foods and beverages. *Development in Industrial Microbiology* 17: 117–128.
- Behre, K.-E. 1984. Zur Geschichte der Bierwürzen nach Fruchtfinden und Schriftlichen Quellen. In: *Plants and Ancient Man. Studies in Palaeoethnobotany*, eds. W. van Zeist and W.A. Casparie, 115–122. Rotterdam: A.A. Balkema.
- Berger, B., Pridmore RD, Barretto C, Delmas-Julien F, Schreiber K, Arigoni, F., and H. Brüßow. 2007. Similarity and differences in the *Lactobacillus acidophilus* group identified by polyphasic analysis and comparative genomics. *Journal of Bacteriology* 189: 1311–1321.
- Bernardeau, M., Guguen, M., and J.P. Vernoux. 2006. Beneficial lactobacilli in food and feed: Long-term use, biodiversity and proposals for specific and realistic safety assessments. *FEMS Microbiology Review* 30: 487–513.
- Bernardo, E.B., Neilan, B.A., and I. Couperwhite. 1998. Characterization, differentiation and identification of wild-type cellulose-synthesizing *Acetobacter* strains involved in nata de coco production. *Systematic and Applied Microbiology* 21(4): 599–608.
- Blandino, A., Al-Aseeri, M.E., Pandiella, S.S., Cantero, D., and C. Webb. 2003. Cereal-based fermented foods and beverages. *Food Research International* 36: 527–543.
- Briggiler-Marcó M, Capr, M.L., Quiberoni, A., Vinderola, G., Reinheimer, J.A., and E. Hynes. 2007. Nonstarter *Lactobacillus* strains as adjunct cultures for cheese making: *in vitro* characterization and performance in two model cheese. *Journal of Dairy Science* 90: 4532–4542.
- Bo, T.A. 1984a. Origin of douchi and its production technology (Part 1). *Brewing Society of Japan* 79: 221–223.
- Bo, T.A. 1984b. Origin of douchi and its production technology (Part 2). *Brewing Society of Japan* 79: 395–402.
- Bose, D.K. 1922. *Wine in Ancient India*. Kolkata: Connor.
- Bourdichon, F., Casaregola, S., Farrokh, C., Frisvad, J.C., Gerds, M.L., Hammes, W.P., Harnett, J. et al. 2012. Food fermentations: Microorganisms with technological beneficial use. *International Journal of Food Microbiology* 154(3): 87–97.
- Campbell-Platt, G. 1987. *Fermented Foods of the World: A Dictionary and Guide*. London: Butterworths.
- Carr, F.J., Chill, D., and N. Maida. 2002. The lactic acid bacteria: A literature survey. *Critical Review of Microbiology* 28: 281–370.
- Capece, A., Siesto, G., Poeta, C., Pietrafesa, R., and P. Romano. 2013. Indigenous yeast population from Georgian aged wines produced by traditional “Kakhetian” method. *Food Microbiology* 36: 447–455.
- Chang, C.H., Chen, Y.S., and F. Yanagida. 2011. Isolation and characterisation of lactic acid bacteria from *yan-jiang* (fermented ginger), a traditional fermented food in Taiwan. *Journal of Science of Food and Agriculture* 91: 1746–1750.
- Chang, H.W., Kim, K.H., Nam, Y.D., Roh, S.W., Kim, M.S., Jeon, C.O., Oh, H.M., and J.W. Bae. 2008. Analysis of yeast and archaeal population dynamics in kimchi using denaturing gradient gel electrophoresis. *International Journal of Food Microbiology* 126:159–166.
- Chakrabarty, J., Sharma, G.D., and J.P. Tamang. 2014. Traditional technology and product characterization of some lesser-known ethnic fermented foods and beverages of North Cachar Hills District of Assam. *Indian Journal of Traditional Knowledge* 13(4): 706–715.
- Chao, S-H., Wu, R-J., Watanabe, K., and Y-C. Tsai. 2009. Diversity of lactic acid bacteria in *suan-tsai* and *fu-tsai*, traditional fermented mustard products of Taiwan. *International Journal of Food Microbiology* 135: 203–210.



- Chao, S-H., Tomii, Y., Watanabe, K., and Y-C. Tsai. 2008. Diversity of lactic acid bacteria in fermented brines used to make stinky tofu. *International Journal of Food Microbiology* 123: 134–141.
- Chao, S.H., Kudo, Y., Tsai, Y.C., and K. Watanabe. 2012. *Lactobacillus futsaii* sp. nov., isolated from traditional fermented mustard products of Taiwan, *fu-tsai* and *suan-tsai*. *International Journal of Systematic and Evolutionary Microbiology* 62: 489–494.
- Chen, Y.S., Yanagida, F., and J.S. Hsu. 2006. Isolation and characterization of lactic acid bacteria from *suan-tsai* (fermented mustard), a traditional fermented food in Taiwan. *Journal of Applied Microbiology* 101: 125–130.
- Chen, Y.S., Wu, H.C., Liu, C.H., Chen, H.C., and F. Yanagida. 2010. Isolation and characterization of lactic acid bacteria from *jiang-sun* (fermented bamboo shoots), a traditional fermented food in Taiwan. *Journal of Science of Food and Agriculture* 90: 1977–1982.
- Chang, J.H. 1975. Studies on the origin of Korean vegetable pickles. *Thesis Collection of Sung-Sim Women College* 6: 149–174.
- Cheigh, H.S. 2002. *Kimchi Culture and Dietary Life in Korea*. Seoul: Hyoil Publishing Co.
- Chettri, R. 2013. *Microbiological Evaluation of turangbai and bekaung, ethnic fermented soybean foods of North East India*. Ph.D. thesis, North Bengal University, Siliguri, India, p. 380.
- Chettri, R. and J.P. Tamang. 2008. Microbiological evaluation of maseura, an ethnic fermented legume-based condiment of Sikkim. *Journal of Hill Research* 21: 1–7.
- Choi, S.H., Lee, M.H., Lee, S.K., and Oh, M.J. 1995. Microflora and enzyme activity of conventional *meju* and isolation of useful mould. *Journal of Agricultural Science Chungnam National University, Korea* 22: 188–197.
- Choi, U.K., Kim, M.H., and N.H. Lee. 2007. The characteristics of cheonggukjang, a fermented soybean product, by the degree of germination of raw soybeans. *Food Science and Biotechnology* 16: 734–739.
- Chokesajjawatee, N., Pornaem, S., Zo, Y.G., Kamdee, S., Luxananil, P., Wanasen, S., and R. Valysevi. 2009. Incidence of *Staphylococcus aureus* and associated risk factors in Nham, a Thai fermented pork product. *Food Microbiology* 26: 547–551.
- Chunhachart, O., Itoh, T., Sukchotiratana, M., Tanimoto, H., and Y. Tahara. 2006. Characterization of  $\gamma$ -glutamyl hydrolase produced by *Bacillus* sp. isolated from Thai thua-nao. *Bioscience, Biotechnology and Biochemistry* 70: 2779–2782.
- Cocolin, L., Aggio, D., Manzano, M., Cantoni, C., and G. Comi. 2002. An application of PCR-DGGE analysis to profile the yeast populations in raw milk. *International Dairy Journal* 12: 407–411.
- Cocolin, L., Alessandria V., Dolci, P., Gorra, R., and K. Rantsiou. 2013. Culture independent methods to assess the diversity and dynamics of microbiota during food fermentation. *International Journal of Food Microbiology* 167(1): 29–43.
- Cocolin, L. and D. Ercolini, (Editors). 2008. *Molecular Techniques in the Microbial Ecology of Fermented Foods*. New York: Springer.
- Cocolin, L., Dolci, P., and K. Rantsiou. 2011. Biodiversity and dynamics of meat fermentations: The contribution of molecular methods for a better comprehension of a complex ecosystem. *Meat Science* 89(3): 296–302.
- Coppola, S., Fusco, V., Andolfi, R., Aponte, M., Aponte, M., Blalotta, G., Ercolini, D., and G. Moschetti. 2006. Evaluation of microbial diversity during the manufacture of Fior di Latte di Agerola, a traditional raw milk pasta-filata cheese of the Naples area. *Journal of Dairy Research* 73: 264–272.
- Corsetti, A. and L. Settanni. 2007. Lactobacilli in sourdough fermentation. *Food Research International* 40: 539–558.
- Coton, E., Desmonts, M.H., Leroy, S., Coton, M., Jamet, E., Christeans, S., Donnio, P.Y., Lebert, I., and R. Talon. 2010. Biodiversity of coagulase-negative staphylococci in French cheeses, dry fermented sausages, processing environments and clinical samples. *International Journal of Food Microbiology* 137: 221–229.
- Curtis, R.I. 2001. *Ancient Food Technology*. Leiden: Brill.
- Dajanta, K., Apichartsrangkoon, A., Chukeatirote, E., Richard, A., and R.A. Frazier. 2011. Free-amino acid profiles of *thua nao*, a Thai fermented soybean. *Food Chemistry* 125: 342–347.
- Dalmacio, L.M.M., Angeles, A.K.J., Larcia, L.L.H., Balolong, M., and R. Estacio. 2011. Assessment of bacterial diversity in selected Philippine fermented food products through PCR-DGGE. *Beneficial Microbes* 2: 273–281.

- Davis, J.G. 1964. *Cheese, vol 1, Basic Technology*. Edinburg: Churchill Livingstone.
- de Bruyne, K., Camu, N., Lefebvre, K., De Vuyst, L., and P. Vandamme. 2008. *Weissella ghanensis* sp. nov., isolated from a Ghanaian cocoa fermentation. *International Journal of Systematic and Evolutionary Microbiology* 58: 2721–2725.
- de Bruyne, K., Camu, N., De Vuyst, L., and P. Vandamme. 2009. *Lactobacillus fabifermentans* sp. nov. and *Lactobacillus cacaonum* sp. nov., isolated from Ghanaian cocoa fermentations. *International Journal of Systematic and Evolutionary Microbiology* 59: 7–12.
- de Bruyne, K., Camu, N., de Vuyst, L., and P. Vandamme. 2010. *Weissella fabaria* sp. nov., from a Ghanaian cocoa fermentation. *International Journal of Systematic and Evolutionary Microbiology* 60: 1999–2005.
- de Bruyne, K., Franz, C.M., Vancanneyt, M., Schillinger, U., Mozzi, F., de Valdez, G.F., de Vuyst, L., and P. Vandamme. 2008. *Pediococcus argentinicus* sp. nov. from Argentinean fermented wheat flour and identification of *Pediococcus* species by pheS, rpoA and atpA sequence analysis. *International Journal of Systematic and Evolutionary Microbiology* 58: 2909–2916.
- de Bruyne, K., Schillinger, U., Caroline, L., Boehringer, B., Cleenwerck, I., Vancanneyt, M., De Vuyst, L., Franz C.M., and P. Vandamme. 2007. *Leuconostoc holzappelii* sp. nov., isolated from Ethiopian coffee fermentation and assessment of sequence analysis of housekeeping genes for delineation of *Leuconostoc* species. *International Journal of Systematic and Evolutionary Microbiology* 57: 2952–2959.
- De León-Rodríguez, A., González-Hernández, L., Barba de la Rosa, A., Escalante-Minakata, P., and López, M. 2006. Characterization of volatile compounds of mezcal, an ethnic alcoholic beverage obtained from *Agave salmiana*. *Journal of Agriculture and Food Chemistry* 54: 1337–1341.
- Delong, D. 1992. *How to Dry Foods*. Penguin Group, New York, p. 79.
- Denter, J. and B. Bisping. 1994. Formation of B-vitamins by bacteria during the soaking process of soybeans for tempe fermentation. *International Journal of Food Microbiology* 22(1): 23–31.
- de Ramesh, C.C., White, C.H., Kilara, A., and Y.H. Hui. 2006. *Manufacturing Yogurt and Fermented Milks*. Blackwell Publishing, Oxford.
- de Vuyst, L., Vrancken G, Ravyts F, Rimaux T., and S. Weckx. 2009. Biodiversity, ecological determinants, and metabolic exploitation of sourdough microbiota. *Food Microbiology* 26: 666–675.
- Dewan, S. and J.P. Tamang. 2006. Microbial and analytical characterization of *Chhu*, a traditional fermented milk product of the Sikkim Himalayas. *Journal of Scientific and Industrial Research* 65: 747–752.
- Dewan, S. and J.P. Tamang. 2007. Dominant lactic acid bacteria and their technological properties isolated from the Himalayan ethnic fermented milk products. *Leeuwenhoek International Journal of General and Molecular Microbiology* 92: 343–352.
- Diancourt, L., Passet, V., Chervaux, C., Garault, P., Smokvina, T., and S. Brisse. 2007. Multilocus sequence typing of *Lactobacillus casei* reveals a clonal population structure with low levels of homologous recombination. *Applied Environmental Microbiology* 73: 6601–6611.
- Dirar, H.A., Harper, D.B., and M.A. Collins. 2006. Biochemical and microbiological studies on kawal, a meat substitute derived by fermentation of *Cassia obtusifolia* leaves. *Journal of Science of Food and Agriculture* 36: 881–892.
- Dirar, H.A. 1993. *The Indigenous Fermented Foods of Sudan*. Wallingford: CAB International.
- Dufour, J., Verstrepen, K., and G. Derdelinckx. 2003. Brewing yeasts, In: *Yeast in Foods*, Boekhout T, Rober V. eds. Woodhead, Cambridge, 347–388.
- Dung, N.T.P., 2004. Defined fungal starter granules for purple glutinous rice wine, Ph.D. Thesis. Wageningen University, Wageningen, The Netherlands, p. 110.
- Dung, N.T.P., Rombouts, F.M., and M.J.R. Nout. 2006. Functionality of selected strains of moulds and yeasts from Vietnamese rice wine starters. *Food Microbiology* 23: 331–340.
- Dung, N.T.P., Rombouts, F.M., and Nout, M.J.R. 2007. Characteristics of some traditional Vietnamese starch-based rice wine starters (*Men*). *LWT—Food Science and Technology* 40: 130–135.
- Dušková, M., Šedo, O., Kšicová, K., Zdráhal, Z., and R. Karpíšková. 2012. Identification of lactobacilli isolated from food by genotypic methods and MALDI-TOF MS. *International Journal of Food Microbiology* 159: 107–114.
- Encinas, J.P., Lopez-Diaz, T.M., Garcia-Lopez, M.L., Otero, A, and Moreno, B. 2000. Yeast populations on Spanish fermented sausages. *Meat Science* 54: 203–208.

- Endo, A., Mizuno, H., and S. Okada. 2008. Monitoring the bacterial community during fermentation of sunki, an unsalted, fermented vegetable traditional to the Kiso area of Japan. *Letters in Applied Microbiology* 47: 221–226.
- Escalante, A., Rodríguez, M.E., Martínez, A., López-Munguía, A., Bolívar, F., and G. Gosset. 2004. Characterization of bacterial diversity in *Pulque*, a traditional Mexican alcoholic fermented beverage, as determined by 16S rDNA analysis. *FEMS Microbiology Letters* 2: 273–279.
- Escalante, A.M., Giles-Gómez, G., Hernández, M.S., Córdova-Aguilar, A., López-Munguía, F., Gosset, G., and F. Bolívar. 2008. Analysis of bacterial community during the fermentation of pulque, a traditional Mexican alcoholic beverage, using a polyphasic approach. *International Journal of Food Microbiology* 124: 126–134.
- Escobar, A. 1977. The American maize beverage chichi. MSc thesis. New York: Cornell University.
- Escobar, A., Gardner, A., and K.H. Steinkraus. 1996. Studies of South American chichi, In: *Handbook of Indigenous Fermented Food*, 2nd edition, ed., Steinkraus K.H., Marcel Dekker, Inc., New York, 402–406.
- Fahrasame, L. and B. Ganow-Parfeit. 1998. Microbial flora of rum fermentation media. *Journal of Applied Microbiology* 84: 921–928.
- Farhad, M., Kailasapathy, K., and Tamang, J.P. 2010. Health aspects of fermented foods, In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 391–414.
- FAO, 1998. Fermented fruits and vegetables. A global perspective. *FAO Agricultural Services Bulletin* 134.
- FAO, 1999. Fermented cereals. A global perspective. *FAO Agricultural Services Bulletin* No. 138.
- Feng, X.M., Eriksson, A.R.B., and J. Schnürer. 2005. Growth of lactic acid bacteria and *Rhizopus oligosporus* during barley tempeh fermentation. *International Journal of Food Microbiology* 104: 249–256.
- Fernandez-Lopez, J., Sendra, E., Sayas-Barbera, E., Navarro, C., and J.A. Perez-Alvarez. 2008. Physico-chemical and microbiological profiles of “Salchichón” (Spanish dry fermented sausage) enriched with orange fiber. *Meat Science* 80: 410–417.
- Foschino, R., Gallina, S., Andrighetto, C., Rossetti, L., and A. Galli. 2004. Comparisons of cultural methods for the identification and molecular investigations of yeasts from sourdoughs for Italian sweet baked products. *FEMS Yeasts Research* 4: 609–618.
- Fujii, T., Wu, Y.C., Suzuk, T., and B. Kimura. 1999. Production of organic acids by bacteria during the fermentation of squid shiokara. *Fisheries Science* 65: 671–672.
- Gadaga, T.H., Mutukumira, A.N., and Narvhus, J.A. 2000. Enumeration and identification of yeast isolates from Zimbabwean traditional fermented milk. *International Dairy Journal* 10: 459–466.
- Ganasen, P. and S. Benjakul. 2010. Physical properties and microstructure of pidan yolk as affected by different divalent and monovalent cations. *LWT—Food Science and Technology* 43: 77–85.
- Gänzle, M.G., Ehmann, M., and W.P. Hammes. 1998. Modeling of growth of *Lactobacillus sanfranciscensis* and *Candida milleri* in response to process parameters of sourdough fermentation. *Applied Environmental Microbiology* 64: 2616–2623.
- García-Fontan, M.C., Lorenzo, J.M., Parada, A., Franco, I., and J. Carballo. 2007. Microbiological characteristics of “Androlla”, a Spanish traditional pork sausage. *Food Microbiology* 24: 52–58.
- García-Varona, M., Santos, E.M., Jaime, I., and J. Rovira. 2000. Characterization of Micrococcaceae isolated from different varieties of chorizo. *International Journal of Food Microbiology* 54: 189–195.
- Gençelep, H., Kaban, G., Aksu, M.I., Oz, F., and M. Kaya. 2008. Determination of biogenic amines in sucuk. *Food Control* 19(9): 868–872.
- Ghosh, J. and G.S. Rajorhia. 1990. Selection of starter culture for production of indigenous fermented milk product (*Misti dahi*). *Lait* 70: 147–154.
- Giraffa, G. and D. Carminati. 2008. Molecular techniques in food fermentation: Principles and applications. In: *Molecular Techniques in the Microbial Ecology of Fermented Foods*. (Eds: Coccolin, L. and Ercolini, D.). Springer Science + Business Media, LCC, New York. Chapter 1, pp. 1–30.
- Gode, P.K. 1943. Some notes on the history of Indian dietetics with special reference to the history of jalebi. *New Indian Antique* 169–181.
- Goncalves de Lima, O. 1975. *Pulque, balche e pajuaru*. Recife: Universidade Federal de Pernambuco.
- Greppi, A., Rantsiou, K., Padonou, W., Hounhouigan, J., Jespersen, L., Jakobsen, M., and L. Coccolin. 2013a. Determination of yeast diversity in ogi, mawè, gowé and tchoukoutou by using culture-dependent and -independent methods. *International Journal of Food Microbiology* 165(2): 84–88.



- Greppi, A., Rantisou, K., Padonou, W., Hounhouigan, J., Jespersen, L., Jakobsen, M., and L. Cocolin. 2013b. Yeast dynamics during spontaneous fermentation of mawè and tchoukoutou, two traditional products from Benin. *International Journal of Food Microbiology* 165(2): 200–207.
- Gorer, G. 1938. *The Lepchas of Sikkim*. Delhi: Gian Publishing House.
- Guan, L., Cho, K.H., and J.H. Lee. 2011. Analysis of the cultivable bacterial community in *jeotgal*, a Korean salted and fermented seafood, and identification of its dominant bacteria. *Food Microbiology* 28: 101–113.
- Guarner, F., Perdigon, G., Corthier G, Salminen S, Koletzko, B., and L. Morelli. 2005. Should yoghurt cultures be considered probiotic? *British Journal of Nutrition* 93: 783–786.
- Gupta, M., Khetarpaul, N., and B.M. Chauhan. 1992. Preparation, nutritional value and acceptability of barley *rabadi*—an indigenous fermented food of India. *Plant Foods for Human Nutrition* 42: 351–358.
- Gupta, R.C., Mann, B., Joshi, V.K., and D.N. Prasad. 2000. Microbiological, chemical and ultrastructural characteristics of misti doi (sweetened dahi). *Journal of Food Science and Technology* 37(1): 54–57.
- Guyot, J.P. 2010. Fermented cereal products, In: Tamang, J.P., and Kailasapathy, K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 247–261.
- Hamad, S.H., Dieng, M.M.C., Ehrmann, M.A., and R.F. Vogel. 1997. Characterisation of the bacterial flora of Sudanese sorghum flour and sorghum sourdough. *Journal of Applied Microbiology* 83: 764–770.
- Hammes, W.P., Brandt, M.J., Francis, K.L., Rosenheim, J., Seitter, M.F.H., and S.A. Vogelmann. 2005. Microbial ecology of cereal fermentations. *Trends in Food Science and Technology* 16: 4–11.
- Haard, N.F., Odunfa, S.A., Lee, C.H., Quintero-Ramírez, R., Lorence-Quiñones, A., and C. Wachter-Radarte. 1999. *Fermented Cereals: A Global Perspective*. FAO Agricultural Service Bulletin 138, 63–97. Rome: Food and Agriculture Organization.
- Han, B.Z., Beumer, R.R., Rombouts, F.M., and Nout, M.J.R. 2001. Microbiological safety and quality of commercial sufu—A Chinese fermented soybean food. *Food Control* 12: 541–547.
- Hansen, E.B. 2002. Commercial bacterial starter cultures for fermented foods of the future. *International Journal of Food Microbiology* 78: 119–131.
- Hansen, E.B. 2004. Micro-organisms. In: *Handbook of Food and Beverage Fermentation Technology*, Eds. Hui, Y.H., Meunier-Goddik, L., Hansen, Å.S., Josephsen, J., Nip, W.K., Stanfield, P.S., and Toldrá, F., Marcel Dekker, Inc., 270 Madison Avenue, New York, 2, pp. 9–21.
- Hao, Y., Zhao, L., Zhang, H., and Z. Zhai. 2010. Identification of the bacterial biodiversity in koumiss by denaturing gradient gel electrophoresis and species-specific polymerase chain reaction. *Journal of Dairy Science* 93: 1926–1933.
- Hara, T., Chetanachit, C., Fujio, Y., and S. Ueda. 1986. Distribution of plasmids in polyglutamate-producing *Bacillus* strains isolated from “natto”-like fermented soybeans, “thua nao,” in Thailand. *Journal of General and Applied Microbiology* 32: 241–249.
- Hara, T., Hiroyuki, S., Nobuhide, I., and K. Shinji. 1995. Plasmid analysis in polyglutamate-producing *Bacillus* strain isolated from nonsalty fermented soybean food, “kinema”, in Nepal. *Journal of General and Applied Microbiology* 41: 3–9.
- Hara, T., Zhang, J.R., and S. Ueda. 1983. Identification of plasmids linked with polyglutamate production in *Bacillus subtilis* (natto). *Journal of General and Applied Microbiology* 29: 345–354.
- Harun-ur-Rashid, M., Togo, K., Useda, M., and T. Miyamoto. 2007. Probiotic characteristics of lactic acid bacteria isolated from traditional fermented milk “Dahi” in Bangladesh. *Pakistan Journal of Nutrition* 6: 647–652.
- Haruta, S., Ueno, S., Egawa, I., Hashiguchi, K., Fujii, A., Nagano, M., and Y.I. Igarashi. 2006. Succession of bacterial and fungal communities during a traditional pot fermentation of rice vinegar assessed by PCR-mediated denaturing gradient gel electrophoresis. *International Journal of Food Microbiology* 109: 79–87.
- Hayashida, F.M. 2008. Ancient beer and modern brewers: Ethnoarchaeological observations of *chicha* production in two regions of the North Coast of Peru. *Journal of Anthropological Archaeology* 27(2): 161–174.
- Hesseltine, C.W. 1983. Microbiology of oriental fermented foods. *Annual Review of Microbiology* 37: 575–601.
- Hesseltine, C.W. 1991. Zygomycetes in food fermentations. *Mycologist* 5: 162–169.
- Hesseltine, C.W. and C.P. Kurtzman. 1990. Yeasts in amylolytic food starters. *Anns Inst Biol Uni Nac Antón México Ser Botany* 60: 1–7.

- Hesseltine, C.W. and M.L. Ray. 1988. Lactic acid bacteria in murcha and ragi. *Journal of Applied Bacteriology* 64: 395–401.
- Hesseltine, C.W., Rogers, R., and F.G. Winarno. 1988. Microbiological studies on amylolytic Oriental fermentation starters. *Mycopathologia* 101: 141–155.
- Hirasawa, T., Yamada, K., Nagahisa, K., Dinh, T.N., Furusawa, C., Katakura, Y., SHioya, S., and Shimizu, H. 2009. Proteomic analysis of responses to osmotic stress in laboratory and sake-brewing strains of *Saccharomyces cerevisiae*. *Process Biochemistry* 44: 647–653.
- Holzappel, W.H. and G. Müller. 2007. Kaffee. In: *Lebensmittel pflanzlicher Herkunft*, Holzappel, W.H. ed. e Behr's Verlag, Hamburg, 457–467.
- Holzappel, W.H., Schillinger, U., Toit, M.D., and L. Dicks. 1997. Systematics of probiotic lactic acid bacteria with reference to modern phenotypic and genomic methods. *Microecology and Therapy* 26: 1–10.
- Hong, S.W., Choi, J.Y., and K.S. Chung. 2012. Culture-based and denaturing gradient gel electrophoresis analysis of the bacterial community from chungkookjang, a traditional Korean fermented soybean food. *Journal of Food Science* 77(10): M572–M578.
- Ho, C.C. 1986. Identity and characteristics of *Neurospora intermedia* responsible for *oncom* fermentation in Indonesia. *Food Microbiology* 3: 115–132.
- Hooker, J.D. 1854. *Himalayan Journals: Notes of a Naturalist in Bengal, the Sikkim and Nepal Himalayas, the Khasia Mountains*. London: John Murray.
- Hounhouigan, D., Nout, M., Nago, C., Houben, J., and F. Rombouts. 1993. Characterization and frequency distribution of species of lactic acid bacteria involved in the processing of mawe, a fermented maize dough from Benin. *International Journal of Food Microbiology* 18: 279–287.
- Huang, Y., Luo, Y., Zhai, Z., Zhang, H., Yang, C., Tian, H., Li, Z., Feng, J., Liu, H., and Y. Hao. 2009. Characterization and application of an anti-*Listeria* bacteriocin produced by *Pediococcus pentosaceus* 05–10 isolated from Sichuan Pickle, a traditionally fermented vegetable product from China. *Food Control* 20: 1030–1035.
- Hulse, J.H. 2004. Biotechnologies: Past history, present state and future prospects. *Trends in Food Science* 15: 3–18.
- Humblot, C. and J.P. Guyot. 2009. Pyrosequencing of tagged 16S rRNA gene amplicons for rapid deciphering of the microbiomes of fermented foods such as pearl millet slurries. *Applied and Environmental Microbiology* 75: 4354–4361.
- Hwanhlem, N., Buradaleng, S., Wattanachant, S., Benjakul, S., Tani, A., and Maneerat, S. 2011. Isolation and screening of lactic acid bacteria from Thai traditional fermented fish (*Plasom*) and production of *Plasom* from selected strains. *Food Control* 22: 401–407.
- Iacumin, L., Cecchini, F., Manzano, M., Osualdini, M., Boscolo, D., Orlic, S., and G. Comi. 2009. Description of the microflora of sourdoughs by culture-dependent and culture-independent methods. *Food Microbiology* 26: 128–135.
- Ijong, F.G. and Y. Ohta. 1996. Physicochemical and microbiological changes associated with bakasang processing—A traditional Indonesian fermented fish sauce. *Journal of the Science of Food and Agriculture* 71: 69–74.
- Inatsu, Y., Nakamura N, Yuriko Y, Fushimi T, Watanasritum L, Kawanmoto S. 2006. Characterization of *Bacillus subtilis* strains in Thua nao, a traditional fermented soybean food in northern Thailand. *Letters in Applied Microbiology* 43: 237–242.
- Ishige, N. 1993. Cultural aspects of fermented fish products in Asia. In: *Fish Fermentation Technology*, eds. C.-H. Lee, K.H. Steinkraus and P.J. Alan Reilly, 13–32. Tokyo: United Nations University Press.
- Itoh, H., Tachi, H., and S. Kikuchi. 1993. Fish fermentation technology in Japan. In: *Fish Fermentation Technology* (Eds: Lee, C.H., Steinkraus, K.H. and Alan Reilly, P.J.), pp. 177–186. United Nations University Press, Tokyo.
- Itoh, H., Tong, J., and Y. Li. 1996. Chinese dauchi, from itohiki natto to nonmashed miso. *Miso Science Technology* 44: 224–250.
- Iyengar, S.H. 1950. *Lokopakara of Chavundarava*. Madras: Oriental Manuscripts Library.
- Jagannath, A., Raju, P.S., and A.S. Bawa. 2010. Comparative evaluation of bacterial cellulose (natta) as a cryoprotectant and carrier support during the freeze drying process of probiotic lactic acid bacteria. *LWT—Food Science and Technology* 43: 1197–1203.

- Jeng, K.C., Chen, C.S., Fang, Y.P., Hou, R.C.W., and Y.S. Chen. 2007. Effect of microbial fermentation on content of statin, GABA, and polyphenols in Puerh tea. *Journal of Agriculture and Food Chemistry* 55: 8787–8792.
- Jennessen, J., Schnürer, J., Olsson, J., Samson, R.A., and J. Dijksterhuis. 2008. Morphological characteristics of sporangiospores of the *tempeh* fungus *Rhizopus oligosporus* differentiate it from other taxa of the *R. microsporus* group. *Mycology Research* 112: 547–563.
- Jeyaram, J., Anand Singh, Th., Romi, W., Ranjita Devi, A., Mohendro Singh, W., Dayanidhi, H., Rajmuhon Singh, N., and J.P. Tamang. 2009. Traditional fermented foods of Manipur. *Indian Journal of Traditional Knowledge* 8(1): 115–121.
- Jeyaram, K., Romi, W., Singh, TAh., Devi, A.R., and S.S. Devi. 2010. Bacterial species associated with traditional starter cultures used for fermented bamboo shoot production in Manipur state of India. *International Journal of Food Microbiology* 143: 1–8.
- Jeyaram, K., Mohendro Singh, W., Premarani, T., Ranjita Devi, A., Selina Chanu, K., Talukdar, N.C., and M. Rohinikumar Singh. 2008a. Molecular identification of dominant microflora associated with “Hawaijar” – a traditional fermented soybean (*Glycine max* L.) food of Manipur, India. *International Journal of Food Microbiology* 122: 259–268.
- Jeyaram, K., Singh, W.M., Capece, A., and P. Romano. 2008b. Molecular identification of yeast species associated with “Hamei”—A traditional starter used for rice wine production in Manipur, India. *International Journal of Food Microbiology* 124: 115–125.
- Jeyaram, K., Tamang, J.P., Capece, A., and P.P. Romano. 2011. Geographical markers for *Saccharomyces cerevisiae* strains with similar technological origins domesticated for rice-based ethnic fermented beverages production in North East India. *Leeuwenhoek International Journal of General and Molecular Microbiology* 100: 569–578.
- Jianzhong, Z., Xiaolia, L., Hanhub, J., and D. Mingshengb. 2009. Analysis of the microflora in Tibetan kefir grains using denaturing gradient gel electrophoresis *Food Microbiology* 26: 770–775.
- Jiang, J. J., Zeng, Q.X., Zhu, Z.W., and L.Y. Zhang. 2007. Chemical and sensory changes associated Yu-lu fermentation process—A traditional Chinese fish sauce. *Food Chemistry* 104: 1629–1634.
- Johanningsmeier, S., McFeeters, R.F., Fleming, H.P., and R.L. Thompson. 2007. Effects of *Leuconostoc mesenteroides* starter culture on fermentation of cabbage with reduced salt concentrations. *Journal of Food Science* 72: M166–M172.
- Josephsen, J. and L. Jespersen L. 2004. Handbook of food and beverage fermentation technology. In: *Starter Cultures and Fermented Products*, Eds. Hui, Y.H., Meunier-Goddik, L., Hansen, Å.S., Josephsen, J., Nip, W.K., Stanfield, P.S., and Toldrá, F., Marcel Dekker, Inc., 270 Madison Avenue, New York, 3, pp. 23–49.
- Jung, D.H., 2012. *Great Dictionary of Fermented Food*. Yuhanmunwhasa, Inc., Kayang-dong, Seoul.
- Jung, J.Y., Lee, S.H., Kim, J.M., Park, M.S., Bae, J.W., Hahn, Y., Madsen, E.L., and C.O. Jeon. 2011. Metagenomic analysis of kimchi, a traditional Korean fermented food. *Applied and Environmental Microbiology* 77: 2264–2274.
- Jung, J.Y., Lee, S.H., Lee, H.J., and C.O. Jeon. 2013. Microbial succession and metabolite changes during fermentation of *saeu-jeot*: Traditional Korean salted seafood. *Food Microbiology* 34(2): 360–368.
- Jung, M.J., Nam, Y.D., Roh, S.W., and J.W. Bae. 2012. Unexpected convergence of fungal and bacterial communities during fermentation of traditional Korean alcoholic beverages inoculated with various natural starters. *Food Microbiology* 30: 112–123.
- Kahala M., Mäki, M., Lehtovaara, A., Tapanainen, J.M., Katiska, R., Juuruskorpi, M., Juhola, J., and V. Joutsjoki. 2008. Characterization of starter lactic acid bacteria from the Finnish fermented milk product viili. *Journal of Applied Microbiology* 105: 1929–1938.
- Karim, M.I.A. 1993. Fermented fish products in Malaysia. In: *Fish Fermentation Technology* (Eds: Lee, C.H., Steinkraus, K.H. and Alan Reilly, P.J.), pp. 95–106. United Nations University Press, Tokyo.
- Karki, T., Okada, S., Baba, T., Itoh, H. and M. Kozaki. 1983. Studies on the microflora of Nepalese pickles gundruk. *Nippon Shokuhin Kogyo Gakkaishi* 30: 357–367.
- Karagozlu, N., Ergonul, B., and C. Karagozlu. 2008. Microbiological attributes of instant tarhana during fermentation and drying. *Bulgarian Journal of Agricultural Science* 14: 535–541.

- Khanh, T.M., May, B.K., Smooker, P.M., Van, T.T.H., and P.J. Coloe. 2011. Distribution and genetic diversity of lactic acid bacteria from traditional fermented sausage. *Food Research International* 44: 338–344.
- Khatri, P.K. 1987. *Nepali Samaj ra Sanskriti (Prachin-Madhyakal)*. Shaja Prakashan, Kathmandu (Nepali).
- Kiers, J.L., Van laeken, A.E.A., Rombouts, F.M., and M.J.R. Nout. 2000. *In vitro* digestibility of *Bacillus* fermented soya bean. *International Journal of Food Microbiology* 60: 163–169.
- Kim, J.O. and J.G. Kim. 1993. Microbial and enzymatic properties related to brewing of traditional ewhaju. *Korean Journal of Society of Food Science* 9(4): 266–271.
- Kim, T.W., Lee, J.W., Kim, S.E., Park, M.H., Chang, H.C., and H.Y. Kim. 2009. Analysis of microbial communities in *doenjang*, a Korean fermented soybean paste, using nested PCR-denaturing gradient gel electrophoresis. *International Journal of Food Microbiology* 131: 265–271.
- Kim, Y.B., Seo, Y.G., and C.H. Lee. 1993. Growth of microorganisms in dorsal muscle of gulbi during processing and their effect on its quality. In: *Fish Fermentation Technology* (Eds: Lee, C.H., Steinkraus, K.H. and Alan Reilly, P.J.), pp. 281–289. United Nations University Press, Tokyo.
- Kimura, K. and Y. Itoh. 2007. Determination and characterization of IS4*Bsu1*-insertion loci and identification of a new insertion sequence element of the IS256 family in a natto starter. *Bioscience, Biotechnology and Biochemistry* 71: 2458–2464.
- Kingston, J.J., Radhika, M., Roshini, P.T., Raksha, M.A., Murali, H.S., and H.V. Batra. 2010. Molecular characterization of lactic acid bacteria recovered from natural fermentation of beet root and carrot Kanji. *Indian Journal of Microbiology* 50: 292–298.
- Kisworo, D. 2003. Characteristics of lactic acid bacteria from raw milk and white soft cheese. *Philippine Agricultural Scientist* 86: 56–64.
- Kitamoto, K. 2002. Molecular biology of the *Koji* molds. *Advances in Applied Microbiology* 51: 129–153.
- Kiuchi, K. 2001. Miso and natto. *Food Culture* 3: 7–10.
- Kiyohara, M., Koyanagi, T., Matsui, H., Yamamoto, K., Take, H., Katsuyama, Y., Tsuji, A., Miyamae, H., Kondo, T., Nakamura, S., Katayama, T., and H. Kumagai. 2012. Changes in microbiota population during fermentation of Narezushi as revealed by pyrosequencing analysis. *Bioscience, Biotechnology, Biochemistry* 76: 48–52.
- Kobayashi, T., Kimura, B., and Fujii, T. 2000a. Strictly anaerobic halophiles isolated from canned Swedish fermented herrings (Suströmming). *International Journal of Food Microbiology* 54: 81–89.
- Kobayashi, T., Kimura, B., and T. Fujii. 2000b. *Haloanaerobium fermentans* sp. nov., a strictly anaerobic, fermentative halophile isolated from fermented puffer fish ovaries. *International Journal of Systematic and Evolutionary Microbiology* 50: 1621–1627.
- Kobayashi, T., Kimura, B., and T. Fujii. 2000c. Differentiation of *Tetragenococcus* populations occurring in products and manufacturing processes of puffer fish ovaries fermented with rice-bran. *International Journal of Food Microbiology* 56: 211–218.
- Kolawole, O.M., Kayode, R.M.O., and Akinduyo, B. 2013. Proximate and microbial analyses of burukutu and pito produced in Ilorin, Nigeria. *African Journal of Microbiology* 1(1): 15–17.
- Kosikowski, F.V. and V.V. Mistry. 1997. *Cheese and Fermented Milk Foods*. Connecticut: LLC Westport.
- Kotaka, A., Bando, H., Kaya, M., Kato-Murai, M., Kuroda, K., Sahara, H., Hata, Y., Kondo, A., and Ueda, M. 2008. Direct ethanol production from barley  $\beta$ -glucan by sake yeast displaying *Aspergillus oryzae*  $\beta$ -glucosidase and endoglucanase. *Journal of Bioscience and Bioengineering* 105: 622–627.
- Kozaki, M. 1976. Fermented foods and related microorganisms in Southeast Asia. *Proceedings of Japanese Association of Mycotoxicology* 2: 1–9.
- Kozaki, M., Tamang, J.P, Kataoka, J., Yamanaka, S., and S. Yoshida. 2000. Cereal wine (*jaanr*) and distilled wine (*naksi*) in Sikkim. *Journal of Brewing Society of Japan* 95: 115–122.
- Kozaki, M. and T. Uchimura. 1990. Microorganisms in Chinese starter “bubod” and rice wine “tapuy” in the Philippines. *Journal of Brewing Society of Japan* 85(11): 818–824.
- Kubo, Y., Rooney, A.P., Tsukakoshi, Y., Nakagawa, R., Hasegawa, H., and K. Kimura. 2011. Phylogenetic analysis of *Bacillus subtilis* strains applicable to natto (fermented soybean) production. *Applied Environmental Microbiology* 77: 6463–6469.
- Kurtzman, C.P., Fell JW, and T. Boekhout (Editor), 2011. *The Yeasts: A Taxonomic Study*, 5th edition. Elsevier: London.



- Kurtzman, C.P. and C.J. Robnett. 2003. Phylogenetic relationship among yeasts of the “*Saccharomyces* complex” determined from multigene sequence analyses. *FEMS Yeast Research* 3: 417–432.
- Kurtzman, C.P., Robnett, C.J., and Basehoar-Powers, E. 2001. *Zygosaccharomyces kombuchaensis*, a new ascosporogenous yeast from “Kombucha tea”. *FEMS Yeast Research* 1: 133–138.
- Kuda, T., Izawa, Y., Yoshida, S., Koyanagi, T., Takahashi, H., and B. Kimura. 2014. Rapid identification of *Tetragenococcus halophilus* and *Tetragenococcus muriaticus*, important species in the production of salted and fermented foods, by matrix-assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF MS). *Food Control* 35(1): 419–425.
- Kutyauripo, J., Parawira, W., Tinofa, S., Kudita, I., and C. Ndengu. 2009. Investigation of shelf-life extension of sorghum beer (*Chibuku*) by removing the second conversion of malt. *International Journal of Food Microbiology* 129: 271–276.
- Kwon, G.H., Lee, H.A., Park, J.Y., Kim, J.S., Lim, J., Park, C.S., Kwon, D.Y., and J.H. Kim. 2009. Development of a RAPD-PCR method for identification of *Bacillus* species isolated from Cheonggukjang. *International Journal of Food Microbiology* 129: 282–287.
- Lactic Acid Bacteria. 2013. <http://www.metamicrobe.com/lactobacteria/>
- Lappe-Oliveras, P., Moreno-Terrazas, R., Arrizón-Gaviño, J., Herrera-Suárez, T., Garcia-Mendoza, A., and A. Gschaedler-Mathis. 2008. Yeasts associated with the production of Mexican alcoholic nondistilled and distilled *Agave* beverages. *FEMS Yeast Research* 8: 1037–1052.
- Lee, C.H. 1993. Fish fermentation technology in Korea. In: *Fish Fermentation Technology* (Eds: Lee, C.H., Steinkraus, K.H. and Alan Reilly, P.J.), pp. 187–201. United Nations University Press, Tokyo.
- Lee, C.H. 1995. An introduction to Korean food culture. *Korean and Korean American Studies Bulletin* 6(1): 6–10.
- Lee, S.W. 1984. *Hankuk sikpum munhwasa (Korean Dietary Culture)*. Seoul: Kyomunsa. (in Korean).
- Lee, I.H., Hung, Y.H., and C.C. Chou. 2007. Total phenolic and anthocyanin contents, as well as antioxidant activity, of black bean *koji* fermented by *Aspergillus awamori* under different culture conditions. *Food Chemistry* 3: 936–942.
- Lee, Z.S. and T.W. Rhee. 1970. Studies on the microflora of Takju brewing. *Korean Journal of Microbiology* 8: 116–133.
- Lefebvre, T., Janssens, M., Camu, N., and L. De Vuyst. 2010. Kinetic analysis of strains of lactic acid bacteria and acetic acid bacteria in cocoa pulp simulation media toward development of a starter culture for cocoa bean fermentation. *Applied Environmental Microbiology* 76: 7708–7716.
- Lepage, P., Leclerc, M.C., Joossens, M., Mondot, S., Blottière, H.M., Raes, J., Ehrlich, D., and J. Doré. 2013. A metagenomic insight into our gut’s microbiome *Gut* 62(1): 146–158.
- Leroy, F. and L. De Vuyst. 2004. Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science and Technology* 15: 67–78.
- Lindqvist, R. and M. Lindblad. 2009. Inactivation of *Escherichia coli*, *Listeria monocytogenes* and *Yersinia enterocolitica* in fermented sausages during maturation/storage. *International Journal of Food Microbiology* 129: 59–67.
- Longo, M.A. and M.A. Sanromán. 2006. Production of food aroma compounds: Microbial and enzymatic methodologies. *Food Technology and Biotechnology* 44: 335–353.
- Lopetcharat, K., Choi, Y.J., Park, J.W., and M.A. Daeschel. 2001. Fish sauce products and manufacturing: A review. *Food Reviews International* 17: 65–88.
- Lücke, F.K. 1985. Fermented sausages. In: *Microbiology of Fermented Foods*, vol. 2, ed. B.J.B. Wood, 41–83. London: Elsevier Applied Science Publishers.
- Lv, X-C., Huang, X-L., Zhang, W., Rao, P-F., and L. Ni. 2013. Yeast diversity of traditional alcohol fermentation starters for Hong Qu glutinous rice wine brewing, revealed by culture-dependent and culture-independent methods. *Food Control* 34: 183–190.
- Mabesa, R.C. and J.S. Babaan. 1993. Fish fermentation technology in the Philippines. In: *Fish Fermentation Technology* (Eds: Lee, C.H., Steinkraus, K.H., and Alan Reilly, P.J.), pp. 85–94. United Nations University Press, Tokyo.
- Marsh, A.J., O’Sullivan, O., Hill, C., R. Paul Ross, R.P., and Cotter, D. 2014. Sequence-based analysis of the bacterial and fungal compositions of multiple kombucha (tea fungus) samples. *Food Microbiology* 38: 171–178.

- Martín, B., Garriga, M., Hugas, M., Bover-Cid, S., Veciana-Noqués, M.T., and Aymerich, T. 2006. Molecular, technological and safety characterization of Gram-positive catalase-positive cocci from slightly fermented sausages. *International Journal of Food Microbiology* 107: 148–158.
- Marty, E., Buchs J, Eugster-Meier E, Lacroix, C., and L. Meile. 2011. Identification of staphylococci and dominant lactic acid bacteria in spontaneously fermented Swiss meat products using PCR–RFLP. *Food Microbiology* 29: 157–166.
- Masoud, W., Cesar, L.B., Jespersen, L., and M. Jakobsen. 2004. Yeast involved in fermentation of *Coffea arabica* in East Africa determined by genotyping and direct denaturing gradient gel electrophoresis. *Yeast* 21: 549–556.
- Mathara, J.M., Schillinger, U., Kutima, P.M., Mbugua, S.K., and W.H. Holzapfel. 2004. Isolation, identification and characterisation of the dominant microorganisms of kule naoto: The Maasai traditional fermented milk in Kenya. *International Journal of Food Microbiology* 94: 269–278.
- Matsushita, M., Tada, S., Suzuki, S., Kusumoto, K., and Y. Kashiwagi. 2009. Deletion analysis of the promoter of *Aspergillus oryzae* gene encoding heat shock protein 30. *Journal of Bioscience and Bioengineering* 107: 345–351.
- Mayo, B., Ammor, M.S., Delgado, S., and A. Alegría. 2010. Fermented milk products. In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 263–288.
- Meerak, J., Iida, H., Watanabe, Y., Miyashita, M., Sato, H., Nakagawa, Y., and Y. Tahara. 2007. Phylogeny of  $\gamma$ -polyglutamic acid-producing *Bacillus* strains isolated from fermented soybean foods manufactured in Asian countries. *Journal of General and Applied Microbiology* 53: 315–323.
- Meerak, J., Yukphan, P., Miyashita, M., Sato, H., Nakagawa, Y., and Y. Tahara. 2008. Phylogeny of  $\gamma$ -polyglutamic acid-producing *Bacillus* strains isolated from a fermented locust bean product manufactured in West Africa. *Journal of General and Applied Microbiology* 54: 159–166.
- Merican, Z. 1996. Malaysian pickles. In: *Handbook of Indigenous Fermented Food*, 2nd edition, ed. K.H. Steinkraus, 138–141. New York: Marcel Dekker, Inc.
- Merican, Z. 1977. Budu (fish sauce). In: *The Proceedings of the Symposium on Indigenous Fermented Foods*, November 21–27, 1977, GIAM-V, Bangkok.
- Merican, Z. and Q.L. Yeoh. 1989. Tapai proceeding in Malaysia: A technology in transition. In: *Industrialization of Indigenous Fermented Foods*, ed. K.H. Steinkraus, 169–189. New York: Marcel Dekker, Inc.
- Mitsuoka, T. 1992. Intestinal flora and aging. *Nutrition Review* 50: 438–446.
- Mo, H., Zhu, Y., and Z. Chen. 2008. Microbial fermented tea—A potential source of natural food preservatives. *Trends in Food Science and Technology* 19: 124–130.
- Mohammed, S.I., Steenson, L.R., and A.W. Kirleis. 1991. Isolation and characterization of microorganisms associated with the traditional sorghum fermentation for production of Sudanese kisra. *Applied Environmental Microbiology* 57: 2529–2533.
- Mogensen, G., Salminen, S., O'Brien, J., Ouwehand, A., Holzapfel, W., Shortt, C., Fonden, R. et al. 2002. Inventory of micro-organisms with a documented history of use in food. *Bulletin of IDF* 377: 10–19.
- Moreira, N., Mendes, F., Hogg, T., and I. Vasconcelos. 2005. Alcohols, esters and heavy sulphur compounds produced by pure and mixed cultures of apiculture wine yeasts. *International Journal of Food Microbiology* 103: 285–294.
- Moroni, A.V., Arendt, E.K., and F.D. Bello. 2011. Biodiversity of lactic acid bacteria and yeasts in spontaneously-fermented buckwheat and teff sourdoughs. *Food Microbiology* 28: 497–502.
- Mozzi, F., Eugenia Ortiz, M., Bleckwedel, J., De Vuyst, L., and P. Micaela. 2013. Metabolomics as a tool for the comprehensive understanding of fermented and functional foods with lactic acid bacteria. *Food Research International* 54: 1152–1161.
- Mshvildadze, M. and J. Neu. 2010. The infant intestinal microbiome: Friend or foe? *Early Human Development* 86: 67–71.
- Mugula, J.K., Ninko, S.A.M., Narvhus, J.A., and T. Sorhaug. 2003. Microbiological and fermentation characteristics of *togwa*, a Tanzanian fermented food. *International Journal of Food Microbiology* 80: 187–199.
- Murray, M.A. 2000. Viticulture and wine production. In: *Ancient Egyptian Materials and Technology*, eds. P.T. Nicholson and I. Shaw, 577–608. Cambridge: Cambridge University Press.

- Nagai, T., Koguchi, K., and Y. Itoh. 1997. Chemical analysis of poly- $\gamma$ -glutamic acid produced by plasmid-free *Bacillus subtilis* (*natto*): evidence that plasmids are not involved in poly- $\gamma$ -glutamic acid production. *Journal of General and Applied Microbiology* 43: 139–143.
- Nagai, T. and J.P. Tamang. 2010. Fermented soybeans and non-soybeans legume foods. In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*. New York: CRC Press, Taylor & Francis Group, pp. 191–224.
- Nagai, T., Tran, L.S., Inatsu, Y., and Y. Itoh. 2000. A new IS4 family insertion sequence, IS4*Bsu1*, responsible for genetic instability of poly- $\gamma$ -glutamic acid production in *Bacillus subtilis*. *Journal of Bacteriology* 182: 2387–2392.
- Nakao, S. 1972. Mame no ryori. In: *Ryori no kigen*. Tokyo: Japan Broadcast Publishing, Tokyo (in Japanese), 115–126.
- Nam, Y-D., Chang, H.W., Kim, K.H., Roh, S.W., and J.W. Bae. 2009. Metatranscriptome analysis of lactic acid bacteria during *kimchi* fermentation with genome-probing microarrays. *International Journal of Food Microbiology* 130: 140–146.
- Nam, Y-D., Lee, So-Y., and S-I. Lim. 2011. Microbial community analysis of Korean soybean pastes by next-generation sequencing. *International Journal of Food Microbiology* 155(1–2): 36–42.
- Nam, Y-D., Yi, S-H., and S-I. Lim. 2012. Bacterial diversity of *cheonggukjang*, a traditional Korean fermented food, analyzed by barcoded pyrosequencing. *Food Control* 28(1): 135–142.
- Naser, S.M., Thompson, F.L., Hoste, B., Gevers, D., Dawyndt, P., Vancanneyt, M., and J. Swings. 2005. Application of multilocus sequence analysis (MLSA) for rapid identification of *Enterococcus* species based on *pheS* and *rpaA* genes. *Microbiology* 151: 2141–2150.
- Ngaba, R.R. and Lee, J.S. 1979. Fermentation of cassava. *Journal of Food Science* 44: 1570–1571.
- Nguyen, H., Elegado, F., Librojo-Basilio, N., Mabesa, R., and Dizon, E. 2011. Isolation and characterisation of selected lactic acid bacteria for improved processing of *nem chua*, a traditional fermented meat from Vietnam. *Beneficial Microbes* 1(1): 67–74.
- Nguyen, D.T.L., Van Hoorde, K., Cnockaert, M., de Brandt, E., de Bruyne, K., Le, B.T., and P. Vandamme. 2013a. A culture-dependent and -independent approach for the identification of lactic acid bacteria associated with the production of *nem chua*, a Vietnamese fermented meat product. *Food Research International* 50(1): 232–240.
- Nguyen, D.T. L., Van Hoorde, K., Cnockaert, M., de Brandt, E., Aerts, M., Thanh, L.B., and P. Vandamme. 2013b. A description of the lactic acid bacteria microbiota associated with the production of traditional fermented vegetables in Vietnam. *International Journal of Food Microbiology* 163(1): 19–27.
- Nielsen, D.S., Schillinger, U., Franz, C.M.A.P., Bresciani, J., Amoa-Awua, W., Holzapfel, W.H., and M. Jakobsen. 2007. *Lactobacillus ghanensis* sp. nov., a motile lactic acid bacterium isolated from Ghanaian cocoa fermentations. *International Journal of Systematic and Evolutionary Microbiology* 57: 1468–1472.
- Nikkuni, S., Karki, T.B., Terao, T., and C. Suzuki. 1996. Microflora of mana, a Nepalese rice koji. *Journal of Fermentation and Bioengineering* 81: 168–170.
- Nishito, Y., Osana, Y., Hachiya, T., Popendorf, K., Toyoda, A., Fujiyama, A., Itaya, M., and Sakakibara, Y. 2010. Whole genome assembly of a *natto* production strain *Bacillus subtilis natto* from very short read data. *BMC Genomics* 11: 243–255.
- Norman, J. M., Handley, S. A., and W. Virgin. 2014. Kingdom-agnostic metagenomics and the importance of complete characterization of enteric microbial communities. *Gastroenterology* 146(6): 1459–1469.
- Nout, M.J.R. and K.E. Aidoo. 2002. Asian fungal fermented food, In: *The Mycota*, ed. Osiewacz, H.D., Springer-Verlag, New York, 23–47.
- Nout, M.J.R. and J.L. Kiers. 2005. Tempeh fermentation, innovation and functionality: Update into the third millenium. *Journal of Applied Microbiology* 98: 789–805.
- Odufa, S.A. 1988. Review: African fermented foods: From art to science. *MIRCEN Journal* 4: 259–273.
- Odufa, S.A. and O.B. Oyewole. 1997. *African Fermented Foods*. Blackie Academic and Professional: London.
- Oguntoyinbo, F.A. and C.E.R. Dodd. 2010. Bacterial dynamics during the spontaneous fermentation of cassava dough in *gari* production. *Food Control* 21: 306–312.
- Oguntoyinbo, F.A., Huch, M., Cho, G.S., Schillinger, U., Holzapfel, W.H., Sanni, A.I., and C.M.A.P. Franz. 2010. Diversity of *Bacillus* species isolated from okpehe, a traditional fermented soup condiment from Nigeria. *Journal of Food Protection* 73: 870–878.

- Oguntoyinbo, F.A., Sanni, A.I., Franz, S.M.A.P., and W.H. Holzapfel. 2007. *In vitro* fermentation studies for selection and evaluation of *Bacillus* strains as starter cultures for the production of okpehe, a traditional African fermented condiment. *International Journal of Food Microbiology* 113: 208–218.
- Oguntoyinbo, F.A., Tourlomousis, P., Gasson, M.J., and Nrabad, A. 2011. Analysis of bacterial communities of traditional fermented West African cereal foods using culture independent methods. *International Journal of Food Microbiology* 145: 205–210.
- Okafor, N. and A.O. Ejiofor. 1990. Rapid detoxification of cassava mash fermenting for garri production following inoculation with a yeast simultaneously producing linamarase and amylase. *Proceeding of Biochemical Institute* 25: 82–86.
- Oki, K., Dugersuren, J., Demberel, S., and K. Watanabe. 2014. Pyrosequencing analysis of the microbial diversity of *airag*, *khoormog* and *tarag*, traditional fermented dairy products of Mongolia. *Bioscience of Microbiota, Food and Health* 33(2): 53–64.
- Oki, K., Kudo, Y., and K. Watanabe. 2012. *Lactobacillus saniviri* sp. nov. and *Lactobacillus senioris* sp. nov., isolated from human faeces. *International Journal of Systematic and Evolutionary Microbiology* 62: 601–607.
- Oki, K., Rai, A.K., Sato, S., Watanabe, K., and J.P. Tamang. 2011. Lactic acid bacteria isolated from ethnic preserved meat products of the Western Himalayas. *Food Microbiology* 28: 1308–1315.
- Olasupo, N.A., Odunfa, S.A., and O.S. Obayori. 2010. Ethnic African fermented foods, In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 323–352.
- Ongol, M.P. and K. Asano. 2009. Main microorganisms involved in the fermentation of Uganda ghee. *International Journal of Food Microbiology* 133: 286–291.
- Ouoba, L., Cantor, M., Diawara, B., Traoré, A., and Jakobsen, M., 2003a. Degradation of African locust bean oil by *Bacillus subtilis* and *Bacillus pumilus* isolated from soumbala, a fermented African locust bean condiment. *Journal of Applied Microbiology* 95: 868–873.
- Ouoba, L., Diawara, B., Annan, N., Poll, L., and Jakobsen, M., 2005. Volatile compounds of Soumbala, a fermented African locust bean (*Parkia biglobosa*) food condiment. *Journal of Applied Microbiology* 99: 1413–1421.
- Ouoba, L.I.I., Diawara, B., Christensen, T., Dalgaard Mikkelsen, J., and Jakobsen, M., 2007a. Degradation of polysaccharides and non-digestible oligosaccharides by *Bacillus subtilis* and *Bacillus pumilus* isolated from Soumbala, a fermented African locust bean (*Parkia biglobosa*) food Condiment. *European Food Research and Technology* 224: 689–694.
- Ouoba, L.I.I., Diawara, B., Jespersen, L., and Jakobsen, M., 2007b. Antimicrobial activity of *Bacillus subtilis* and *Bacillus pumilus* during the fermentation of African locust bean (*Parkia biglobosa*) for Soumbala production. *Journal of Applied Microbiology* 102: 963–970.
- Ouoba, L.I., Nyanga-Koumou, C.A., Parkouda, C., Sawadogo, H., Kobawila, S.C., Keleke, S., Diawara, B., Louembe, D., and J.P. Sutherland. 2010. Genotypic diversity of lactic acid bacteria isolated from African traditional alkaline-fermented foods. *Journal of Applied Microbiology* 108: 2019–2029.
- Ouoba, L., Parkouda, C., Diawara, B., Scotti, C., and A. Varnam. 2008. Identification of *Bacillus* spp. from Bikalga, fermented seeds of *Hibiscus sabdariffa*: Phenotypic and genotypic characterization. *Journal of Applied Microbiology* 104: 122–131.
- Ouoba, L., Rechinger, K., Barkholt, V., Diawara, B., Traore, A., and Jakobsen, M., 2003b. Degradation of proteins during the fermentation of African locust bean (*Parkia biglobosa*) by strains of *Bacillus subtilis* and *Bacillus pumilus* for production of Soumbala. *Journal of Applied Microbiology* 94: 396–402.
- Ouoba, L., Diawara, B., Wk, A.A., Traore, A., and Moller, P., 2004. Genotyping of starter cultures of *Bacillus subtilis* and *Bacillus pumilus* for fermentation of African locust bean (*Parkia biglobosa*) to produce Soumbala. *International Journal of Food Microbiology* 90: 197–205.
- Oyewole, O.B., Olatunji, O.O., and S.A. Odunfa. 2004. A process technology for conversion of dried cassava chips into “gari”. *Nigeria Food Journal* 22: 65–76.
- Papalexandratou, Z., Vrancken, G., De Bruyne, K., Vandamme, P., and L. De Vuyst. 2011. Spontaneous organic cocoa bean box fermentations in Brazil are characterized by a restricted species diversity of lactic acid bacteria and acetic acid bacteria. *Food Microbiology* 28: 1326–1338.
- Parente, E. and Cogan, T.M. 2004. Starter cultures: general aspects, In: *Cheese: Chemistry, Physics and Microbiology*, 3rd edition, ed. Fox, P. O., Elsevier, Oxford, 123–147.



- Park, E.J., Chang, H.W., Kim, K.H., Nam, Y.D., Roh, S.W., and J.W. Bae. 2009. Application of quantitative real-time PCR for enumeration of total bacterial, archaeal, and yeast populations in kimchi. *Journal Microbiology* 47: 682–685.
- Park, C., Choi, J.C., Choi, Y.H., Nakamura, H., Shimanouchi, K., Horiuchi, T., Misono, H., Sewaki, T., Soda, K., Ashiuchi M., and M.H. Sung. 2005. Synthesis of super-high-molecular-weight poly- $\gamma$ -glutamic acid by *Bacillus subtilis* subsp. *chungkookjang*. *Journal of Molecular Catalysis B: Enzymatic* 35: 128–133.
- Park, E.J., Chun, J., Cha, C.J., Park, W.S., Jeon, C.O., and J.W. Bae. 2012. Bacterial community analysis during fermentation of ten representative kinds of *kimchi* with barcoded pyrosequencing. *Food Microbiology* 30: 197–204.
- Park, J.M., Shin, J.H., Lee, D.W., Song, J.C., Suh, H.J., Chang, U.J., and J.M. Kim. 2010. Identification of the lactic acid bacteria in kimchi according to initial and over-ripened fermentation using PCR and 16S rRNA gene sequence analysis. *Food Science and Biotechnology* 19: 541–546.
- Parkouda C, Nielsen, D.S., Azokpota, P., Ouoba, L.I.I., Amoa-Awua, W.K., Thorsen, L., Houhouigan, J.D., Jensen, J.S., Tano-Debrah, K., Diawara, B., and M. Jakobsen. 2009. The microbiology of alkaline-fermentation of indigenous seeds used as food condiments in Africa and Asia. *Critical Review in Microbiology* 35: 139–156.
- Parente, E.S., Di Matteo, M., Spagna Musso, S., and M.A. Crudele. 1994. Use of commercial starter cultures in the production of soppressa lucana, a fermented sausage from Basilicata. *Italian Journal of Science* 6: 59–69.
- Parente, E.S., Grieco, S., and M.A. Crudele. 2001a. Phenotypic diversity of lactic acid bacteria isolated from fermented sausages produced in Basilicata (Southern Italy). *Journal of Applied Microbiology* 90: 943–952.
- Parente, E., Martuscelli, M., Gardini, F., Grieco, S., Crudele, M.A., and G. Suzzi. 2001b. Evolution of microbial populations and biogenic amine production in dry sausages produced in Southern Italy. *Journal of Applied Microbiology* 90: 882–891.
- Patidar, S.K. and J.B. Prajapati. 1998. Standardization and evaluation of lassi prepared using *Lactobacillus acidophilus* and *Streptococcus thermophilus*. *Journal of Food Science and Technology* 35(5): 428–431.
- Patil, M.M., Pal, A., Anand, T., and K.V. Ramana. 2010. Isolation and characterization of lactic acid bacteria from curd and cucumber. *Indian Journal of Biotechnology* 9: 166–172.
- Patrignani, F., Lanciotti R, Mathara J.M., Guerzoni M.E., and W.H. Holzapfel. 2006. Potential of functional strains, isolated from traditional *Maasai* milk, as starters for the production of fermented milks. *International Journal of Food Microbiology* 107: 1–11.
- Pederson, C.S. 1979. *Microbiology of Food Fermentations*, 2nd edition. Westport: AVI Publishing Company.
- Pederson, C.S. and M.N. Albury. 1969. The sauerkraut fermentation. *Food Technology* 8: 1–5.
- Phithakpol, B. 1993. Fish fermentation technology in Thailand. In: *Fish Fermentation Technology* (Eds: Lee, C.H., Steinkraus, K.H., and Alan Reilly, P.J.), pp. 155–166. United Nations University Press, Tokyo.
- Phithakpol, B., Varayanond, W., Reungmaneepeatoon, S., and H. Wood. 1995. *The Traditional Fermented Foods of Thailand*. Kuala Lumpur: ASEAN Food Handling Bureau.
- Picozzi, C., Bonacina, G., Vigentini, I., and Foschino, R. 2010. Genetic diversity in Italian *Lactobacillus sanfranciscensis* strains assessed by multilocus sequence typing and pulsed field gel electrophoresis analyses. *Microbiology* 156: 2035–2045.
- Plengvidhya, V., Breidt, F., and H.P. Fleming. 2007. Use of RAPD-PCR as a method to follow the progress of starter cultures in sauerkraut fermentation. *International Journal of Food Microbiology* 93: 287–296.
- Prakash, O. 1961. *Food and Drinks in Ancient India*. Delhi: Munshi Ram Monoharlal Publ.
- Prajapati, J.B. and B.M. Nair. 2003. The history of fermented foods. In: *Handbook of Fermented Functional Foods*, ed. R. Farnworth, 1–25. New York: CRC Press.
- Pretorius, I.S. 2000. Review: Tailoring wine yeast for the new millennium: Novel approaches to the ancient art of wine making. *Yeast* 16: 675–729.
- Purchiaroni, F., Tortora, A., Gabrielli, M., Bertucci, F., Gigante, G., Ianiro, G., Ojetti, V., Scarpellini, E., and A. Gasbarrini. 2013. The role of intestinal microbiota and the immune system. *European Review for Medical and Pharmacological Sciences* 17: 323–333.
- Puspito, H. and G.H. Fleet. 1985. Microbiology of *sayur asin* fermentation. *Applied Microbiology and Biotechnology* 22: 442–445.
- Putro, S. 1993. Fish fermentation technology in Indonesia. In: *Fish Fermentation Technology* (Eds: Lee, C.H., Steinkraus, K.H., and Alan Reilly, P.J.), pp. 107–128. United Nations University Press, Tokyo.

- Quigley, L., O'Sullivan, O., Beresford, T.P., Ross, R.P., Fitzgerald, G.F., and P.D. Cotter. 2011. Molecular approaches to analysing the microbial composition of raw milk and raw milk cheese. *International Journal of Food Microbiology* 150: 81–94.
- Rai, A.K., Palni, U., and J.P. Tamang. 2009. Traditional knowledge of the Himalayan people on production of indigenous meat products. *Indian Journal of Traditional Knowledge* 8(1): 104–109.
- Rai, A.K., Palni, U., and J.P. Tamang. 2010. Microbiological studies of ethnic meat products of the Eastern Himalayas. *Meat Science* 85: 560–567.
- Ramakrishnan, C.V. 1979. Studies on Indian fermented foods. *Baroda Journal of Nutrition* 6: 1–54.
- Ramos, C.L., de Almeida, E.G., de Melo Pereira, G.V., Cardoso, P.G., Dias, E.S., and R.F. Schwan. 2010. Determination of dynamic characteristics of microbiota in a fermented beverage produced by Brazilian Amerindians using culture-dependent and culture-independent methods. *International Journal of Food Microbiology* 140(2–3): 225–231.
- Rani, D.K. and S.K. Soni. 2007. Applications and commercial uses of microorganisms. In: *Microbes: A Source of Energy for 21st Century*, ed. Soni, S.K., Jai Bharat Printing Press, Tohtas Nagar, Shahdara, Delhi. Chapter 2, pp. 71–126.
- Rantsiou, K. and L. Coccolin. 2006. New developments in the study of the microbiota of naturally fermented sausages as determined by molecular methods: A review. *International Journal of Food Microbiology* 108: 255–267.
- Rasic, J.L. and J.A. Kurmann. 1978. *Yoghurt—Scientific Grounds, Technology, Manufacture and Preparations*. Copenhagen: Technical Dairy Publishing House.
- Renfrew, J.W. 1999. Palaeoethnobotany and the archaeology of wine. In: *The Oxford Companion to Wine*, 2nd edition, ed. J. Robinson, 508–509. Oxford: Oxford University Press.
- Rhee, S.J., Lee, J.E., and C.H. Lee. 2011. Importance of lactic acid bacteria in Asian fermented foods. *Microbial Cell Factories* 10: 1–13.
- Risley, H.H. 1928. *The Gazetteer of Sikkim*. New Delhi: D. K. Publishing Distributors (P) Ltd.
- Robert, H., Gabriel, V., and C. Fontagné-Faucher. 2009. Biodiversity of lactic acid bacteria in French wheat sourdough as determined by molecular characterization using species-specific PCR. *International Journal of Food Microbiology* 135: 53–59.
- Robinson, R.K. and A.Y. Tamime. 2006. Types of fermented milks, In: *Fermented Milks*, Tamime, A.Y. ed. Tamime, Wiley, Hoboken, 1–10.
- Romano, P., Capace, A., and L. Jespersen. 2006. Taxonomic and ecological diversity of food and beverage yeasts, In: Querol, A., Fleet, G.H. eds. *The Yeast Handbook—Yeasts in Food and Beverages*, Springer-Verlag, Berlin, 13–53.
- Romano-Keeler, J., Moore, D.J., Wang, C., Brucker, R.M., Fonnesbeck, C., Slaughter, J.C., Li, H., Curran, D.P., Meng, S., and H. Correa et al. 2014. Early life establishment of site-specific microbial communities in the gut. *Gut Microbes* 5: 192–201.
- Rossi, L.P.R., Almeida, R.C.C., Lopes, L.S., Figueiredo, A.C.L., Ramos, M.P.P., and P.F. Almeida. 2011. Occurrence of *Listeria* spp. in Brazilian fresh sausage and control of *Listeria monocytogenes* using bacteriophage P100. *Food Control* 22: 954–958.
- Ruddle, K. 1993. The availability and supply of fish for fermentation in Southeast Asia. In: *Fish Fermentation Technology*, eds. C.-H. Lee, K.H. Steinkraus and P.J. Alan Reilly, 45–84. Tokyo: United Nations University Press.
- Sakai, H., Caldo, G. A., and M. Kozaki. 1983a. Yeast-flora in red *burong-isda* a fermented fish food from the Philippines. *Journal of Agricultural Science (Tokyo)* 28: 181–185.
- Sakai, H., Caldo, G. A., and M. Kozaki. 1983b. The fermented fish food, *burong-isda*, in the Philippines. *Journal of Agricultural Science (Tokyo)* 28: 138–144.
- Saisithi, P. 1987. Traditional fermented fish products with special reference to Thai products. *ASEAN Food Journal* 3: 3–10.
- Saithong, P., Panthavee, W., Boonyaratanakornkit, M., and C. Sikkhamondhol. 2010. Use of a starter culture of lactic acid bacteria in *plaa-som*, a Thai fermented fish. *Journal of Bioscience and Bioengineering* 110: 553–557.
- Sakamoto, N., Tanaka, S., Sonomoto, K., and J. Nakayama. 2011. 16S rRNA pyrosequencing-based investigation of the bacterial community in nukadoko, a pickling bad of fermented rice bran. *International Journal of Food Microbiology* 144: 352–359.

- Salampeyy, J., Kailasapathy, K., and N. Thapa. 2010. Fermented fish products. In: Tamang, J.P. and Kailasapathy, K. eds. *Fermented Foods and Beverages of the World*. New York: CRC Press, Taylor & Francis Group, pp. 289–307.
- Salminen, S., Wright, A.V., and A. Ouwehand. 2004. *Lactic Acid Bacteria Microbiology and Functional Aspects*, 3rd edition, New York: Marcel Dekker.
- Samuel, D. 1996. Archaeology of ancient Egyptian beer. *Journal of American Society of Brewing Chemists* 54(1): 3–12.
- Samuel, D. 2000. Brewing and baking. In: *Ancient Egyptian Materials and Technology*, eds. P.T. Nicholson and I. Shaw, 537–576. Cambridge: Cambridge University Press.
- Samuel, D. 2002. Bread in archaeology. *Civilisations* 49: 28–36.
- Sarkar, P.K., Hasenack, B., and M.J.R. Nout. 2002. Diversity and functionality of *Bacillus* and related genera isolated from spontaneously fermented soybeans (Indian Kinema) and locust beans (African Soumbala). *International Journal of Food Microbiology* 77: 175–186.
- Sarkar, S. 2008. Innovations in Indian fermented milk products—A review. *Food Biotechnology* 22(1): 78–97.
- Sarkar, P.K. and J.P. Tamang. 1994. The influence of process variables and inoculum composition on the sensory quality of kinema. *Food Microbiology* 11: 317–325.
- Sarkar, P.K., Tamang, J.P., Cook, P.E., and J.D. Owens. 1994. Kinema—A traditional soybean fermented food: Proximate composition and microflora. *Food Microbiology* 11: 47–55.
- Sarma, P.J. 1939. The art of healing. *Rigveda Annals of Medical History*, 3rd Series: 1, 538.
- Sato, H., Torimura, M., Kitahara, M., Ohkuma, M., Hotta, Y., and H. Tamura. 2012. Characterization of the *Lactobacillus casei* group based on the profiling of ribosomal proteins coded in S10-spc-alpha operons as observed by MALDI-TOF MS. *Systematic and Applied Microbiology* 35: 447–454.
- Savado, A., Tapi, A., Chollet M, Wathelet B, Traoré A.S., and Jacques, P. 2011. Identification of surfactin producing strains in *Soumbala* and *Bikalga* fermented condiments using polymerase chain reaction and matrix assisted laser desorption/ionization-mass spectrometry methods. *International Journal of Food Microbiology* 151: 299–306.
- Sawamura, S. 1906. On the micro-organisms of natto. Bulletin College Agri. Tokyo Imperial University, 7, 107–110.
- Schillinger, U., Ban-Koffi, L., and C.M.A.P. Franz. 2010. Tea, coffee and cacao. In: *Fermented Foods and Beverages of the World*, eds. J.P. Tamang and K. Kailasapathy, 353–375. New York: CRC Press, Taylor & Francis Group.
- Schillinger, U., Böhringer, B., Wallbaum, S., Caroline, L., Gonfa, A., Kostinek, M., Holzapfel W.H., and Franz, C.M.A.P. 2008. A genus-specific PCR for differentiation between *Leuconostoc* and *Weissella* and its application in identification of heterofermentative lactic acid bacteria from coffee fermentation. *FEMS Microbiology Letters* 286: 222–226.
- Schillinger, U., Yousif, N.M.K., Sesar, L., and C.M.A.P. Franz. 2003. Use of group-specific and RAPD-PCR analyses for rapid differentiation of *Lactobacillus* strains from probiotic yogurts. *Current Microbiology* 47: 453–456.
- Schuijt, T.J., Poll, T., de Vos, W.M., and W.J. Wiersinga. 2013. Human microbiome: The intestinal microbiota and host immune interactions in the critically ill. *Trends in Microbiology* 21: 221–229.
- Schuller, D., Pereira, L., Alves, H., Cambon, B., Dequin, S., and M. Casal. 2007. Genetic characterization of commercial *Saccharomyces cerevisiae* isolates recovered from vineyard environment. *Yeast* 24: 625–636.
- Scott, R. 1986. *Cheese Making Practice*, 2nd edition. London: Elsevier.
- Sealey, P.R. and P.A. Tyers. 1989. Olives from Roman Spain: A unique amphora find in British waters. *The Antiquaries Journal* 69: 54–72.
- Sekwati-Monang, B. and M.G. Gänzle. 2011. Microbiological and chemical characterisation of *ting*, a sorghum-based sourdough product from Botswana. *International Journal of Food Microbiology* 150: 115–121.
- Sengun, I.Y. and S. Karabiyikli. 2011. Importance of acetic acid bacteria in food industry. *Food Control* 22: 647–665.
- Sengun, I.Y., Nielsen, D.S., Karapinar, M., and M. Jakobsen. 2009. Identification of lactic acid bacteria isolated from Tarhana, a traditional Turkish fermented food. *International Journal of Food Microbiology* 135: 105–111.

- Sha, S.P., Thakur, N., Tamang, B., and J.P. Tamang. 2012. *Haria*, a traditional rice fermented alcoholic beverage of West Bengal. *International Journal of Agriculture Food Science and Technology* 3(2): 157–160.
- Shi, Z., Zhang, Y., Phillips, G.O., and Yang, G. 2014. Utilization of bacterial cellulose in food. *Food Hydrocolloids* 35: 539–545.
- Shin, M.S., Han, S.K., Ryu, J.S., Kim, K.S., and W.K. Lee. 2008. Isolation and partial characterization of a bacteriocin produced by *Pediococcus pentosaceus* K23–2 isolated from kimchi. *Journal of Applied Microbiology* 105: 331–339.
- Shin, D., Kwon, D., Kim, Y., and D. Jeong. 2012. *Science and Technology of Korean Gochujang*. Seoul: Public Health Edu, pp. 10–133.
- Shon, M.Y., Lee, J., Choi, J.H., Choi, S.Y., Nam, S.H., Seo, K.I., Lee, S.W., Sung, N.J., and S.K. Park. 2007. Antioxidant and free radical scavenging activity of methanol extract of chungkukjang. *Journal of Food Composition and Analysis* 20: 113–118.
- Shrestha, H., Nand, K., and E.R. Rati. 2002. Microbiological profile of *murcha* starters and physico-chemical characteristics of *poko*, a rice based traditional food products of Nepal. *Food Biotechnology* 16: 1–15.
- Shurtleff, W. and A. Aoyagi. 2010. History of soybeans and soyfoods in South Asia/Indian subcontinent (1656–2010): Extensively annotated, bibliography and sourcebook. [www.soyinfocenter.com/books](http://www.soyinfocenter.com/books).
- Silva, C.F., Batista, L.R., Abreu, L.M., Dias, E.S., and R.F. Schwan. 2008. Succession of bacterial and fungal communities during natural coffee (*Coffea arabica*) fermentation. *Food Microbiology* 25: 951–957.
- Shamala, T.R. and K.R. Sreekantiah. 1988. Microbiological and biochemical studies on traditional Indian palm wine fermentation. *Food Microbiology* 5: 157–162.
- Simoncini, N., Rotelli, D., Virgili, R., and S. Quintavalla. 2007. Dynamics and characterization of yeasts during ripening of typical Italian dry-cured ham. *Food Microbiology* 24(6): 577–584.
- Singh, P.K. and K.I. Singh. 2006. Traditional alcoholic beverage, *Yu* of *Meitei* communities of Manipur. *Indian Journal of Traditional Knowledge* 5(2): 184–190.
- Singh, T.A., Devi, K.R., Ahmed, G., and K. Jeyaram. 2014. Microbial and endogenous origin of fibrinolytic activity in traditional fermented foods of Northeast India. *Food Research International* 55: 356–362.
- Solieri, L. and Giudici, P. 2008. Yeasts associated to traditional balsamic vinegar: Ecological and technological features. *International Journal of Food Microbiology* 125: 36–45.
- Sonar, R.N. and P.M. Halami. 2014. Phenotypic identification and technological attributes of native lactic acid bacteria present in fermented bamboo shoot products from North-East India. *Journal of Food Science and Technology* DOI 10.1007/s13197-014-1456-x.
- Soni, S.K., Sandhu, D.K., Vilku, K.S., and N. Kamra. 1986. Microbiological studies on dosa fermentation. *Food Microbiology* 3: 45–53.
- Sridevi, J., Halami, P.M., and Vijayendra, S.V.N. 2010. Selection of starter cultures for *idli* batter fermentation and their effect on quality of *idli*. *Journal of Food Science and Technology* 47: 557–563.
- Srinivasa, P.T.I. 1930. *Pre-Aryan Tamil Culture*. Madras: University of Madras.
- Steinkraus, K.H. 1996. *Handbook of Indigenous Fermented Food*, 2nd edition. New York: Marcel Dekker, Inc.
- Steinkraus K.H., 2004. *Industrialization of Indigenous Fermented Foods*. (Eds: Holzapfel, W.H. and Talijsaard, J.L.) Marcel Dekker, Inc, New York, Chapter 7, pp. 363–405.
- Steinkraus, K.H., van Veer, A.G., and D.B. Thiebeau. 1967. Studies on idli—An Indian fermented black gram-rice food. *Food Technology* 21: 110–113.
- Stevens, H.C. and L. Nabors. 2009. Microbial food cultures: A regulatory update. *Food Technology (Chicago)* 63: 36–41.
- Stiles, M.E. and W.H. Holzapfel. 1997. Lactic acid bacteria of foods and their current taxonomy. *International Journal of Food Microbiology* 36: 1–29.
- Suganuma, T., Fujita, K., and K. Kitahara. 2007. Some distinguishable properties between acid-stable and neutral types of  $\alpha$ -amylases from acid-producing *koji*. *Journal of Bioscience and Bioengineering* 104: 353–362.
- Sugawara, E. 2010. Fermented soybean pastes *miso* and *shoyu* with reference to aroma, In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 225–245.
- Sujaya, I., Antara, N., Sone, T., Tamura, Y., Aryanta, W., Yokota, A., Asano, K., and F. Tomita, F. 2004. Identification and characterization of yeasts in brem, a traditional Balinese rice wine. *World Journal of Microbiology and Biotechnology* 20: 143–150.



- Sukontasing, S., Tanasupawat, S., Moonmangmee, S., Lee, J.S., and K. Suzuki. 2007. *Enterococcus cameliae* sp. nov., isolated from fermented tea leaves in Thailand. *International Journal of Systematic and Evolutionary Microbiology* 57: 2151–2154.
- Sumino, T., Endo, E., Kageyama, A. S., Chihihara, R., and Yamada, K., 2003. Various Components and Bacteria of Furu (Soybean Cheese). *Journal of Cookery Science of Japan* 36: 157–163.
- Sun, S.Y., Gong, H.S., Jiang, X.M., and Y.P. Zhao. 2014. Selected non-*Saccharomyces* wine yeasts in controlled multistarter fermentations with *Saccharomyces cerevisiae* on alcoholic fermentation behaviour and wine aroma of cherry wines. *Food Microbiology* 44: 15–23.
- Sun, Z., Liu, W., Gao, W., Yang, M., Zhang, J., Wang, J., Menghe, B., Sun, T., and H. Zhang. 2010. Identification and characterization of the dominant lactic acid bacteria from *kurut*: The naturally fermented yak milk in Qinghai, China. *Journal of General and Applied Microbiology* 56: 1–10.
- Suprianto, Ohba, R., Koga, T., and S. Ueda. 1989. Liquefaction of glutinous rice and aroma formation in tapé preparation by ragi. *Journal of Fermentation and Bioengineering* 64(4): 249–252.
- Sura, K., Garg, S., and F.C. Garg. 2001. Microbiological and biochemical changes during fermentation of *Kanji*. *Journal of Food Science and Technology* 38: 165–167.
- Takahashi, M., Masaki, K., Mizuno, A., and N. Goto-Yamamoto. 2014. Modified COLD-PCR for detection of minor microorganisms in wine samples during the fermentation. *Food Microbiology* (39): 74–80.
- Takeda, S., Yamasaki K, Takeshita M, Kikuchi Y, Tsend-Ayush C, Dashnyam B, Ahhmed AM, Kawahara, S., and M. Muguét Ruma. 2011. The investigation of probiotic potential of lactic acid bacteria isolated from traditional Mongolian dairy products. *Animal Science Journal* 82: 571–579.
- Tamang, J.P. 1998. Role of microorganisms in traditional fermented foods. *Indian Food Industry* 17: 162–167.
- Tamang, J.P. 2001. Kinema. *Food Culture* 3: 11–14.
- Tamang, J.P. 2003. Native microorganisms in fermentation of *kinema*. *Indian Journal of Microbiology* 43: 127–130.
- Tamang, J.P. 2005. *Food Culture of Sikkim*. Sikkim Study Series volume IV. Information and Public Relations Department, Government of Sikkim, Gangtok, p. 120.
- Tamang, J.P. 2010a. *Himalayan Fermented Foods: Microbiology, Nutrition, and Ethnic Values*. New: CRC Press, Taylor & Francis Group.
- Tamang, J.P. 2010b. Diversity of fermented foods, In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 41–84.
- Tamang, J.P. 2010c. Diversity of fermented beverages, In: Tamang JP, Kailasapathy K. eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 85–125.
- Tamang, J.P. 2012a. Ancient food culture: History and ethnicity. In: *Globalisation and Cultural Practices in Mountain Areas: Dynamics, Dimensions and Implications*. ed. Lama, M.P., Sikkim University Press and Indus Publishing Company, New Delhi, pp. 43–63.
- Tamang, J.P. 2012b. Chapter 4. Plant-based fermented foods and beverages of Asia. In: *Handbook of Plant-Based Fermented Food and Beverage Technology*, Second Edition (Eds: Hui, Y.H. and Özgül, E.). CRC Press, Taylor & Francis Group, New York, pp. 49–90.
- Tamang, J.P. 2014. Biochemical and modern identification techniques—microfloras of fermented foods. In: *Encyclopaedia of Food Microbiology*, 2nd edition (Eds: Batt, C. and Tortorello, M.A.). Elsevier Ltd., Oxford, pp. 250–258.
- Tamang, J.P., Chettri, R., and R.M. Sharma. 2009. Indigenous knowledge of Northeast women on production of ethnic fermented soybean foods. *Indian Journal Traditional Knowledge* 8: 122–126.
- Tamang, J.P., Dewan, S., Tamang, B., Rai, A., Schillinger, U., and W.H. Holzapfel. 2007. Lactic acid bacteria in *Hamei* and *Marcha* of North East India. *Indian Journal of Microbiology* 47: 119–125.
- Tamang, J.P., Dewan, S., Thapa, S., Olasupo, N.A., Schillinger, U., Wijaya, A., and W.H. Holzapfel. 2000. Identification and enzymatic profiles of predominant lactic acid bacteria isolated from soft-variety *chhurpi*, a traditional cheese typical of the Sikkim Himalayas. *Food Biotechnology* 14: 99–112.
- Tamang, J.P. and G.H. Fleet. 2009. Yeasts diversity in fermented foods and beverages. In: Satyanarayana T, Kunze G. eds. *Yeasts Biotechnology: Diversity and Applications*, Springer, New York, 169–198.
- Tamang, J.P. and S. Nikkuni. 1996. Selection of starter culture for production of kinema, fermented soybean food of the Himalaya. *World Journal of Microbiology and Biotechnology* 12: 629–635.

- Tamang, J.P. and D. Samuel. 2010. Dietary culture and antiquity of fermented foods and beverages, In: J.P. Tamang and K. Kailasapathy, eds. *Fermented Foods and Beverages of the World*, CRC Press, Taylor & Francis Group, New York, 1–40.
- Tamang, J.P. and P.K. Sarkar. 1993. Sinki—A traditional lactic acid fermented radish tap root product. *Journal of General and Applied Microbiology* 39: 395–408.
- Tamang, J.P. and P.K. Sarkar. 1995. Microflora of murcha: An amylolytic fermentation starter. *Microbios* 81: 115–122.
- Tamang, J.P. and P.K. Sarkar. 1996. Microbiology of mesu, a traditional fermented bamboo shoot product. *International Journal of Food Microbiology* 29: 49–58.
- Tamang, J.P., Sarkar, P.K., and C.W. Hesseltine. 1988. Traditional fermented foods and beverages of Darjeeling and Sikkim—A review. *Journal of Science of Food and Agriculture* 44: 375–385.
- Tamang, B. and J.P. Tamang. 2007. Role of lactic acid bacteria and their functional properties in *Goyang*, a fermented leafy vegetable product of the Sherpas. *Journal of Hill Research* 20: 53–61.
- Tamang, B. and J.P. Tamang. 2009. Lactic acid bacteria isolated from indigenous fermented bamboo products of Arunachal Pradesh in India and their functionality. *Food Biotechnology* 23: 133–147.
- Tamang, B. and J.P. Tamang. 2010. *In situ* fermentation dynamics during production of *gundruk* and *khalpi*, ethnic fermented vegetables products of the Himalayas. *Indian Journal of Microbiology* 50(Suppl 1): 93–98.
- Tamang, B., Tamang, J.P., Schillinger, U., Franz, C.M.A.P., Gores, M., and W.H. Holzapfel. 2008. Phenotypic and genotypic identification of lactic acid bacteria isolated from ethnic fermented tender bamboo shoots of North East India. *International Journal of Food Microbiology* 121: 35–40.
- Tamang, J.P., Tamang, B., Schillinger, U., Franz, C.M.A.P., Gores, M., and W.H. Holzapfel. 2005. Identification of predominant lactic acid bacteria isolated from traditional fermented vegetable products of the Eastern Himalayas. *International Journal of Food Microbiology* 105: 347–356.
- Tamang, J.P., Tamang, N., Thapa, S., Dewan, S., Tamang, B.M., Yonzan, H., Rai, A.K., Chettri, R., Chakrabarty, J., and N. Kharel. 2012. Microorganisms and nutritional value of ethnic fermented foods and alcoholic beverages of North East India. *Indian Journal of Traditional Knowledge* 11(1): 7–25.
- Tamang, J.P., Thapa, S., Dewan, S., Jojima, Y., Fudou, R., and S. Yamanaka. 2002. Phylogenetic analysis of *Bacillus* strains isolated from fermented soybean foods of Asia: *Kinema*, *chungkokjang* and *natto*. *Journal of Hill Research* 15: 56–62.
- Tamang, J.P. and S. Thapa. 2006. Fermentation dynamics during production of bhaati jaanr, a traditional fermented rice beverage of the Eastern Himalayas. *Food Biotechnology* 20: 251–261.
- Tamang, J.P., Thapa, S., Tamang, N., and B. Rai. 1996. Indigenous fermented food beverages of the Darjeeling hills and Sikkim: Process and product characterization. *Journal of Hill Research* 9: 401–411.
- Tamime, A.Y. 2005. *Probiotic Dairy Products*. Oxford: Blackwell Publishing.
- Tamime, A.Y. and R.K. Robinson. 2007. *Yoghurt Science and Technology*. Cambridge: Woodhead Publishing Ltd.
- Tanaka, N., Traisman, E., Lee, M.S., Cassens, R.G., and E.M. Foster. 1980. Inhibition of botulinum toxin formation in bacon by acid development. *Journal of Food Protection* 43: 450–457.
- Tanasupawat, S., Pakdeeto, A., Thawai, C., Yukphan, P., and S. Okada. 2007. Identification of lactic acid bacteria from fermented tea leaves (miang) in Thailand and proposals of *Lactobacillus thailandensis* sp. nov., *Lactobacillus camelliae* sp. nov., and *Pediococcus siamensis* sp. nov. *Journal of General and Applied Microbiology* 53: 7–15.
- Tanganurat, W., Quinquis, B., Leelawatcharamas, V., and A. Bolotin. 2009. Genotypic and phenotypic characterization of *Lactobacillus plantarum* strains isolated from Thai fermented fruits and vegetables. *Journal of Basic Microbiology* 49: 377–385.
- Tanigawa, K., Kawabata, H., and K. Watanabe. 2010. Identification and typing of *Lactococcus lactis* by matrix-assisted laser desorption ionization—time-of-flight mass spectrometry. *Applied and Environmental Microbiology* 76: 4055–4062.
- Tanigawa, K. and K. Watanabe. 2011. Multilocus sequence typing reveals a novel subspeciation of *Lactobacillus delbrueckii*. *Microbiology* 157: 727–738.
- Tanimura, W., Sanchez, P.C., and M. Kozaki. 1977. The fermented food in the Philippines (Part 1) Tapuy (rice wine). *Journal of Agricultural Science of the Tokyo University of Agriculture* 22(1): 118–134.

- Tanimura, W., Sanchez, P.C., and M. Kozaki. 1978. The fermented foods in the Philippines. (Part-II). Basi (sugarcane wine). *Journal of Agricultural Society (Japan)* 22: 118–133.
- Taylor, J.R.N. 2003. Beverages from sorghum and millet. In: *Encyclopedia of Food Sciences and Nutrition*, 2nd edition, eds. B. Caballero, L.C. Trugo and P. M. Finglas, 2352–2359. London: Academic Press.
- Teoh, A.L., Heard, G., and J. Cox. 2004. Yeasts ecology of Kombucha fermentation. *International Journal of Food Microbiology* 95: 119–126.
- Thakur, N., Savitri, and T.C. Bhalla. 2004. Characterisation of some traditional fermented foods and beverages of Himachal Pradesh. *Indian Journal of Traditional Knowledge* 3(3): 325–335.
- Thanh, V.N., Mai, L.T., and D.A. Tuan. 2008. Microbial diversity of traditional Vietnamese alcohol fermentation starters (*banh men*) as determined by PCR-mediated DGGE. *International Journal of Food Microbiology* 128: 268–273.
- Thapa, N., Pal, J., and J.P. Tamang. 2004. Microbial diversity in *ngari*, *hentak* and *tungtap*, fermented fish products of Northeast India. *World Journal of Microbiology and Biotechnology* 20: 599–607.
- Thapa, N., Pal, J., and J.P. Tamang. 2006. Phenotypic identification and technological properties of lactic acid bacteria isolated from traditionally processed fish products of the Eastern Himalayas. *International Journal of Food Microbiology* 107: 33–38.
- Thapa, N., Pal, J., and J.P. Tamang. 2007. Microbiological profile of dried fish products of Assam. *Indian Journal of Fisheries* 54: 121–125.
- Thapa, S. and J.P. Tamang. 2004. Product characterization of kodo ko jaanr: Fermented finger millet beverage of the Himalayas. *Food Microbiology* 21: 617–622.
- Thapa, S. and J.P. Tamang. 2006. Microbiological and physico-chemical changes during fermentation of *kodo ko jaanr*, a traditional alcoholic beverage of the Darjeeling hills and Sikkim. *Indian Journal of Microbiology* 46: 333–341.
- Toldra, F. 2007. Handbook of fermented meat and poultry. Blackwell Publishing, Oxford.
- Topisirovic, L., Kojic, M., Fira, D., Golic, N., Strahinic, I., and J. Lozo. 2006. Potential of lactic acid bacteria isolated from specific natural niches in food production and preservation. *International Journal of Food Microbiology* 112: 230–235.
- Tou, E.H., Mouquet-River, C., Rochette, I., Traoré. A.S., Treche. S., and J.P. Guyot. 2007. Effect of different process combinations on the fermentation kinetics, microflora and energy density of *ben-saalga*, a fermented gruel from Burkina Faso. *Food Chemistry* 100: 935–943.
- Tsuyoshi, N., Fudou, R., Yamanaka, S., Kozaki, M., Tamang, N., Thapa, S., and J.P. Tamang. 2005. Identification of yeast strains isolated from marcha in Sikkim, a microbial starter for amylolytic fermentation. *International Journal of Food Microbiology* 99: 135–146.
- Tyn, M.T. 1993. Trends of fermented fish technology in Burma. In: *Fish Fermentation Technology* (Eds: Lee, C.H., Steinkraus, K.H. and Alan Reilly, P.J.), pp. 129–153. United Nations University Press, Tokyo.
- Uchimura, T., Okada, S., and M. Kozaki. 1991. Identification of lactic acid bacteria isolated from Indonesian Chinese starter, “ragi”. Microorganisms in Chinese starters from Asia (Part-4). *Journal of Brewing Society of Japan* 86: 55–61.
- Urushibata, Y., Tokuyama, S., and Y. Tahara. 2002. Characterization of the *Bacillus subtilis ywsC* gene, involved in  $\lambda$ -polyglutamic acid production. *Journal of Bacteriology* 184: 337–343.
- van Veen, A.G. 1965. Fermented and dried sea food products in South-East Asia. In: *Fish as Food*, volume 3 ed. Borgstrom, G., pp. 227–250. Academic Press, New York.
- Vachanavinich, K., Kim, W.J., and Y.I. Park. 1994. Microbial study on krachae, Thai rice wine, In: *Lactic Acid Fermentation of Non-Alcoholic Dairy Food and Beverages*, Lee CH, Adler-Nissen J, Bärwald G. eds. Ham Lim Won, Seoul, 233–246.
- Vallejo, J.A., Miranda, P., Flores-Félix, J.D., Sánchez-Juanes, F., Ageitos, J.M., González-Buitrago, J.M., Velázquez, E., and T.G. Villa. 2013. Atypical yeasts identified as *Saccharomyces cerevisiae* by MALDI-TOF MS and gene sequencing are the main responsible of fermentation of *chicha*, a traditional beverage from Peru. *Systematic and Applied Microbiology* 36(8): 560–564.
- van Hijum, S.A.F.T., Vaughan, E.E., and R.F. Vogel. 2013. Application of state-of-art sequencing technologies to indigenous food fermentations. *Current Opinion in Biotechnology* 24(2): 178–186.

- Vaughn, R.H. 1985. The microbiology of vegetable fermentations. In: *Microbiology of Fermented Foods*, vol. 1, (Ed: Wood, B.J.B.), pp. 49–109. Elsevier Applied Science Publishers, London.
- Veinocchi, P. Valmossi, S., Dalai, I., Torrsani, S., et Gianotti A, Suzzi G, Guerzoni ME, Mastrocola, D., and F. Gardini. 2006. Characterization of the yeast populations involved in the production of a typical Italian bread. *Journal of Food Science* 69: M182–M186.
- Versavaud, A., Coucoux, P., Roulland, C., Dulac, L., and J.N. Hallet. 1995. Genetic diversity and geographical distribution of wild *Saccharomyces cerevisiae* strains from the wine producing area of charentes, France. *Applied and Environmental Microbiology* 61: 3521–3529.
- Vieira-Dalodé, G., Jespersen L, Hounhouigan J, Moller PL, Nago C.M., and M. Jakobsen. 2007. Lactic acid bacteria and yeasts associated with *gowé* production from sorghum in Bénin. *Journal of Applied Microbiology* 103: 342–349.
- Vilanova, M. and C. Sieiro. 2006. Contribution by *Saccharomyces cerevisiae* yeast to fermentative flavour compounds in wines from cv. Albariño. *Journal of Industrial Microbiology and Biotechnology* 33: 929–933.
- Vogel, R.F., Hammes, W.P., Habermeyer, M., Engel, K.H., Knorr, D., and G. Eisenbrand. 2011. Microbial food cultures—Opinion of the Senate Commission on Food Safety (SKLM) of the German Research Foundation (DFG). *Molecular Nutrition and Food Research* 55: 654–662.
- Von Mollendorff, J.W. 2008. Characterization of bacteriocins produced by lactic acid bacteria from fermented beverages and optimization of starter cultures. MSc thesis, University of Stellenbosch, South Africa.
- Wacher, C., Díaz-Ruiz, G., and J.P. Tamang 2010. Fermented vegetable products. In: *Fermented Foods and Beverages of the World*, eds. J.P.Tamang and K. Kailasapathy, 149–190. New York: CRC Press, Taylor & Francis Group.
- Walker, G.M. 2014. Microbiology of Winemaking. In: *Encyclopaedia of Food Microbiology*, 2nd edition (Eds: Batt, C. and Tortorello, MA). Elsevier Ltd., Oxford, pp. 787–792.
- Wallace, T.C., Guarner, F., Madsen, K., Cabana, M.D., Gibson, G., Hentges, E., and M.E. Sanders. 2011. Human gut microbiota and its relationship to health and disease. *Nutrition Reviews* 69(7): 392–403.
- Wang, J. and D.Y.C. Fung. 1996. Alkaline-fermented foods: A review with emphasis on pidan fermentation. *Critical Review in Microbiology* 22: 101–138.
- Wang, C.T., Ji, B.P., Li, B., Nout, R., Li, P.L., Ji, H., and L.F. Chen. 2006. Purification and characterization of a fibrinolytic enzyme of *Bacillus subtilis* DC33, isolated from Chinese traditional *Douchi*. *Industrial Microbiology and Biotechnology* 33: 750–758.
- Watanabe, K., Fujimoto, J., Sasamoto, M., Dugersuren, J., Tumursuh, T., and Demberel, S. 2008. Diversity of lactic acid bacteria and yeasts in *airag* and *tarag*, traditional fermented milk products from Mongolia. *World Journal of Microbiology and Biotechnology* 24: 1313–1325.
- Watanabe, K., Fujimoto, J., Tomii, Y., Sasamoto, M., Makino, H., Kudo, Y., and Okada, S. 2009a. *Lactobacillus kisonensis* sp. nov., *Lactobacillus otakiensis* sp. nov., *Lactobacillus rapi* sp. nov. and *Lactobacillus sunkii* sp. nov., heterofermentative species isolated from sunki, a traditional Japanese pickle. *International Journal of Systematic and Evolutionary Microbiology* 59: 754–760.
- Watanabe, K., Makino, H., Sasamoto, M., Kudo, Y., Fujimoto, J., and S. Demberel. 2009b. *Bifidobacterium mongoliense* sp. nov., from *airag*, a traditional fermented mare's milk product from Mongolia. *International Journal of Systematic and Evolutionary Microbiology* 59: 1535–1540.
- Weckx, S., Allemeersch, J., Van der Meulen, R., Vrancken, G., Huys, G., Vandamme, P., Van Hummelen, P., and L. de Vuyst. 2009. Development and validation of a species-independent functional gene microarray that targets lactic acid bacteria. *Applied Environmental Microbiology* 75: 6488–6495.
- Weckx, S., Van der Meulen, R., Maes, D., Scheirlinck, I., Huys, G., Vandamme, P., and L. De Vuyst. 2010. Lactic acid bacteria community dynamics and metabolite production of rye sourdough fermentations share characteristics of wheat and spelt sourdough fermentations. *Food Microbiology* 27: 1000–1008.
- Wei, D. and S. Jong. 1983. Chinese rice pudding fermentation: Fungal flora of starter cultures and biochemical changes during fermentation. *Journal of Fermentation Technology* 61: 573–579.
- Wilson, H. 1999. Origins of viticulture. In: *The Oxford Companion to Wine*, 2nd edition, ed. J. Robinson, 505–506. Oxford: Oxford University Press.
- Winarno, F.G., Fardiaz, S., and Daulay, D. 1973. Indonesian Fermented Foods, Department of Agricultural Product Technology, Fatema, Bogor Agricultural University, Indonesia.



- Winarno, F.G., Fardiaz, S., and Daulay, D. 1973. Indonesian Fermented Foods, Department of Agricultural Product Technology, Fatema, Bogor Agricultural University, Indonesia.
- Wong, P.P. and H. Jakson. 1977. Malaysian belechan. In: *The Proceedings of the Symposium on Indigenous Fermented Foods*, November 21–27, 1977, GIAM-V, Bangkok.
- Wongkhalaung, C. 2004. Industrialization of Thai fish sauce (nam pla) In: *Industrialization of Indigenous Fermented Foods*. 2nd edition, Revised and Expanded, ed. K. H. Steinkraus. New York: Marcell Dekker, Inc.
- Wu, Y.C., Kimura, B., and T. Fujii. 2000. Comparison of three culture methods for the identification of *Micrococcus* and *Staphylococcus* in fermented squid shiokara. *Fisheries Science* 66: 142–146.
- Wu, R., Wang, L., Wang, J., Li, H., Menghe, B., Wu, J., Guo, M., and H. Zhang. 2009. Isolation and preliminary probiotic selection of lactobacilli from Koumiss in Inner Mongolia. *Journal of Basic Microbiology* 49: 318–326.
- Yan, Y., Qian, Y., Ji, F., Chen, J., and B. Han. 2013. Microbial composition during Chinese soy sauce *koji*-making based on culture dependent and independent methods. *Food Microbiology* 34(1): 189–195.
- Yan, P.M., Xue, W.T., Tan, S.S., Zhang, H., and X.H. Chang. 2008. Effect of inoculating lactic acid bacteria starter cultures on the nitrite concentration of fermenting Chinese paocai. *Food Control* 19: 50–55.
- Yegna Narayan Aiyar, A.K. 1953. Dairying in ancient India. *Indian Dairyman* 5: 77–83.
- Ying, A., Yoshikazu, A., and Yasuki, O. 2004. Classification of lactic acid bacteria isolated from chigee and mare milk collected in Inner Mongolia. *Animal Science Journal* 75: 245–252.
- Yokotsuka, T. 1985. Fermented protein foods in the Orient, with emphasis on shoyu and miso in Japan. In: *Microbiology of Fermented Foods*. Vol 1, ed. B.B. Wood, 197–247. London: Elsevier Applied Sciences.
- Yonzan, H. and J.P. Tamang. 2009. Traditional processing of *Selroti*—a cereal-based ethnic fermented food of the *Nepalis*. *Indian J Traditional Knowledge* 8(1): 110–114.
- Yonzan, H. and J.P. Tamang. 2010. Microbiology and nutritional value of *selroti*, an ethnic fermented cereal food of the Himalayas. *Food Biotechnology* 2: 227–247.
- Yonzan, H. and J.P. Tamang. 2013. Optimization of traditional processing of *Selroti*, a popular cereal-based fermented food. *Journal of Scientific and Industrial Research* 72: 43–47.
- Yoon, M.Y., Kim, Y.J., and H.J. Hwang. 2008. Properties and safety aspects of *Enterococcus faecium* strains isolated from *Chungkukjang*, a fermented soy product. *LWT-Food Science Technology* 41: 925–933.
- Yousif, N.M.K., Huch, M., Schuster, T., Cho, G.S., Dirar, H.A. Holzappel, W.H., and C.M.A.P. Franz. 2010. Diversity of lactic acid bacteria from Hussuwa, a traditional African fermented sorghum food. *Food Microbiology* 27: 757–768.
- Yu, J., Wang, W.H., Menghe, B.L., Jiri, M.T., Wang, H.M., Liu, W.J., Bao, Q.H. et al. 2011. Diversity of lactic acid bacteria associated with traditional fermented dairy products in Mongolia. *Journal of Dairy Science* 94: 3229–3241.
- Zhang, S. and Y. Liu. 2000. *Prepare Technology of Seasoning in China*. Beijing: South China University of Technology Press (in Chinese).
- Zhang, J.H., Tatsumi, E., Fan, J.F., and Li, L.T. 2007. Chemical components of *Aspergillus*-type Douchi, a Chinese traditional fermented soybean product, change during the fermentation process. *International Journal of Food Science and Technology* 42: 263–268.
- Zhu, Y.P., Cheng, Y.Q., Wang, L.J., Fan, J.F., and Li, L.T., 2008. Enhanced antioxidative activity of Chinese traditionally fermented Okara (Meitauza) prepared with various microorganism. *International Journal of Food Properties* 11: 519–529.
- Zhu, Y. and J. Trampe. 2013. Koji – where East meets West in fermentation. *Biotechnology Advances* 31(8): 1448–1457.
- Zhu, L., Xu, H., Zhang, Y., Fu, G., Wu, P.Q., and Y. Li. 2014. BOX-PCR and PCR-DGGE analysis for bacterial diversity of a naturally fermented functional food (enzymes). *Food Bioscience* 5: 115–122.

